

TW1  
07

# The Ore Bin



**LIBRARY**  
Marine Science Laboratory  
Oregon State University

DEC 29 1975

Vol. 37, No. 12  
December 1975

**STATE OF OREGON**  
**DEPARTMENT OF GEOLOGY AND MINERAL INDUSTRIES**

# The Ore Bin

Published Monthly By

STATE OF OREGON

DEPARTMENT OF GEOLOGY AND MINERAL INDUSTRIES

Head Office: 1069 State Office Bldg., Portland, Oregon - 97201

Telephone: [503] - 229-5580

## FIELD OFFICES

2033 First Street  
Baker 97814

521 N. E. "E" Street  
Grants Pass 97526

X X X X X X X X X X X X X X X X X X X

## Subscription Rate

1 year - \$3.00; 3 years - \$8.00

Available back issues - \$.25 at counter; \$.35 mailed

Second class postage paid at Portland, Oregon

X X X X X X X X X X X X X X X X X X X

## GOVERNING BOARD

R. W. deWeese, Portland, Chairman

Leeanne MacColl, Portland

H. Lyle Van Gordon, Grants Pass

## STATE GEOLOGIST

R. E. Corcoran

## GEOLOGISTS IN CHARGE OF FIELD OFFICES

Howard C. Brooks, Baker

Len Ramp, Grants Pass

X X X X X X X X X X X X X X X X X X X

Permission is granted to reprint information contained herein.

Credit given the State of Oregon Department of Geology and Mineral Industries  
for compiling this information will be appreciated.

## THE WALLOWA "ICE CAP" OF NORTHEASTERN OREGON an exercise in the interpretation of glacial landforms

John E. Allen  
Emeritus Professor of Geology, Portland State University

### Introduction

The geology of the Wallowa Mountains was first mapped in 1938 (Ross) and 1941 (Smith and Allen). Since that time, little work has been done on the effects of the Pleistocene ice which mantled much of the Wallowa high country more than 15,000 years ago. From elevations above 8,000 feet, the ice flowed down the canyons, reaching as low as 3,000 feet on Pine Creek. It has been the custom to call this glacial cover an "ice cap," implying that a more or less continuous sheet of névé covered the central area of the mountains. Only one map showing the inferred maximum extent of the ice has been published (Crandell, 1965). The accompanying map (Figure 1) is a demonstration of what can tentatively be deduced from a geomorphic study of topography on maps which were not available in 1941. Undoubtedly field checks will revise the map in detail, but the large picture should remain valid.

### Procedures

Seven 15-minute and four 7½-minute U.S. Geological Survey topographic quadrangle maps (see list at end of text) were used to determine the extent of the ice (Figure 1). These maps reveal many glacial landforms, both erosional and depositional (see Figure 2); the following can be easily recognized (definitions are adapted from Gary and others, 1972):

#### Erosional landforms (see accompanying photographs)

arête - a narrow, jagged, serrate mountain crest, or a narrow, rocky, sharp-edged ridge or spur, commonly present above the snow line in glaciated mountains

bastion - a prominent mass of bedrock extending from the mouth of a glacial trough and projecting far out into the glacial valley



Above: Cirque and tarn; terminal moraine from "little ice age" about 4,000 years ago. Eagle Cap on skyline and arête to right. (Oregon Hwy. Div. photo)

Below: Ice Lake dammed by recessional moraine. Matterhorn Peak (upper left) and Sacajawea (upper right) are highest peaks in Wallawas. (Oregon Hwy. Div. photo)





cirque - a deep, steep-walled, flat- or gently-floored, half-bowl-like recess or hollow commonly at the head of a glacial valley or saddle-shaped depression in a ridge

col - a deep pass formed by the headward erosion of two cirques

hanging valley - a glacial valley whose mouth is at a relatively high level on the steep side of a larger glacial valley

matterhorn - a high peak with prominent faces bounded by intersecting walls of three or more cirques

tarn - a relatively small, steep-banked lake or pool occupying an ice-gouged rock basin

U-shaped valley - a valley having a pronounced parabolic cross profile suggesting the form of a broad letter U, with steep parallel walls and a broad, nearly flat floor; a glacial trough

### Depositional landforms (see accompanying photographs)

moraine - a mound, ridge, or other distinct accumulation of unsorted, unstratified glacial drift, predominantly till, deposited chiefly by direct action of glacier ice in a variety of topographic landforms that are independent of control by the surface on which the drift lies

lateral moraine - a low, ridge-like moraine deposited at or near the side margin of a mountain glacier

recessional moraine - an end moraine built during a temporary but significant halt or pause in the final retreat of a glacier

terminal moraine - an end moraine, extending across a glacial valley as an arcuate or crescentic ridge, that marks the farthest advance or maximum extent of a glacier


morainal lake - a glacial lake occupying a depression and dammed by a terminal or recessional moraine

outwash - stratified detritus removed or "washed out" from a glacier by melt-water streams and deposited in front of or beyond the terminal moraine

In studying the topographic maps, the elevation of lateral moraines was particularly helpful in determining the extent of the ice in the valleys. Evidences of glacial extent, recognizable in the field but not on topographic maps, were noted by Smith and Allen (1941). These included the location of glacial erratics and the elevations of ice-scoured and polished surfaces (see photographs).

Interpolation of projected thicknesses of the ice, carefully checked by remnants of lateral moraines and medial moraine spurs at tributary valley mouths, permitted drawing appropriately spaced contour lines for the surface of the ice (Figure 1).

Below the firn area (zone of accretion), the thickness of the ice is inferred to have decreased gradually (in the zone of ablation) to the terminus. Rock steps and steeper stretches in the present valley profile were taken into



Above: U-shaped valley of Eagle Creek near Boulder Park.  
(Oregon Hwy. Div. photo)

Below: Bastions project into U-shaped valley of West Fork of  
Wallowa River. Small cirques and hanging valley on left.  
Sentinel Peak in distance is a "matterhorn."



consideration in estimating the thickness of the ice. Glacial cirque ice is mapped as being halfway up the cirque walls.

In a few instances, landslides may have been interpreted as *névé* fields or incipient cirques at lower elevations on the periphery of the area, and bastions may have been interpreted as moraines at glacial junctions.

### Conclusions

The so-called "ice cap" which fed the radiating glacial streams during the Pleistocene turns out to be multiple in nature. The Lake Basin ice field had a surface at least 8,500 feet above sea level; over the ridge to the west the Minam Lake ice field was a few hundred feet lower. The Aneroid Lake ice field and the several Imnaha ice fields were at about 8,400 feet elevation. Nunataks (isolated rock knobs projecting above the ice) are surprisingly rare; most of the high peaks appear to have been connected by narrow unglaciated ridges.

Thin ice fields covered much of the plateau surface south of the Imnaha valley, feeding glaciers down Lake and Clear Creeks. This area may have been more extensively covered by ice than is shown (Ross, 1938, p. 58).

The nine large glaciers (each more than 10 miles long) and their tributaries covered about 279 square miles (see Figure 1). Other isolated ice fields and glaciers added 58 square miles for a total of about 337 square miles covered by ice during the Pleistocene. This is a closer approximation than "200 square miles" or "500 square miles" - figures which have been bandied about for years - but it is a minimum figure, since thickness of the ice was conservatively estimated where definite evidence was lacking on the topographic maps.

In length, the Lostine glacier (22 miles) was slightly longer than the Minam (21 miles) and the Imnaha (20 miles). The other six major glaciers were from 12 to 13 miles in length.

In area covered, the Minam glacier was the largest, with 67 square miles, followed by the Lostine (55) and the Imnaha (50). The Wallowa glacier covered 35 square miles. The other five major glaciers covered between 11 and 18 square miles each.

In elevation reached by the lower ends of the glaciers, the Lostine came down to 3,380 feet above sea level; the Minam was a close second at 3,600 feet. The six other major glaciers reached between 4,000 and 4,200 feet. The Pine Creek glacier apparently reached to 2,960 feet at some time (P. Frazier) during the Pleistocene (Crandell, 1965).


In thickness of ice, without field-checking, the maximum seems to have been about 2,500 feet in the upper Lostine and Minam glaciers. The Wallowa was thickest at its junction with the East Fork, about 1,500 feet. The Hurricane was less than 1,000 feet thick throughout.



Above: Moccasin Lake, a tarn lake in upper Wallowa River drainage. Eagle Cap on left.

Below: Glacier Lake, a tarn at head of West Fork of Wallowa River. Cusick Mountain in upper right. Three cols on skyline. (Oregon Hwy. Div. photo)





Above: Terminal moraine of Wallowa River glacier dammed river to form Wallowa Lake. (Oregon Hwy. Div. photo)

Below: Lateral and terminal moraines hem in Wallowa Lake. U-shaped valley of West Fork of Wallowa River in upper left and many cirque basins on skyline. (U.S. Forest Service photo)



FIGURE 1

# EXTENT of PLEISTOCENE GLACIERS

in the

## WALLOWA MOUNTAINS, OREGON

SUMMARY OF DATA ON WALLOWA GLACIERS

Name	Length (miles)	No. of major branches	Elevation at lower end (feet)	Area of ice (sq. miles)
A. Glaciers more than 10 miles long:				
Lostine River	22	4	3,380	55.0
Minam River	21	5	3,600	67.5
Imnaha River	20	5	4,200	50.5
Wallowa River	13	2	4,200	35.4
Hurricane Creek	13	1	4,200	11.5
East Eagle Creek	13	1	4,100	12.2
Eagle Creek	13	1	4,000	18.5
Bear Creek	12	2	4,200	15.5
Pine Creek	12.6	2	2,690	12.6
				<u>278.7</u>

### B. Between 5 and 9 miles long (tributary or isolated):

Sheep Creek	9.0	3	5,400	11.1
Lake Fork Creek	8.0	1	5,200	10.2
North Fork Minam	8.0	2		(9.4)*
Goot Creek (Bear)	6.5	2		(6.5)*
McCully Creek	6.0	3	5,200	4.7
East Fork Wallowa	5.5	1		(4.7)*
West Eagle Creek	5.5	3		(6.0)*
				<u>26.0</u>

### C. Between 2 and 5 miles long:

Catherine Creek	4.6		5,300	2.6
Scotch Creek	3.5		6,400	2.0
Boulder Creek	3.2		5,200	2.5
S. Fork Catherine Creek	3.2		5,600	1.8
Little Sheep Creek	2.9		5,900	.9
Clear Creek	2.6 (Lake Fork)			(1.0)*
Duck Creek	2.5 (Imnaha)			(2.3)*
Sturgill Creek	2.3		6,000	1.0
				<u>10.8</u>


### D. 86 isolated glaciers or neve' fields less than 2 miles long; if they averaged $\frac{1}{2}$ mile in area:

				21.5
Total area:				337.0

\* Included in area of glaciers listed under "A".









USGS QUADRANGLE SHEETS : JOSEPH







SCALE  
0 1 MILE  
EXCEPT AS NOTED

— PROX. MAX. BOUNDARY of GLACIER

— MORAINES

FIGURE 2





The Frazier glaciation, the latest major advance of alpine glaciers in the Wallowas, ended about 15,000 years ago. Since that time, climatic fluctuations have caused small glacial advances, such as the "little ice age" about 4,000 years ago, which left terminal moraines high in the major glacial valleys.



Moraine on Thorp Creek, tributary of Hurricane Creek, is a prominent topographic feature. "Ballet dancer" is Warren D. Smith, noted Oregon geologist, deceased.



Outwash from Wallowa glaciers formed broad plains of sand and gravel at foot of mountains near Joseph. (Oregon Hwy. Div. photo)



Above: Striated, ice-scoured rock surface exposed in Lostine Canyon. Note geologic pick for scale.

Below: Erratic boulders of granodiorite perched on limestone bedrock near Marble Point, 2,500 feet above Lostine River, indicate elevation of glacial ice.



## References

- Crandell, D. R., 1965, The glacial history of western Washington and Oregon, in Wright, H. E., Jr., and Fry, D. G. (eds.), *The Quaternary of the United States*: Princeton Univ. Press, p. 349-350 (map after Lowell, 1940).
- Gary, M., McAfee, R., Jr., and Wolf, C. L. (eds.), 1972, *Glossary of Geology*: Am. Geol. Inst.
- Lowell, W. R., 1940, *Glaciation in the Wallowa Mountains of Oregon*: Univ. of Chicago, master's thesis, unpub.
- Ross, C. P., 1938, The geology of a part of the Wallowa Mountains: Ore. Dept. Geol. and Mineral Indus. Bull. 3, 74 p.
- Smith, W. D., Allen, J. E., 1941, Geology and physiography of the northern Wallowa Mountains, Oregon: Ore. Dept. Geol. and Mineral Indus. Bull. 12, 64 p.
- Stovall, J. C., 1929, Pleistocene geology and physiography of the Wallowa Mountains, with special reference to Wallowa and Hurricane Canyons: Univ. Oregon master's thesis, unpub.

## Topographic Quadrangle Maps

<u>15-minute</u>		<u>7½-minute</u>
Enterprise	Homestead	Fox Point
Joseph	Sparta	Jim White Ridge
Eagle Cap	Halfway	China Cap
Cornucopia		Flagstaff Butte

\* \* \* \* \*

## MOUNT RAINIER HISTORY DESCRIBED

"Quaternary Stratigraphy and Extent of Glaciation in the Mount Rainier Region, Washington," by D. R. Crandell and R. D. Miller, is a recent publication by the U.S. Geological Survey and designated Professional Paper 847. The 59-page report includes a geologic map of the region and a topographic map of Mount Rainier National Park. According to the authors, a nearly continuous ice cap once mantled the Cascade Range, with ice extending down the Cowlitz River about 120 km (193 miles). Today each of the five major river valleys contains glacial deposits of repeated glaciations, and drift from at least two ancient glaciations has been recognized.

Professional Paper 847 is for sale by U.S.G.S. Branch of Distribution, 1200 S. Eads St., Arlington, VA 22202. The price is \$1.60.

\* \* \* \* \*



## POTENTIAL GEOTHERMAL LAND WITHDRAWN FROM MINERAL EXPLORATION

Recent news release from the U.S. Department of the Interior, Bureau of Land Management, announces that 31,114.7 acres (approximately 50 square miles) of land in Malheur County centering around Jordan Craters has been established as a "Research Natural Area." According to the news release, the arrangement will "aid BLM in managing and protecting the area to insure preservation of the total environment." Not stated in the release was the fact that this action also withdraws the land from mineral entry.

As noted in the October 1975 ORE BIN, about two-thirds of all public lands are now completely or partially withdrawn from mining activities. Here is another instance of a tract of land essentially reserved for ecological study, a single-use purpose which is contrary to the multiple-use concept espoused for many years by the Federal government.

The Jordan Craters lie within a large area of relatively young volcanic rocks, one of the most promising regions for future geothermal development. Even though the Jordan Craters Research Natural Area of approximately 31,000 acres is only a small part of the total volcanic field, its withdrawal has a negative effect on mineral exploration activities in the surrounding territory; past experience has shown that companies looking for geothermal mineral resources always give a wide berth to public lands withdrawn "to insure preservation of the total environment."

Every public land withdrawal is justified in the eyes of the Federal agency responsible for managing it, and in most instances, the area involved for any one withdrawal is comparatively small. But as has been clearly pointed out by Bennethum and Lee in the October 1975 ORE BIN, the cumulative effect of withdrawals is tremendous. To quote from these authors: "We think some attention will have to be paid to the trend toward accelerated withdrawals because it seriously erodes the long-range mineral position of the country. It affects our economy, our ability to protect jobs, and it is forcing American industry to look elsewhere for minerals. It makes us vulnerable to mineral cartels like the OPEC oil cartel."

--- R. E. Corcoran

\* \* \* \* \*

## INTERIOR DEPARTMENT POSTS FILLED

Thomas S. Kleppe, former U.S. Representative from North Dakota, was made Secretary of the Interior on October 9, 1975, succeeding Stanley K. Hathaway, who resigned in July. On November 20, 1975, the Senate confirmed the nomination of D. Kent Frizzell to be Under-Secretary of the Interior. Frizzell, a Kansan who has been solicitor of the Interior Department, fills a post that has been vacant since May 1.

\* \* \* \* \*

\*\*\*\*\*

# ARE YOU EXPIRING?

We Don't Want to Lose YOU!

CHECK BACK COVER OF OCTOBER ISSUE  
IS DEC. 1975 MARKED? THEN THIS IS YOUR LAST COPY!

\*\*\*\*\*

Please use renewal form on centerfold of October issue --  
or cut out form below and mail to:

ORE BIN Subscriptions  
Oregon Dept. of Geology and Mineral Indus.  
1069 State Office Building  
Portland, Oregon 97201

ORE BIN RENEWAL FORM  
\$3 - year; \$8 - 3 years

Enclosed is \$ \_\_\_\_\_ to extend my ORE BIN subscription for  
\_\_\_\_\_ years:

Name \_\_\_\_\_

Street \_\_\_\_\_

City, State \_\_\_\_\_ Zip \_\_\_\_\_

# AVAILABLE PUBLICATIONS

(Please include remittance with order; postage free. All sales are final - no returns. Upon request, a complete list of Department publications, including out-of-print, will be mailed.)

## BULLETINS

26. Soil: Its origin, destruction, preservation, 1944: Twenhofel . . . . .	\$0.45
33. Bibliography (1st suppl.) geology and mineral resources of Oregon, 1947: Allen . . . . .	1.00
35. Geology of Dallas and Valsetz quadrangles, Oregon, rev. 1964: Baldwin . . . . .	3.00
36. Papers on Tertiary foraminifera: Cushman, Stewart & Stewart. . . . .	vol. 2-1.25
39. Geology and mineralization of Morning mine region, 1948: Allen and Thayer . . . . .	1.00
44. Bibliography (2nd suppl.) geology and mineral resources of Oregon, 1953: Steere. . . . .	1.00
46. Ferruginous bauxite deposits, Salem Hills, 1956: Corcoran and Libbey . . . . .	1.25
49. Lode mines, Granite mining district, Grant County, Oregon, 1959: Koch . . . . .	1.00
52. Chromite in southwestern Oregon, 1961: Ramp . . . . .	5.00
53. Bibliography (3rd suppl.) geology and mineral resources of Oregon, 1962: Steere, Owen . . . . .	3.00
57. Lunar Geological Field Conf. guidebook, 1965: Peterson and Groh, editors . . . . .	3.50
60. Engineering geology of Tualatin Valley region, 1967: Schlicker and Deacon . . . . .	7.50
61. Gold and silver in Oregon, 1968: Brooks and Ramp . . . . .	7.50
62. Andesite Conference Guidebook, 1968: Dole . . . . .	3.50
64. Geology, mineral, and water resources of Oregon, 1969 . . . . .	3.00
65. Proceedings of the Andesite Conference, 1969: McBirney, editor (photocopy) . . . . .	10.00
66. Geology and mineral resources of Klamath and Lake Counties, 1970. . . . .	6.50
67. Bibliography (4th suppl.) geology and mineral industries, 1970: Roberts . . . . .	3.00
68. Seventeenth biennial report of the Department, 1968-1970 . . . . .	1.00
69. Geology of the southwestern Oregon Coast, 1971: Dott . . . . .	4.00
70. Geologic formations of western Oregon, 1971: Beaulieu . . . . .	2.00
71. Geology of selected lava tubes in the Bend area, 1971: Greeley . . . . .	2.50
72. Geology of Mitchell quadrangle, Wheeler County, 1972: Oles and Enlows . . . . .	3.00
73. Geologic formations of eastern Oregon, 1972: Beaulieu . . . . .	2.00
75. Geology, mineral resources of Douglas County, 1972: Ramp . . . . .	3.00
76. Eighteenth biennial report of the Department, 1970-1972. . . . .	1.00
77. Geologic field trips in northern Oregon and southern Washington, 1973. . . . .	5.00
78. Bibliography (5th suppl.) geology and mineral industries, 1973: Roberts and others . . . . .	3.00
79. Environmental geology inland Tillamook Clatsop Counties, 1973: Beaulieu. . . . .	7.00
80. Geology and mineral resources of Coos County, 1973: Baldwin and others . . . . .	6.00
81. Environmental geology of Lincoln County, 1973: Schlicker and others . . . . .	9.00
82. Geol. Hazards of Bull Run Watershed, Mult. Clackamas Counties, 1974: Beaulieu . . . . .	6.50
83. Eocene stratigraphy of southwestern Oregon, 1974: Baldwin . . . . .	4.00
84. Environmental geology of western Linn Co., 1974: Beaulieu and others. . . . .	12.00
85. Environmental geology of coastal Lane Co., 1974: Schlicker and others . . . . .	12.00
86. Nineteenth biennial report of the Department, 1972-1974 . . . . .	1.00
87. Environmental geology of western Coos and Douglas Counties, Oregon, 1975 . . . . .	in press
88. Geology and mineral resources of upper Chetco River drainage, 1975: Ramp . . . . .	in press

## GEOLOGIC MAPS

Geologic map of Oregon west of 121st meridian, 1961: Wells and Peck . . . . .	\$2.00; mailed - 2.50
Geologic map of Oregon (12" x 9"), 1969: Walker and King . . . . .	0.25
Geologic map of Albany quadrangle, Oregon, 1953: Allison (from Bulletin 37) . . . . .	1.00
Geologic map of Galice quadrangle, Oregon, 1953: Wells and Walker . . . . .	1.50
Geologic map of Lebanon quadrangle, Oregon, 1956: Allison and Felts . . . . .	1.50
Geologic map of Bend quadrangle, and portion of High Cascade Mtns., 1957: Williams . . . . .	1.50
GMS-1: Geologic map of the Sparta quadrangle, Oregon, 1962: Prostka . . . . .	2.00
GMS-2: Geologic map, Mitchell Butte quadrangle, Oregon: 1962 . . . . .	2.00
GMS-3: Preliminary geologic map, Durkee quadrangle, Oregon, 1967: Prostka . . . . .	2.00
GMS-4: Gravity maps, Oregon onshore & offshore; [set only]: at counter \$3.00, mailed . . . . .	3.50
GMS-5: Geology of the Powers quadrangle, 1971: Baldwin and Hess . . . . .	2.00
GMS-6: Preliminary report, geology of part of Snake River Canyon, 1974: Vallier. . . . .	6.50


[Continued on back cover]

Marine Science Center  
Marine Science Drive  
Newport, Oregon 97365

The ORE BIN  
1069 State Office Bldg., Portland, Oregon 97201

## The Ore Bin

Second Class Matter  
POSTMASTER: Form 3579 requested



### Available Publications, Continued:

#### SHORT PAPERS

- |   |        |
|---|--------|
| 18. Radioactive minerals prospectors should know, 1955: White and Schafer . . . . . | \$0.30 |
| 19. Brick and tile industry in Oregon, 1949: Allen and Mason . . . . .              | 0.20   |
| 21. Lightweight aggregate industry in Oregon, 1951: Mason . . . . .                 | 0.25   |
| 24. The Almeda mine, Josephine County, Oregon, 1967: Libbey . . . . .               | 3.00   |

#### MISCELLANEOUS PAPERS

- |   |      |
|---|------|
| 1. Description of some Oregon rocks and minerals, 1950: Dole . . . . .                      | 1.00 |
| 2. Oregon mineral deposits map (22 x 34 inches) and key (reprinted 1973): . . . . .         | 1.00 |
| 4. Rules and regulations for conservation of oil and natural gas (rev. 1962) . . . . .      | 1.00 |
| 5. Oregon's gold placers (reprints), 1954 . . . . .   | 0.50 |
| 6. Oil and gas exploration in Oregon, rev. 1965: Stewart and Newton . . . . .               | 3.00 |
| 7. Bibliography of theses on Oregon geology, 1959: Schlicker . . . . .                      | 0.50 |
| (Supplement) Bibliography of theses, 1959 to Dec. 31, 1965: Roberts . . . . .               | 0.50 |
| 8. Available well records of oil and gas exploration in Oregon, rev. 1963: Newton . . . . . | 1.00 |
| 11. A collection of articles on meteorites, 1968 (reprints from The ORE BIN) . . . . .      | 1.50 |
| 12. Index to published geologic mapping in Oregon, 1968: Corcoran . . . . .                 | 0.50 |
| 13. Index to The ORE BIN, 1950-1974. . . . .  | 1.50 |
| 14. Thermal springs and wells, 1970: Bowen and Peterson . . . . .                           | 1.50 |
| 15. Quicksilver deposits in Oregon, 1971: Brooks . . . . .                                  | 1.50 |
| 16. Mosaic of Oregon from ERTS-1 imagery, 1973: . . . . .                                   | 2.50 |
| 18. Proceedings of Citizens' Forum on potential future sources of energy, 1975 . . . . .    | 2.00 |

#### OIL AND GAS INVESTIGATIONS

- |  |      |
|--|------|
| 1. Petroleum geology, western Snake River basin, 1963: Newton and Corcoran . . . . .         | 3.50 |
| 2. Subsurface geology, lower Columbia and Willamette basins, 1969: Newton . . . . .          | 3.50 |
| 3. Prelim. identifications of foraminifera, General Petroleum Long Bell No. 1 well . . . . . | 2.00 |
| 4. Prelim. identifications of foraminifera, E. M. Warren Coos Co. 1-7 well: Rau . . . . .    | 2.00 |

#### MISCELLANEOUS PUBLICATIONS

- |   |                                       |
|---|---------------------------------------|
| Landforms of Oregon: a physiographic sketch (17" x 22"), 1941 . . . . . | 0.25                                  |
| Mining claims (State laws governing quartz and placer claims) . . . . . | 0.50                                  |
| Oregon base map (22" x 30"). . . . .                                    | 0.50                                  |
| Geologic time chart for Oregon, 1961 . . . . .                          | free                                  |
| Postcard - geology of Oregon, in color . . . . .                        | 10¢ each; 3 - 25¢; 7 - 50¢; 15 - 1.00 |
| The ORE BIN - Annual subscription . . . . . (\$8.00 for 3 yrs.)         | 3.00                                  |
| Available back issues, each . . . . .                                   | 25¢; mailed 0.35                      |
| Accumulated index - see Misc. Paper 13                                  |                                       |