


7/15/60

B6f  
Pam

# FOREST STATISTICS *for* WALLOWA COUNTY, OREGON



PACIFIC NORTHWEST  
FOREST AND RANGE EXPERIMENT STATION  
U. S. DEPT. OF AGRICULTURE • FOREST SERVICE

PREPARED BY  
THE DIVISION OF FOREST ECONOMICS RESEARCH

Carl A. Newport, Chief

Field and Office Work in

Wallowa County, Oregon, by:

C. E. Mayer, In Charge  
Ben Spada, Field Supervisor

James T. Bones  
Peter H. Geiger  
John W. Hazard  
Charles H. Harden  
Earl Nelson  
John P. Kennedy  
Hilton D. Muntz

Kathryn Flaherty  
Inga E. Fulkerson  
Marlin E. Plank  
James E. Simonson  
Thomas J. Spolar  
Paul E. Hightree  
Wayne E. Orr

Richard D. Schimel

Acknowledgment is made of cooperation from the staff of the Wallowa-Whitman and Umatilla National Forests who participated in the measurement of field data, and from public and private agencies in furnishing cutting and ownership records.

Forest Survey Report 134

FOREST STATISTICS  
FOR  
WALLOWA COUNTY, OREGON

by

James T. Bones  
and  
Richard D. Schimel

April 1960

PACIFIC NORTHWEST  
FOREST AND RANGE EXPERIMENT STATION  
R. W. Cowlin, Director                      Portland, Oregon  
FOREST SERVICE                      U.S. DEPARTMENT OF AGRICULTURE



## PREFACE

This publication summarizes the results of a 1957 reinventory of the forests of Wallowa County, Oreg. The reinventory is a part of the maintenance phase of the Forest Survey, a nationwide project of the Forest Service authorized by the McSweeney-McNary Forest Research Act of 1928, amended June 25, 1949. The purpose of the Forest Survey is to periodically inventory the extent and condition of forest lands and the timber and other products on them to ascertain rates of forest growth and depletion, to estimate present consumption of timber products and determine probable future trends in timber requirements, to analyze and make available survey information needed in the formulation of forest policies and programs, and to make resurveys as necessary to keep the basic information up to date.

The Forest Survey is conducted in the various forest regions of the Nation by the regional experiment stations of the Forest Service. In the Pacific Northwest region of Oregon and Washington, it is an activity of the Pacific Northwest Forest and Range Experiment Station at Portland, Oreg.

The initial inventory of forest lands in Wallowa County took place in 1935 and 1936. A statistical report, "Forest Statistics for Wallowa County, Oregon," and a forest type map on a 1-inch-to-the-mile scale were released in 1937.

The forest type maps have been revised as a result of the 1957 reinventory and are available on scales of 1 and 2 inches to the mile. A single 1-inch-scale map was prepared for the county; 2-inch-scale maps were prepared in sections (see index map, inside rear cover). Prints of these maps are available at cost of blue-printing. For information write Director, Pacific Northwest Forest and Range Experiment Station, P. O. Box 4059, Portland 8, Oreg.

## CONTENTS

	<u>Page</u>
Preface	
Map: Generalized Forest Type Map, Wallowa County, Oreg., 1957	
DESCRIPTION OF THE COUNTY . . . . .	1
SIGNIFICANT FINDINGS IN THE FOREST INVENTORY . . . . .	2
Land Classification . . . . .	2
Commercial Forest Land Area . . . . .	2
Major Types . . . . .	2
Stand-Size Classes . . . . .	3
Stocking of Young-Growth Stands . . . . .	3
Commercial Forest Land Timber Volumes . . . . .	3
Volume of Sawtimber by Species . . . . .	4
Volume of Sawtimber by Diameter Classes . . . . .	4
Volume of Growing Stock by Species . . . . .	4
Forest Ownership . . . . .	5
Commercial Forest Land . . . . .	5
Sawtimber Volume . . . . .	5
Log Production . . . . .	6
SUMMARY TABLES . . . . .	7
FOREST SURVEY PROCEDURE . . . . .	20
Initial Inventory . . . . .	20
Reinventory . . . . .	20
ACCURACY OF 1957 REINVENTORY DATA . . . . .	22
Forest Area . . . . .	22
Timber Volume . . . . .	22
DIFFERENCES IN RESULTS OF INVENTORIES . . . . .	22
Forest Area . . . . .	23
Timber Volume . . . . .	23

	<u>Page</u>
DEFINITION OF TERMS . . . . .	24
Land Area . . . . .	24
Forest Land Classes . . . . .	25
Types . . . . .	25
Tree Classes . . . . .	26
Stand-Size Classes . . . . .	27
Stocking . . . . .	28
Timber Volume . . . . .	28
Timber Cut . . . . .	29
TREE SPECIES . . . . .	30


#### List of Tables

1. Land area in Wallowa County, by major class of land, 1957 . . . . .	7
2. Area of commercial forest land in Wallowa County, by ownership and stand-size classes, 1957 . . . . .	8
3. Area of commercial forest land in Wallowa County, by major forest type and stand-size class, 1957 . . . . .	9
4. Land area in Wallowa County, by cover type, ownership class, and land-use class, 1957 . . . . .	10-11
5. Area of commercial forest land in Wallowa County, by forest-condition and ownership classes, 1957 . . . . .	12
6. Area of young-growth timber stands on commercial forest land in Wallowa County, by stand-size class, species group, and stocking class, 1957 . . . . .	13
7. Net volume of live sawtimber and growing stock on commercial forest land in Wallowa County, by ownership class, 1957 . . . . .	14
8. Net volume of live sawtimber and growing stock on commercial forest land in Wallowa County, by stand-size class, 1957 . . . . .	15
9. Net volume of live sawtimber and growing stock on commercial forest land in Wallowa County, by species, 1957 . . . . .	16
10. Net volume of live sawtimber on commercial forest land in Wallowa County, by diameter class and species group, 1957 . . . . .	17
11. Net volume of all timber on commercial forest land in Wallowa County, by class of material and species group, 1957 . . . . .	18
12. Average annual cut of live sawtimber and growing stock on commercial forest land in Wallowa County, by species group, 1953-1958 . . . . .	19
13. Comparison of forest area statistics for Wallowa County, initial inventory and reinventory . . . . .	23
14. Comparison of timber volume statistics for Wallowa County, initial inventory and reinventory . . . . .	24

# GENERALIZED FOREST TYPE MAP







WALLOWA COUNTY, OREGON

1957



SCALE: 5 0 5 10 15 20 MILES

## LEGEND:

	Douglas-fir		Fir-spruce
	Ponderosa pine		Noncommercial
	Lodgepole pine		Nonforest

## DESCRIPTION OF THE COUNTY

Wallowa County is located in the northeastern corner of the State of Oregon. It is bounded by the State of Washington on the north, the Snake River (Idaho boundary) on the east, Baker County on the south, and Union and Umatilla Counties on the west.

All of the county's major streams are tributaries of the Snake River. The Imnaha, Wallowa, and Minam Rivers head in the Wallowa Mountains, which form the county's southern boundary, and drain in a northerly direction. The Imnaha River empties directly into the Snake River while the Minam and Wallowa Rivers join the Grande Ronde River, which in turn empties into the Snake River.

The topography of Wallowa County is quite diverse. The Wallowa Valley, agricultural center of the county, is surrounded on all sides by prominent physical features. The granite peaks of the high Wallows to the south and southwest and the abrupt river canyons of the north and east encircle the valley. The steep river gorges, which limit access for a 20-mile width on the eastern side of the county, are terminated by the 5,600-foot-deep Hells Canyon of the Snake River.

Elevations vary from 1,244 feet on the Snake River to 10,033 feet at the summit of the Sacajawea Peak in the Wallowa Mountains. The total annual precipitation varies with elevation. In 1957, the precipitation in Wallowa County varied from 12 inches in the Wallowa Valley to 50 inches in the mountains.

Wallowa County has no Federal highways traversing it but is served by State Highway 3 from the north and State Highway 82 from the west. A Forest Service road is proposed to link Wallowa County with Baker County on the south. A branch line of the Union Pacific Railroad connects the Wallowa Valley with the main line at La Grande to the west.


In recent years the population of Wallowa County has been steadily decreasing and shifting from the farms to the cities and villages. In 1958, the population was 6,750, a decrease of 7.1 percent from the number reported in the 1950 census. Of the 6,750 total population, 4,110 live in towns and 2,640 are rural dwellers. The town of Enterprise is the county seat.



## SIGNIFICANT FINDINGS IN THE FOREST INVENTORY

### Land Classification

About 1,101,000 acres or 55 percent of Wallowa County is forest land, 80 percent of which is classified as commercial. About two-thirds of the noncommercial forest land is unproductive and includes the high-elevation subalpine type and midelevation noncommercial rocky areas. The remaining one-third consists of productive-reserved land--commercial forest land reserved from cutting--




Forty-five percent of the total land area in Wallowa County is non-forest land. According to the 1954 Census of Agriculture, 37 percent of Wallowa County is farmland, which accounts for 82 percent of the total non-forest area. The remainder is barren cliff areas adjacent to the Snake River and in the Wallowa Mountains.

### Commercial Forest Land Area

#### Major Types


Although ponderosa pine is the predominant tree species throughout the county, it rarely appears in extensive pure stands; on northerly and easterly aspects Douglas-fir, western larch, and lodgepole pine become associates. The higher elevations are dominated by the fir-spruce type and by pure stands of lodgepole pine where fire consumed the original stand.



Other types include white pine at high elevations, hardwoods in the valley stream bottoms, and nonstocked areas.


### Stand-Size Classes

Nearly 47 percent of the total sawtimber stands are in trees that are predominately 21.0 inches d.b.h. or larger. Most of the poletimber stands have resulted from cutting in the lower elevations or fire in the higher ones. Commercial forest land acreage of seedling and sapling stands and nonstocked areas is negligible.



### Stocking of Young-Growth Stands

Most of the young-growth stands of Wallowa County are satisfactorily stocked (40-100 percent); only 6 percent are poorly stocked or nonstocked. (Young-growth stands are those in which the majority of the cubic-foot volume is in trees less than 21.0 inches d.b.h.) Sixty-nine percent of the young-growth sawtimber, 60 percent of the poletimber, and 38 percent of the seedling and sapling areas are well stocked (over 70 percent).




### Commercial Forest Land Timber Volumes

The net volume of live sawtimber trees (11.0 inches d.b.h. and larger) on commercial forest land in Wallowa County is estimated to be 7,289 million board feet, Scribner rule. Ninety-eight percent is in sawtimber stands. The remaining volume is in sawtimber-size trees in poletimber, seedling and sapling, and nonstocked areas.

### Volume of Sawtimber by Species


Softwood species constitute more than 99 percent of the live sawtimber volume in Wallowa County. Sixty-two percent of this sawtimber volume is in ponderosa pine and Douglas-fir. The remaining volume is made up of several softwood and two hardwood species. "Other softwoods" include whitebark pine, lodgepole pine, subalpine fir, Engelmann spruce, and mountain hemlock; the hardwoods are black cottonwood and quaking aspen.



1/ Less than 0.5 percent.

### Volume of Sawtimber by Diameter Classes


Of the two dominant species, 51 percent of the Douglas-fir and 78 percent of the ponderosa pine sawtimber volume is in trees more than 20.9 inches d.b.h.



### Volume of Growing Stock by Species

Growing-stock volume is the net cubic volume of all live sound trees 5.0 inches d.b.h. and larger to a minimum 4-inch top inside bark.


Of the total growing stock in Wallowa County, 78 percent is in sawtimber trees and 22 percent is in poletimber trees. Ponderosa pine and Douglas-fir comprise over half of the county's growing stock.



### Forest Ownership


#### Commercial Forest Land

Nearly 66 percent of the commercial forest land in Wallowa County is federally owned, of which 99 percent is national forest. About 75 percent of the area of large sawtimber stands (trees more than 21.0 inches d.b.h.) and 58 percent of the area of small sawtimber stands (trees 11.0 to 20.9 inches d.b.h.) are on Federal lands. About 69 percent of the area of residual sawtimber stands is in private ownership.



#### Sawtimber Volume

Approximately 79 percent of the live sawtimber volume in Wallowa County is found on Federal lands. Private owners have 21 percent of the total volume on 34 percent of the total sawtimber area.





### Log Production

Annual log production (Scribner rule) for Wallowa County during the period 1935-58 was reported to be:

	<u>Annual production<sup>1/</sup></u> (M b. m.)
1935	16,205
1940	33,067
1945	34,480
1950	58,243
1955	79,906
1956	87,650
1957	99,635
1958	88,327

<sup>1/</sup> Does not include volume removed for poles, piling, or products from woodcutting operations.

A peak log production in Wallowa County of 115 million board feet was reached in 1924, followed by a low of 6 million feet in 1934. Annual log production since 1934 has been generally increasing.

To date most of the logs have been supplied by private lands. In 1958, 65 percent of the total annual log production was provided by these lands that contain only 21 percent of the total sawtimber volume of the county. Indications are that as the need for logs increases and timber values rise, the timber-harvesting program on Federal lands in Wallowa County will accelerate.

Table 1.--Land area in Wallowa County, by major class of land, 1957

Class of land	Area
	<u>Acres</u>
Forest:	
Commercial	878,110
Noncommercial:	
Productive-reserved	70,440
Unproductive	152,770
Total forest	1,101,320
Nonforest	910,510
All classes	2,011,830

Table 2.--Area of commercial forest land in Wallowa County,  
by ownership and stand-size classes, 1957

(In acres)

Ownership class	Total	Sawtimber stands	Poletimber stands	Seedling and sapling stands	Nonstocked areas
Private	296,200	236,720	54,740	4,480	260
State	4,420	3,260	1,130	30	--
County	270	270	--	--	--
<hr/>					
Federal:					
Bureau of Land Mgt.	4,690	4,430	220	40	--
National forest	572,530	469,890	94,810	660	7,170
Total Federal	577,220	474,320	95,030	700	7,170
All ownerships	878,110	714,570	150,900	5,210	7,430

Table 3.--Area of commercial forest land in Wallowa County,by major forest type and stand-size class, 1957

(In acres)

Forest type	Total	Sawtimber stands		Poletimber stands	Seedling and sapling stands	Nonstocked areas
		Large <sup>1/</sup>	Small <sup>2/</sup>			
Ponderosa pine	348,710	193,400	103,830	48,470	3,010	--
White pine <sup>3/</sup>	2,010	10	20	1,980	--	--
Lodgepole pine	58,680	--	19,900	38,380	400	--
Douglas-fir	246,410	82,240	122,900	40,770	500	--
Western larch	60,310	22,280	32,870	4,600	560	--
Fir—spruce	150,620	35,630	98,430	15,820	740	--
Hardwood	3,940	840	2,220	880	--	--
Nonstocked	7,430	--	--	--	--	7,430
Total	878,110	334,400	380,170	150,900	5,210	7,430

<sup>1/</sup> 21.0 inches d.b.h. and larger.<sup>2/</sup> 11.0-20.9 inches d.b.h.<sup>3/</sup> The white pine cover type, as specified by the Forest Survey in the Pacific Northwest, may be Pinus monticola (western white pine), Pinus albicaulis (whitebark pine), or both.



Table 4.--Land area in Wallowa County, by cover

(In

Cover type or land class		Total unreserved and reserved
		PRODUCTIVE
		Total
1	Ponderosa pine, large sawtimber . . . . .	207,430
2	Ponderosa pine, small sawtimber . . . . .	104,680
3	Ponderosa pine, poletimber . . . . .	48,480
4	Ponderosa pine, seedlings and saplings . . . . .	3,010
5	White pine, <sup>1/</sup> large sawtimber . . . . .	10
6	White pine, small sawtimber . . . . .	20
7	White pine, poletimber . . . . .	1,980
8	Lodgepole pine, small sawtimber . . . . .	19,960
9	Lodgepole pine, poletimber . . . . .	43,730
10	Lodgepole pine, seedlings and saplings . . . . .	980
11	Douglas-fir, small old-growth and large young-growth sawtimber (red fir) . . . . .	92,070
12	Douglas-fir, small sawtimber . . . . .	135,910
13	Douglas-fir, poletimber . . . . .	44,750
14	Douglas-fir, seedlings and saplings . . . . .	510
15	Western larch, large sawtimber . . . . .	22,820
16	Western larch, small sawtimber . . . . .	37,760
17	Western larch, poletimber . . . . .	6,510
18	Western larch, seedlings and saplings . . . . .	720
19	White fir, <sup>2/</sup> large sawtimber . . . . .	27,770
20	White fir, small sawtimber . . . . .	50,000
21	White fir, poletimber . . . . .	6,130
22	White fir, seedlings and saplings . . . . .	50
23	True fir--mountain hemlock, large sawtimber . . . . .	1,950
24	True fir--mountain hemlock, small sawtimber . . . . .	30,990
25	True fir--mountain hemlock, poletimber . . . . .	12,030
26	True fir--mountain hemlock, seedlings and saplings . . . . .	830
27	Engelmann spruce, large sawtimber . . . . .	6,630
28	Engelmann spruce, small sawtimber . . . . .	27,350
29	Engelmann spruce, poletimber . . . . .	2,070
30	Hardwoods, large sawtimber . . . . .	890
31	Hardwoods, small sawtimber . . . . .	2,220
32	Hardwoods, poletimber . . . . .	880
33	Nonstocked area . . . . .	7,430
34	Total . . . . .	948,550
		NONCOMMERCIAL
35	Subalpine . . . . .	54,050
36	Noncommercial, rocky . . . . .	96,430
37	Juniper . . . . .	2,290
38	Total . . . . .	152,770
		NONFOREST
39	Vegetative land (cultivated, grass, or brush) . . . . .	735,320
40	Nonvegetative land (including barrens and cities) . . . . .	172,320
41	Unmeandered water . . . . .	2,870
42	Total . . . . .	910,510
		ALL
Forest land:		
43	Commercial . . . . .	878,110
44	Noncommercial (productive-reserved and unproductive) . . . . .	223,210
45	Total forest land . . . . .	1,101,320
46	Nonforest land . . . . .	910,510
47	Total all land . . . . .	2,011,830

<sup>1/</sup> The white pine cover type, as specified by the Forest Survey in the Pacific Northwest, may be *Pinus monticola* (western white pine), *Pinus albicaulis* (whitebark pine), or both.

<sup>2/</sup> See list of tree species, page 30.

type, ownership class, and land-use class, 1957

acres)

Unreserved						Reserved			
Total	Private	State	County	Federal		Total	State	Federal (national forest)	
				Bureau of Land Mgt.	National forest				

# FOREST LAND

Commercial						Noncommercial (productive-reserved)			
193,400	58,610	810	80	2,490	131,410	14,030	140	13,890	1
103,830	75,650	1,870	30	720	25,560	850	--	850	2
48,470	29,420	660	--	50	18,340	10	10	--	3
3,010	2,980	30	--	--	--	--	--	--	4
10	10	--	--	--	--	--	--	--	5
20	20	--	--	--	--	--	--	--	6
1,980	--	--	--	--	1,980	--	--	--	7
19,900	300	--	--	--	19,600	60	--	60	8
38,380	10,700	--	--	--	27,680	5,350	--	5,350	9
400	400	--	--	--	--	580	--	580	10
82,240	16,510	40	60	310	65,320	9,830	50	9,780	11
122,900	41,990	350	60	880	79,620	13,010	--	13,010	12
40,770	10,790	470	--	170	29,340	3,980	10	3,970	13
500	480	--	--	20	--	10	--	10	14
22,280	730	--	--	--	21,550	540	--	540	15
32,870	7,840	60	40	10	24,920	4,890	--	4,890	16
4,600	2,000	--	--	--	2,600	1,910	--	1,910	17
560	540	--	--	20	--	160	--	160	18
27,160	4,550	70	--	--	22,540	610	--	610	19
49,210	25,930	60	--	--	23,220	790	--	790	20
6,060	980	--	--	--	5,080	70	--	70	21
50	50	--	--	--	--	--	--	--	22
1,950	--	--	--	--	1,950	--	--	--	23
25,930	400	--	--	--	25,530	5,060	--	5,060	24
7,690	530	--	--	--	7,160	4,340	--	4,340	25
690	30	--	--	--	660	140	--	140	26
6,520	100	--	--	--	6,420	110	--	110	27
23,290	1,700	--	--	--	21,590	4,060	--	4,060	28
2,070	90	--	--	--	1,980	--	--	--	29
840	840	--	--	--	--	50	50	--	30
2,220	1,540	--	--	20	660	--	--	--	31
880	230	--	--	--	650	--	--	--	32
7,430	260	--	--	--	7,170	--	--	--	33
878,110	296,200	4,420	270	4,690	572,530	70,440	260	70,180	34

# UNPRODUCTIVE FOREST LAND

5,180	1,230	--	--	--	3,950	48,870	--	48,870	35
50,770	6,250	200	80	1,310	42,930	45,660	--	45,660	36
2,290	310	--	--	--	1,980	--	--	--	37
58,240	7,790	200	80	1,310	48,860	94,530	--	94,530	38

# LAND

660,670	509,890	4,000	50	8,760	137,970	74,650	60	74,590	39
61,790	39,890	1,060	--	5,370	15,470	110,530	40	110,490	40
2,210	740	50	--	130	1,290	660	--	660	41
724,670	550,520	5,110	50	14,260	154,730	185,840	100	185,740	42

# LAND

878,110	296,200	4,420	270	4,690	572,530	--	--	--	43
58,240	7,790	200	80	1,310	48,860	164,970	260	164,710	44
936,350	303,990	4,620	350	6,000	621,390	164,970	260	164,710	45
724,670	550,520	5,110	50	14,260	154,730	185,840	100	185,740	46
1,661,020	854,510	9,730	400	20,260	776,120	350,810	360	350,450	47

Table 5.--Area of commercial forest land in Wallowa County, by  
forest-condition and ownership classes, 1957

(In acres)

Forest-condition class	Total	Private	State	County	Federal	
					Bureau of	National
					Land Mgt.	forest
Softwoods:						
Large sawtimber:						
Uncut	279,650	57,550	560	130	2,430	218,980
Residual	53,910	22,960	360	10	370	30,210
Total	333,560	80,510	920	140	2,800	249,190
Small sawtimber:						
Uncut	251,380	52,760	510	100	1,000	197,010
Residual	126,570	101,070	1,830	30	610	23,030
Total	377,950	153,830	2,340	130	1,610	220,040
Poletimber	150,020	54,510	1,130	--	220	94,160
Seedlings and saplings	5,210	4,480	30	--	40	660
Hardwoods	3,940	2,610	--	--	20	1,310
Nonstocked	7,430	260	--	--	--	7,170
Total	878,110	296,200	4,420	270	4,690	572,530

Table 6.--Area of young-growth timber stands on commercial forest land in Wallowa County, by stand-size class, species group, and stocking class, 1957<sup>1/</sup>

(In acres)

Stand-size class and species group	Total	Well stocked	Medium stocked	Poorly stocked
Young-growth sawtimber:				
Softwoods	377,950	260,340	105,990	11,620
Hardwoods	2,220	1,420	730	70
Total	380,170	261,760	106,720	11,690
Poletimber:				
Softwoods	150,020	88,960	46,680	14,380
Hardwoods	880	840	40	--
Total	150,900	89,800	46,720	14,380
Seedlings and saplings:				
Softwoods	5,210	1,960	2,030	1,220
Hardwoods	--	--	--	--
Total	5,210	1,960	2,030	1,220
All classes:				
Softwoods	533,180	351,260	154,700	27,220
Hardwoods	3,100	2,260	770	70
Total	536,280	353,520	155,470	27,290

<sup>1/</sup> Young-growth stands are those in which the majority of the cubic-foot volume is in trees less than 21.0 inches d.b.h.



Table 7.--Net volume of live sawtimber and growing stock  
on commercial forest land in Wallowa County,  
by ownership class, 1957

Ownership class	Live sawtimber volume		Growing stock volume
	Scribner rule	International	
		1/4-inch rule	
	<u>Million bd.ft.</u>	<u>Million bd.ft.</u>	<u>Million cu.ft.</u>
Private	1,488	1,664	399
State	19	21	5
County	2	2	(1/)
Federal:			
Bureau of Land Mgt.	25	28	6
National forest	5,755	6,322	1,393
Total Federal	5,780	6,350	1,399
All ownerships	7,289	8,037	1,803

1/ Less than 0.5 million.

Table 8.--Net volume of live sawtimber and growing stock  
on commercial forest land in Wallowa County,  
by stand-size class, 1957

	:		:
	:	Live sawtimber volume	:
	:		:
Stand-size class	:		Growing stock
	:	:	volume
	:	International	:
	:	Scribner rule	1/4-inch rule
	:	:	:
<hr/>			
	<u>Million bd.ft.</u>	<u>Million bd.ft.</u>	<u>Million cu.ft.</u>
Sawtimber stands	7,161	7,893	1,694
Poletimber stands	123	138	108
Seedling and sapling stands	1	1	(1/)
Nonstocked areas	4	5	1
	<hr/>		
Total	7,289	8,037	1,803

1/ Less than 0.5 million.

Table 9.--Net volume of live sawtimber and growing stock  
on commercial forest land in Wallowa County,  
by species, 1957

	:	:	:
	:	Live sawtimber volume	:
	:	:	:
Species	:	:	Growing stock
	:	International	volume
	:	Scribner rule	1/4-inch rule
	:	:	:
	:	:	:
		<u>Million bd.ft.</u>	<u>Million bd.ft.</u>
			<u>Million cu.ft.</u>
Softwoods:			
Ponderosa pine	2,476	2,695	500
Douglas-fir	2,058	2,295	437
White fir	920	994	276
Western larch	788	873	159
Engelmann spruce	482	544	142
Lodgepole pine	256	300	164
Subalpine fir	254	274	101
Mountain hemlock	17	18	4
Whitebark pine	5	6	2
Total	7,256	7,999	1,785
Hardwoods:			
Black cottonwood	33	38	10
Quaking aspen	(1/)	(1/)	2
Red alder	--	--	3
Northwestern paper birch	--	--	3
Total	33	38	18
Total all species	7,289	8,037	1,803

1/ Less than 0.5 million.

Table 10.--Net volume of live sawtimber on commercial forest land  
in Wallowa County, by diameter class and species group, 1957

(In million board feet)

Diameter class (inches d.b.h.) and log rule	Total	Ponderosa pine	Douglas- fir	White fir	Western larch	Other soft- woods	Hard- woods
11.0-20.9:							
Scribner rule	3,082	538	1,010	430	319	782	3
International 1/4-inch rule	3,524	624	1,171	464	370	892	3
21.0-30.9:							
Scribner rule	2,641	1,017	720	340	340	195	29
International 1/4-inch rule	2,855	1,098	777	368	367	211	34
31.0-40.9:							
Scribner rule	1,209	736	247	93	103	30	--
International 1/4-inch rule	1,284	781	262	101	109	31	--
41.0 and larger:							
Scribner rule	357	185	81	57	26	7	1
International 1/4-inch rule	374	192	85	61	27	8	1
All diameter classes:							
Scribner rule	7,289	2,476	2,058	920	788	1,014	33
International 1/4-inch rule	8,037	2,695	2,295	994	873	1,142	38

Table 11.--Net volume of all timber on commercial forest land  
in Wallowa County, by class of material and species  
group, 1957

(In million cubic feet)

Class of material	:	Total	:	Softwoods	:	Hardwoods
	:		:		:	
Growing stock:						
Sawtimber trees:						
Saw-log portion		1,303		1,296		7
Upper-stem portion		99		98		1
Total		1,402		1,394		8
Poletimber trees		401		391		10
Total growing stock		1,803		1,785		18
Other material:						
Sound cull trees		7		6		1
Rotten cull trees		6		6		(1/)
Salvable dead trees		14		14		--
Total other material		27		26		1
All timber		1,830		1,811		19

1/ Less than 0.5 million.

Table 12.--Average annual cut of live sawtimber and growing stock on commercial forest land in Wallowa County, by species group, 1953-58

Species group <sup>1/</sup>	Live sawtimber						Growing stock		
	Scribner rule			International 1/4-inch rule					
	Annual	Timber	Logging	Annual	Timber	Logging	Annual	Timber	Logging
	cut <sup>2/</sup>	products	residue	cut <sup>2/</sup>	products	residue	cut <sup>2/</sup>	products	residue
	----- <u>Thousand board feet</u> -----						-- <u>Thousand cubic feet</u> --		
Softwoods	90,162	92,759	1,818	99,414	102,278	2,005	24,276	22,945	1,331

<sup>1/</sup> Hardwood cut insignificant.

<sup>2/</sup> Annual cut is the reduction in inventory volume resulting from the production of the given output of timber products.

## FOREST SURVEY PROCEDURE

Procedures used in the reinventory of Wallowa County were materially different from those used in the initial inventory. This change in procedures accounts for some significant differences in the forest-area and timber-volume statistics obtained.

### Initial Inventory

The initial inventory of the county's forests was conducted in 1935 and 1936 by what is known as the "compilation method." In this method, existing information on forest types, timber volumes, areas logged, and other inventory data were collected from private timber owners and various public agencies. These data were checked in the field for reliability and were adjusted to Forest Survey specifications and standards. Forest-type and timber-volume data for areas not covered by reliable existing information were obtained through field reconnaissance.

All land in the county was classified as either forest or nonforest. Forest land was further classified as commercial or noncommercial; the commercial was still further classified by forest type, stand-size or condition class, and--in the case of young-growth stands--by stocking and age classes. All such types and classes were mapped in place on a 1-inch-to-the-mile base map of each forested township. These township type maps were then superimposed over current ownership-status plats and dot counted to obtain forest-type area statistics by ownership class. Type delineations on the township maps were traced to a base map of the county to form a county forest type map. The commercial forest land was also classified as to site quality, or forest productive capacity.

In-place, timber-volume estimates were based on existing cruises collected and adjusted to the Forest Survey standard, on field samples, and on ocular appraisals. Cruises made by commercial cruisers were obtained for most of the privately owned timber and Forest Service cruises were available for a large part of the national-forest lands. Separate volume estimates were computed for each of the commercial tree species and for each ownership class. Methods used in this initial inventory did not permit a statistical computation of accuracy of the estimate.

### Reinventory

In the 1957 reinventory, the forest type map of the county was completely revised. This revision was accomplished through interpretation, classification, and field mapping on aerial photos that covered all of the forest land in the county. In the mapping on aerial photos, types whose classification was difficult were examined more closely in the field. Likewise, species composition of mixed stands was checked on the ground. The use of aerial photos in mapping resulted in type delineations of much greater

accuracy and detail than were possible through the ground reconnaissance employed in the initial inventory. In the preparation of a revised type map, the delineations on the aerial photos were transferred to a 2-inch county base map through use of a reflecting projector.

Forest type areas were obtained by two methods. In the first method area figures were derived by superimposing the type map over a current ownership-status map and dot counting. This was done for the Pine and Union Working Circles of the Wallowa-Whitman National Forests and the "all other ownership" stratum, which included private, county, and State lands.

The second method, based on a systematic grid of plots, was used to determine area figures for the Wenaha Working Circle and Wenaha Land Management Unit of the Umatilla National Forest and the Wallowa Working Circle of the Wallowa-Whitman National Forests. Each of the three subplots in every plot cluster was first classified as commercial forest, non-commercial forest, or nonforest. The ratio of subplots in each class to the total number of subplots was applied to the total land area to determine the acreage of each classification. Subplots falling on commercial forest land were also classified by forest type and stand-size class as indicated by the subplot tally. The percentage of subplots falling in each type was applied to the total area of commercial forest land in the working circle to determine the acreage of land in that type.

Volume estimates for live sawtimber, growing stock, and salvable dead material for the Pine and Union Working Circles and "all other ownership" stratum were calculated by applying average per-acre volumes to the appropriate forest type acreages. The average per-acre volumes for sawtimber stands and poletimber stands were obtained through a grid sampling procedure in which the stands were measured at a 1.7-mile or 3.4-mile interval, varying with accuracy requirements for different strata. Intensity of sampling was designed to produce a total volume estimate in the county to a specified sampling accuracy set by Forest Survey. The sample plots consisted of clusters of three 1/5-acre circular plots spaced at 6-chain intervals.

Average per-acre volumes of sawtimber and poletimber trees in the overstory of seedling and sapling stands and on nonstocked areas were obtained through an aerial-photo plot-sampling procedure for the Pine and Union Working Circles and "all other ownership" stratum. First, 1-acre photo plots were located through random selection. Next, estimates were made of average number of trees per acre, average crown diameter, and average total height for both sawtimber- and poletimber-size trees. Finally, gross volume of the average tree in each class was obtained from volume tables and then adjusted for defect and breakage to obtain net volume.

In the Wallowa and Wenaha Working Circles and the Wenaha Land Management Unit, volume estimates for all size classes were determined



by species for all subplots falling on commercial forest land. Average per-acre volumes for commercial forest land plots were then expanded by the total acreage of commercial forest land, providing total volume by species for each working circle inventoried.

## ACCURACY OF 1957 REINVENTORY DATA

### Forest Area

The in-place mapping of the forests and their classification by forest type, stand-size class, and condition class were based on 100-percent coverage for the Pine and Union Working Circles and "all other ownership" stratum. Thus no error due to sampling was involved. Errors due to techniques or judgment in the field and in office computation of data were possible, but difficult to evaluate. Throughout all phases of the work, however, close supervision and frequent checks assured a high level of accuracy and uniformity of standards.

The Wallowa and Wenaha Working Circles and Wenaha Land Management Unit were sampled for area, consequently an accuracy figure could be calculated. The chances are 19 out of 20 that the actual total area of commercial forest land in the Wallowa County portions of the three above-mentioned strata is within plus or minus 4.7 percent of the estimated total of 565,001 acres, and that the noncommercial unproductive area is within plus or minus 31.8 percent of the estimated total of 48,169 acres.

### Timber Volume

The chances are 19 out of 20 that the total board-foot volume of live sawtimber, if measured by a 100-percent cruise, would be within plus or minus 8.73 percent of the estimated total of 7,289 million board feet, Scribner rule. On the same basis, cubic-foot volume of growing stock from a 100-percent cruise would be within a range of plus or minus 7.43 percent of the estimated 1,803 million cubic feet. Volume estimates by species, stand-size class, or other subdivision probably have greater sampling errors.

## DIFFERENCES IN RESULTS OF INVENTORIES

Some of the differences between forest-type and timber-volume statistics resulting from the initial inventory and the reinventory are due to physical change--such as cutting of stands, restocking of deforested areas, and ingrowth of stands into the next larger size class. Other differences are due to variations in the procedures used to interpret and classify forest conditions, and to variations in standards of utilization. Differences such as these preclude direct comparison of some of the statistics; comparison of other statistics is meaningful only after they have been adjusted to common standards.

### Forest Area

The total area of forest land has decreased slightly between inventories. Clearing forest land for agriculture and differences in inventory techniques are two possible reasons for the change.

The 1957 reinventory showed a decrease in poletimber, seedling and sapling, and noncommercial areas. This was about compensated for by an increase in the area of sawtimber. Some of this change may be due to growth and less intensive cutting; part is attributable to differing utilization standards and the inventory techniques used.

Table 13.--Comparison of forest area statistics for Wallowa

#### County, initial inventory and reinventory

(Thousand acres)

Forest area	: : 1935-36 : :	: : 1957 : :
Sawtimber	635	715
Poletimber; seedlings and saplings	215	156
Nonstocked	7	7
Noncommercial (productive-reserved and unproductive)	253	223
Total	1,110	1,101

### Timber Volume

The reinventory showed a greater volume of sawtimber for all species than did the initial inventory. Some of this difference may be due to a favorable growth-drain relationship, where the gain in volume through ingrowth of poletimber trees and growth of sawtimber trees is greater than the loss in volume through cutting or destruction by fire, insects, wind-throw, or disease. However, it appears likely that most of the difference is due to changes in standards of utilization and in Forest Survey procedures.

Table 14.--Comparison of timber volume statistics for Wallowa

County, initial inventory and reinventory

(Million board feet, Scribner rule)

Species	1935-36	1957
Ponderosa pine	2,243	2,476
Douglas-fir	968	2,058
Western larch	509	788
White fir	331	920
Other softwoods	175	1,014
Hardwoods	3	33
Total	4,229	7,289

The reinventory showed an increase in sawtimber volume for all species, but the proportionate increase in ponderosa pine was slight in comparison with all others. This is due primarily to two reasons: (1) more of the high-elevation country was included as commercial forest land and (2) more conservative cull estimates were given these minor species because of increasing recognition of their value.

# DEFINITION OF TERMS

## Land Area

### Total Land Area

Includes dry land and unmeandered water surfaces.

### Forest Land Area

Includes (a) land that is at least 10 percent stocked by trees of any size and capable of producing timber or other wood products, or of exerting an influence on the climate or on the water regime; and (b) land from which the trees described in "(a)" have been removed to less than 10-percent stocking and that has not been developed for other use. Minimum area of forest land recognized in reinventory of the county is 10 acres.

### Nonforest Land Area

Land that does not qualify as forest land. Minimum area recognized in the reinventory of the county is 10 acres.

## Forest Land Classes

### Commercial Forest Land Area

Forest land that is (a) producing, or is physically capable of producing, usable crops of wood, (b) economically available now or prospectively, and (c) not withdrawn from timber utilization.

### Noncommercial Forest Land Area

Forest land (a) withdrawn from timber utilization through statute, ordinance, or administrative order but that otherwise qualifies as commercial forest land, or (b) incapable of yielding usable wood products (usually sawtimber) because of adverse site conditions, or so physically inaccessible as to be unavailable economically in the foreseeable future.

## Types

### Forest Land Types

Forest land is typed on the basis of the predominant species, as indicated by cubic volume for sawtimber and poletimber stands and number of trees for seedling and sapling stands, or on the basis of forest condition, such as nonstocked cutover or burned-over land. Where none of the indicated species comprise 50 percent or more of a given stand, the stand is classified on the basis of plurality of cubic volume or number of trees. In classifying forest land by type, the minimum area recognized is 10 acres.

### Commercial Forest Land

Major forest types. Local forest types are grouped into generalized types. The major forest types in Wallowa County are as follows:

Ponderosa pine. Forests in which 50 percent or more of the stand is ponderosa pine.

Lodgepole pine. Forests in which 50 percent or more of the stand is lodgepole pine.

Douglas-fir. Forests in which 50 percent or more of the stand is Douglas-fir.

Western larch. Forests in which 50 percent or more of the stand is western larch.

Fir-spruce. Forests in which 50 percent or more of the stand is true fir, Engelmann spruce, or both.

Hardwoods. Forests in which 50 percent or more of the stand is black cottonwood or other hardwoods, singly or in combination.

#### Noncommercial Forest Land

Productive-reserved. Forest land withdrawn from timber utilization through statute, ordinance, or administrative order, but that otherwise qualifies as commercial forest land.

Unproductive. Forest land incapable of yielding usable wood products (usually sawtimber) because of adverse site conditions, or so physically inaccessible as to be unavailable economically in the foreseeable future.

#### Nonforest Land Types

Vegetative. Cultivated land, stump pasture, grass, or brush on nonforest land.

Nonvegetative. Includes barrens and towns.

Unmeandered water. Includes unmeandered streams and lakes, and tideflats.

#### Tree Classes

##### Sawtimber Tree

Tree of commercial species, 11.0 inches d.b.h. and larger, that contains at least one 16-foot coniferous saw log or one 8-foot hardwood saw log to a variable top diameter never less than 8 inches inside the bark. Also, 25 percent or more of the gross board-foot volume must be free from rot or defect.

##### Poletimber Tree

Tree of commercial species, 5.0 to 10.9 inches d.b.h., in which 25 percent or more of the gross cubic-foot volume is free from rot and defect.

##### Seedling and Sapling Trees

Live trees of commercial species, less than 5.0 inches d.b.h., and of good form and vigor.

## Cull Tree

Live tree of sawtimber or poletimber size that is unmerchantable for saw logs, now or prospectively, because of defect, rot, or species.

Sound cull tree. Live tree of sawtimber or poletimber size that contains 25 percent or more of sound volume but will not make at least one merchantable saw log, now or prospectively, because of roughness, poor form, or species.

Rotten cull tree. Live tree of sawtimber or poletimber size in which less than 25 percent of the total volume is sound.

## Salvable Dead Tree

Standing or down dead tree that contains 25 percent or more of sound volume and at least one merchantable 16-foot coniferous or 8-foot hardwood saw log.

## Stand-Size Classes

### Sawtimber Stand

Stand of sawtimber trees having a minimum per-acre net volume of 1,500 board feet (International 1/4-inch rule).

Large sawtimber stand. Stand in which the majority of the volume is in trees 21.0 inches d.b.h. and larger.

Small sawtimber stand. Stand in which the majority of the volume is in trees from 11.0 to 20.9 inches d.b.h.

Uncut sawtimber stand. Uncut sawtimber stand or sawtimber stand in which less than 10 percent of its volume has been removed by cutting.

Residual sawtimber stand. Sawtimber stand in which 10 percent or more of the volume has been removed by cutting, and in which the residual per-acre volume amounts to 1,500 board feet (International 1/4-inch rule).

### Poletimber Stand

Stand failing to meet sawtimber stand specifications but at least 10 percent stocked with poletimber and larger (5.0 inches d.b.h. and larger) trees and with at least half the minimum stocking in poletimber trees.

### Seedling and Sapling Stand

Stand not qualifying as either a sawtimber or poletimber stand but at least 10 percent stocked with trees of commercial species and with at least half the minimum stocking in seedling and sapling trees.

### Nonstocked Area

An area less than 10 percent stocked with present or potential growing-stock trees.

### Stocking

Stocking is the extent to which growing space is effectively utilized by present or potential growing-stock trees of commercial species. "Degree of stocking" is synonymous with "percentage of growing space occupied" and means the ratio of actual stocking to full stocking for comparable sites and stands. Stocking may be measured in terms of number of trees, volume, basal area, cover canopy, or other criterion or combination of criteria.

Well-stocked stand. Stand that is 70 percent or more stocked with present or potential growing-stock trees.

Medium-stocked stand. Stand that is 40 to 69 percent stocked with present or potential growing-stock trees.

Poorly stocked stand. Stand that is 10 to 39 percent stocked with present or potential growing-stock trees.

Nonstocked area. An area less than 10 percent stocked with present or potential growing-stock trees.

### Timber Volume

#### Live Sawtimber Volume

Net volume in board feet of live sawtimber trees of commercial species:

Scribner rule. The common board-foot log rule used in determining volume of sawtimber in the Pacific Northwest.

International 1/4-inch rule. The standard board-foot log rule adopted nationally by the Forest Service in the presentation of Forest Survey volume statistics.

### Growing Stock

Net volume in cubic feet of live sawtimber trees and live poletimber trees from stump to a minimum 4.0-inch top (of central stem) inside bark.

### All-Timber Volume

Net volume in cubic feet of live and salvable dead sawtimber trees and poletimber trees of commercial species, and cull trees of all species from stump to a minimum 4.0-inch top inside bark.

### Timber Cut

#### Annual Cut of Live Sawtimber

The net board-foot volume of live sawtimber trees cut or killed by logging on commercial forest land during a specified year.

Timber products from live sawtimber. The volume of timber products cut from live sawtimber.

Logging residues from live sawtimber. The volume of sound wood in live sawtimber trees cut or killed by logging on commercial forest land and not converted to timber products.

#### Annual Cut of Growing Stock

The net cubic-foot volume of live sawtimber and poletimber trees cut or killed by logging on commercial forest land during a specified year.

Timber products from growing stock. The volume of timber products cut from growing stock.

Logging residues from growing stock. The volume of sound wood in growing stock cut or killed by logging on commercial forest land and not converted to timber products.



## TREE SPECIES

Tree species commonly found in Wallowa County include:


### Softwoods:

Ponderosa pine (Pinus ponderosa)  
Whitebark pine (Pinus albicaulis)  
Lodgepole pine (Pinus contorta)  
Douglas-fir (Pseudotsuga menziesii)  
Western larch (Larix occidentalis)  
White fir (Abies concolor or A. grandis)<sup>1/</sup>  
Subalpine fir (Abies lasiocarpa)  
Engelmann spruce (Picea engelmannii)  
Mountain hemlock (Tsuga mertensiana)

### Hardwoods:

Red alder (Alnus rubra)  
Black cottonwood (Populus trichocarpa)  
Quaking aspen (Populus tremuloides)  
Northwestern paper birch (Betula papyrifera  
var. subcordata)

<sup>1/</sup> No attempt was made to separate A. concolor (white fir) from A. grandis (grand fir). White fir, as specified in the Forest Survey in the Pacific Northwest, may be concolor, grandis, or both.



I N D E X M A P of

W A L L O W A C O .

S T A T E of O R E G O N

showing divisions of 2" = 1 mile maps