



Open Access Articles

Covering Nearly Surface-Embedded Graphs with a Fixed Number of Balls

The Faculty of Oregon State University has made this article openly available.
Please share how this access benefits you. Your story matters.

Citation	Borradaile, G., & Chambers, E. W. (2014). Covering nearly surface-embedded graphs with a fixed number of balls. <i>Discrete & Computational Geometry</i> , 51(4), 979-996. doi:10.1007/s00454-014-9594-5
DOI	10.1007/s00454-014-9594-5
Publisher	Springer
Version	Accepted Manuscript
Terms of Use	http://cdss.library.oregonstate.edu/sa-termsofuse

Covering nearly surface-embedded graphs with a fixed number of balls

Glencora Borradaile

Erin Wolf Chambers

April 1, 2014

Abstract

A recent result of Chepoi, Estellon and Vaxes [Disc. Comp. Geom. '07] states that any planar graph of diameter at most $2R$ can be covered by a constant number of balls of size R ; put another way, there are a constant-sized subset of vertices within which every other vertex is distance half the diameter. We generalize this result to graphs embedded on surfaces of fixed genus with a fixed number of apices, making progress toward the conjecture that graphs excluding a fixed minor can also be covered by a constant number of balls. To do so, we develop two tools which may be of independent interest. The first gives a bound on the density of graphs drawn on a surface of genus g having a limit on the number of pairwise-crossing edges. The second bounds the size of a non-contractible cycle in terms of the Euclidean norm of the degree sequence of a graph embedded on surface.

1 Introduction

Chepoi, Estellon and Vaxès showed there is a constant ρ such that any planar graph of diameter at most $2R$ has a subset of at most ρ vertices such that every vertex in the graph is within distance R of that subset [2]. Since this can be viewed as showing that there is a constant-sized set cover in the set system of balls of radius R , we refer to this property as the *ball-cover* property. Graphs having constant-sized ball covers admit interval routing schemes with dilation $\frac{3}{2} \times \text{diameter}$ and compactness $O(1)$ where dilation measures the indirectness of the routing scheme and compactness measures the size of the routing table [8]. We believe the ball-cover property is an inherently interesting property. Graphs having this property could define an interesting class of graphs and perhaps could have broader utility than previously realized.

We generalize the class of graphs having the ball-cover property to those graph families that can be embedded on a surface of fixed genus after the removal of a constant number of vertices (the *apices*); the number of balls required depends only on the genus of the surface (either orientable or non-orientable) and the number of apices. Since graphs of bounded treewidth are also known to have the ball-cover property [8] by way of the Graph Minor Structure Theorem, our result is a significant step toward proving that fixed-minor-excluded graphs also have the ball-cover property. We discuss this more in Section 5. We start by sketching the proof for the planar case as we use a similar, but more general, tool set here.

1.1 A sketch of the proof of the ball-cover property for planar graphs

Throughout, graphs are simple, undirected and unweighted. Let $B(x)$ be the set of all vertices that are within distance R of vertex x in graph G ; this is the *ball centered at x* . Let $\mathcal{B}(G) = \{B(x) : x \in V(G)\}$; this is the *ball system* of G . We say that $\mathcal{B}' \subset \mathcal{B}$ covers G if \mathcal{B}' is a set cover of $V(G)$.

The dual of a set system \mathcal{S} with ground set U is defined as follows: the ground set of the dual set system is \mathcal{S} and for every element $x \in U$, the dual system has a set representing the sets of \mathcal{S} containing x , i.e., $X = \{S : S \in \mathcal{S}, x \in S\}$. It is easy to see:

Observation 1. *The dual set system of $\mathcal{B}(G)$ is $\mathcal{B}(G)$.*

Since a hitting set of \mathcal{S} (a subset of the ground set that contains an element in every set) is a set cover of the dual set system of \mathcal{S} , we likewise have that the centers of a subset of balls covering G is a hitting set for the ball system. A hitting set of $\mathcal{B}(G)$ is exactly a subset of vertices within which every other vertex is distance R .

Matoušek gives a characterization of set systems that have small hitting sets [13] in terms of the set system's *fractional-Helly* or (p, q) -*property* and the dual set system's *VC-dimension*.

VC-dimension

A set system \mathcal{S} *shatters* a set X if for every subset Y of X there is a set $S \in \mathcal{S}$ such that $S \cap X = Y$. The Vapnik-Chervonenkis dimension or VC-dimension of \mathcal{S} is the maximum size of a set that \mathcal{S} can shatter [25]. Chepoi et al. remark that the VC-dimension of the ball system of a graph excluding K_{r+1} as a minor is at most r [2]. This gives us:

Lemma 1. *The VC-dimension of ball system of a graph excluding H as a minor is at most $|H| - 1$.*

Recall that a minor of a graph G is a graph that is obtained from G by edge contractions and deletions; a forbidden or excluded minor is a graph that *cannot* be obtained this way. It follows from Observation 1 that the dual of the ball system of a graph excluding K_{r+1} as a minor also has VC-dimension at most r .

Fractional Helly theorems

If a set system is such that every d sets has a point in common, then the set system is said to have Helly order d . A *Helly theorem* is one that shows that certain set systems of Helly order d have a non-empty intersection. The first such theorem was given for the Euclidean plane: if a family of convex sets has a nonempty intersection for every triple of sets, then the whole family has a nonempty intersection [12]. A set system has *fractional Helly order* (p, q) , or *has the (p, q) -property*, if among every p sets some q have a point in common. Matoušek gave the following fractional Helly theorem:

Theorem 1 (Fractional Helly Theorem [13]). *Let \mathcal{Q} be a set system having the (p, q) -property (for $p \geq q$) and whose dual set system has VC-dimension $q - 1$. Then there is a constant ρ such that \mathcal{Q} has a hitting set of size at most ρ .*

Given Lemma 1, one could therefore show that, for a fixed minor H , H -minor free graphs have the ball-cover property by showing that the corresponding ball system has fractional Helly order $(p, |H|)$ for some fixed $p \geq |H|$. Chepoi et al. do just this for planar graphs. Starting with p vertices, they consider the pairwise shortest paths between these vertices; each shortest path contains a vertex that is contained by the balls centered on the paths' endpoints. Viewing these shortest paths as edges of a complete graph and drawn on the plane (as inherited from a drawing of the original graph), they invoke a result showing that such a drawing of K_p , for p sufficiently large, must contain at least 7 pairwise crossing edges. The 7 pairwise crossing shortest paths then witness a point in common to 5 of the balls. We use this idea at the heart of our proof for surface-embedded graphs.

1.2 Surface-embedded graphs

We start by extending this result to graphs embedded on more general surfaces. We first give some definitions.

A 2-manifold (or surface) S is a Hausdorff space in which every point has a neighborhood homeomorphic to the Euclidean plane or the closed half plane. A cycle in a surface is a continuous function from S^1 to the surface; the cycle is called simple if the map is injective. A simple cycle γ is separating if $S \setminus \gamma$ is not connected; see Figure 1. The genus g of a surface S is the maximum number of pairwise disjoint non-separating cycles $\gamma_1, \gamma_2, \dots, \gamma_g$ such that $S \setminus (\gamma_1 \cup \dots \cup \gamma_g)$ is connected. Note that cutting a surface along a non-separating cycle reduces the genus by 1; this is a common algorithmic technique for reducing the complexity of a surface. A surface is non-orientable if and only if it contains a subspace homeomorphic to the Möbius band and is otherwise orientable.




Figure 1: Left: An example of homologous cycles: the single dashed red non-separating cycle (above on left) is \mathbb{Z}_2 -homologous to two solid blue cycles. Right: A null-homologous separating cycle.

An embedding of a graph $G = (V, E)$ on a surface S is a drawing of G on S , such that vertices are mapped to distinct points in S and edges are mapped to *internally* disjoint simple paths. A face of an embedding is a maximal connected subset of S that does not intersect the image of G . An embedding is cellular if all of its faces are homeomorphic to a topological open disc. We say that G is a graph of (orientable or non-orientable) genus g if G has a cellular embedding on a surface of (orientable or non-orientable) genus g .

We will briefly use the notion of \mathbb{Z}_2 -homology in this paper and so include a brief description for completeness; we refer the reader to a topology text for full details [11, 16]. A homology cycle is a linear combination of oriented cycles with coefficients from a ring R ; when $R = \mathbb{Z}_2$, these homology cycles are even-degree subgraphs of G . A boundary subgraph is the boundary of a union of faces of G . Two subgraphs are homologous if their symmetric difference is a boundary subgraph, or, more intuitively, if they can be deformed to each other (where the deformation may include merging intersection cycles or splitting at self-intersections or deleting trivial separating cycles); see Figure 1 for an example. Boundary cycles are null-homologous, and since every separating cycle is a boundary cycle, we can view separating cycles as the identity element for homology classes.

1.3 Our contribution

The bulk of this paper focusses on showing that a graph of genus g has the ball-cover property by showing that its ball system has the (p_g, q_g) -property for numbers p_g and q_g that depend only on g (Section 4). Since K_n has orientable genus $\lceil \frac{1}{12}(n-3)(n-4) \rceil$ and non-orientable genus $\lceil \frac{1}{8}(n-3)(n-4) \rceil$ [18], we set $q_g = c \cdot g^2$ (where c depends only on whether the surface in question is orientable). Then, since a graph of genus at most g excludes K_{q_g} as a minor, the VC-dimension for a graph of genus at most g is at most $q_g - 1$. By Observation 1, Lemma 1 and the Fractional Helly Theorem, we will get:

Theorem 2. *There exists a constant ρ_g (depending only on g) such that any graph of genus at most g and diameter at most $2R$ can be covered by at most ρ_g balls of radius R .*

We show that the same holds if the graph additionally has a fixed number of apices and discuss how one might generalize to fixed-minor-excluded graph families in Section 5.

In order to prove that the ball system for a genus- g graph has the (p_g, q_g) -property, we show that there is a small set of edges of a surface-embedded graph whose removal leaves a planar graph (Section 2) and give bounds on the number of edges in a graph drawn on a surface of fixed genus having a limit on the number of crossings (Section 3). The former result can be used to generalize an edge-separator result for planar graphs due to Gazit and Miller [9]. Both these results are likely of more general interest. We give background on these problems in their relevant sections.

The takeaway from these generalizations will allow us to argue that any topological drawing of K_n on a surface of orientable or non-orientable genus g must have a large subset of edges that pairwise cross. In Section 3, we will formally define what constitutes a topological drawing on a surface of genus g and prove this theorem.

2 A norm-sized, planarizing edge set for surface-embedded graphs

In this section, we show there is a small set of edges in a surface-embedded graph whose removal leaves a planar graph. We start by bounding the size of a non-separating cycle:

Theorem 3. *The shortest non-separating cycle of a graph G embedded on a surface has length at most $\frac{1}{2}\|G\|_f$.*

where

$$\|G\|_f = \sqrt{\sum_{f \in \mathcal{F}} |f|^2}$$

is the *face-norm* of G and \mathcal{F} is the set of faces of G . We use a sequence of g non-separating cycles to *planarize* G . The face-norm was used by Gazit and Miller to tighten the bound on the size of edge-separators for planar graphs [9]. Theorem 3 implies an $O(g\|G\|_f)$ -sized edge separator for genus- g graphs. We discuss some open problems in this vein at the end of the paper.

Let G be a graph with a cellular embedding on a surface of genus g (either orientable or not). We start with a shortest non-separating cycle C and generate an ordered family of disjoint cycle sets \mathcal{C} each of which is homologous to C . We use this family to build another non-separating cycle C' formed by one vertex from each set in \mathcal{C} . Since C is shortest, C' acts a witness giving a lower bound on $|\mathcal{C}|$. Overall, this gives a lower bound on the number of edges in \mathcal{C} , and so an upper bound on $|C|$.

We appeal to a combinatorial embedding of the graph which gives, for each vertex v , a clockwise ordering of the edges incident to v as they are embedded around v [5, 26]. We note that any such embedding can be maintained under operations such as contraction, deleting, or cutting along a cycle, via appropriate unions, deletions, or duplications of the vertex lists which maintain the clockwise orderings; full details are described by Mohar and Thomassen [15].

In the following ∂f denotes the boundary of face f .

Lemma 2. *Let G be a graph with a cellular embedding on a surface \mathcal{S} , either orientable or non-orientable. Let \mathcal{F} be a set faces of G . We can add a set L of edges to G such that*

- L can be incorporated into the embedding of G in a noncrossing way.
- The endpoints of L are the set of vertices at distance one from the boundaries of \mathcal{F} .
- L decomposes into a set of cycles that is homologous to the boundaries of \mathcal{F} .

Proof. For a face $f \in \mathcal{F}$, let ∂f denote the cycle in G giving f 's boundary, taken in clockwise order. Let X be the set of vertices at distance 1 from \mathcal{F} in G . If $f, g \in \mathcal{F}$ are adjacent in G (that is, there is an edge uv such that $u \in \partial f$ and $v \in \partial g$ or f and g share a vertex x), then the vertices at distance 1 from ∂f interferes with ∂g . To avoid this, we merge adjacent faces. If f and g share a vertex x , we cut open the graph at x , merging the interiors of f and g and creating two copies of x , both on the boundary of the newly created face. If f and g are connected by an edge uv , we cut open the graph along uv , merge the interior of f with that of g resulting in face h . The edge uv is duplicated and both copies appear in ∂h . We repeat this operation *minimally* until the distance between every pair of faces is at least 2: that is, performing a sequence of such operations will guarantee that the interior of the resulting faces are homeomorphic to a disk. (Note that on a non-orientable surface, this minimality avoids the possibility that the union of neighboring faces spans a Möbius strip, and so the interior remains a topological disk.) Let \mathcal{F}' be the resulting set of faces and G' the resulting graph. Note that the boundaries of \mathcal{F} are \mathbb{Z}_2 -homologous to the boundaries of \mathcal{F}' , since the introduction of uv twice cancels under \mathbb{Z}_2 homology. Note further that interior of each face in \mathcal{F}' is homologous to a disk and thus the boundaries are contractible, and the set of vertices at distance one from \mathcal{F}' is still X , the set of vertices at distance 1 from \mathcal{F} .

Let G'' be the graph obtained by contracting the boundaries of the faces of \mathcal{F}' . Let F' be the vertices resulting from these contractions. Note again that the set of vertices at distance 1 from F' in G'' is still X , since each vertex at distance 1 from F' must also be within distance 1 of some vertex in $\partial F'$, and vice versa.

We will build a cycle that is homologous to each face in \mathcal{F}' whose vertices are among X . Since the faces in \mathcal{F}' are at distance at least two from each other, the cycles we construct will not interact with each other.




Figure 2: Cycles which are connected by an edge are merged into a single face (shaded above, right), and level edges (shown dashed above) are embedded so that the boundary of the face, incident edges and new edge bounds a topological disk (shaded above, left).

Subdivide every self-loop ℓ adjacent to a vertex in \mathcal{F}' into two edges with a vertex v_ℓ . Let G''' be the resulting graph. The set of vertices at distance 1 is now X' , which consists of vertices from X and vertices which came from loop subdivisions.

For each vertex $f \in F'$, consider the cyclic clockwise ordering of the edges incident to the vertex corresponding to f in the embedding of G''' . For every two consecutive edges fu and fv in this order we introduce the edge uv and call it a *level edge*. Edge uv can be embedded to be arbitrarily close to fu followed by fv ; on the original surface, this corresponds to a path following the edge from u to the face f , followed by a (possibly empty) portion of the face boundary ∂f , followed by the edge from f to v ; see Figure 2. Let L be the set of all such edges. Since each such edge can be embedded as described to follow two adjacent edges in the clockwise ordering around the vertex f , $G''' \cup L$ can be embedded in a non-crossing way. Note that self-loops and parallel edges may be introduced this way, e.g. when a vertex $f \in F$ has degree 1 or 2, respectively. See Figure 3.

The level edges corresponding to f inherit a cyclic ordering from the ordering of the edges adjacent to f . That is, uv and vw are consecutive in this ordering if fu, fv, fw are consecutive in the ordering of edges adjacent to f . Further, given how we have embedded uv , we know that the cycle ∂f union the edges fu, uv, fv bounds a topological disk. This implies a partitioning of L into a set of cycles \mathcal{C} that is homologous to the boundaries of \mathcal{F}' : simply replace each portion of a face ∂f with the path fu, uv, fv . Since we are (in \mathbb{Z}_2 homology sense) adding a set of disks to a cycle, each new cycle is homologous to the original. This proves the second and third implications of Lemma 2.

However, the endpoints of L are not necessarily vertices of G , since they include the subdividing vertices. Refer to Figure 3. Consider such a vertex $v_\ell \in X'$ which was used to subdivide self-loop ℓ . Merge any two consecutive edges $uv_\ell, v_\ell w$, creating edge uw and minimally modify the embedding so that uw does not intersect ℓ . This maintains the second and third implications. If there are parallel loops (either on an oriented or non-oriented surface), the connecting level edges consist of bigons between loop vertices; these bigons are null-homologous and hence can be disregarded. The set of level edges may also have included a self-loop centered at a subdividing vertex, the new “edge” will no longer have any endpoints. This “edge” must bound a topological disk, since, if we introduced a level edge centered at v_ℓ , ℓ must have bounded a face in G'' . Therefore, we can remove this “edge” while maintaining the same homology type for our set of cycles. We let L' be the modified and remaining edges. These are the edges satisfying the three implications of Lemma 2. □

2.1 Short non-separating cycles

We are now ready to prove Theorem 3.




Figure 3: A face f (shaded) and its incident edges (solid) with added subdividing vertices (hollow). Left: the level edges L (dashed) added to G''' . Right: the level edges after connections to the subdividing vertices are removed. Note that the outer endpoint-less “edge” is not included in L' .

Let C be the shortest non-separating cycle of G . Cut open the surface and graph along C , duplicating C into copies C_0 and C'_0 ; let G_0 be the cut open graph. Glue a disk onto each hole left from cutting open the graph. C_0 and C'_0 are now the boundaries of faces.

Let V_i be the set of vertices in G_0 that are at distance i from C_0 and let s be the smallest index such that $V_s \cap V(C'_0) \neq \emptyset$. We define sets of cycles C_i in a graph G_i , $i = 0 \dots s$, starting with C_0 , inductively as follows: Given the set of cycles C_{i-1} that are the boundaries of faces (and starting with C_0 as our initial cycle), we define C_i to be the homologous set of cycles going through V_i as guaranteed by Lemma 2. We remove the edges and vertices of C_{i-1} and the edges adjacent to C_{i-1} to make C_i the boundaries of faces.

For any chord uv of a face f , let P_{uv} be the shortest u -to- v path along the boundary of f and let $\ell(uv) = |P_{uv}|$. Gazit and Miller [9] show that for a face f and a set of pairwise non-crossing chords H across f , $\ell(H) \leq \frac{1}{8}|f|^2$. Since the edges of $\cup_{i=1}^s C_i$ are chords of the faces of G , we get

$$\sum_{i=0}^s \ell(C_i) \leq \frac{1}{8}(\|G\|_f)^2 \quad (1)$$

By construction C_i is homologous to C_0 and so to C . Let \bar{C}_i be the set of cycles obtained from C_i by replacing each edge $uv \in C_i$ with P_{uv} . We get $|\bar{C}_i| = \ell(C_i)$. Since \bar{C}_i is homologous to C , \bar{C}_i must contain a non-separating cycle S . Since C is the shortest non-separating cycle,

$$|\bar{C}_i| \geq |S| \geq |C|$$




Figure 4: C is the shortest non-separating cycle, $A \cup B$ is another non-separating cycle.

Let B be the shortest path from C_0 to C'_0 . Let A be the shortest subpath of C that connects B 's endpoints. $A \cup B$ is a non-separating cycle. See Figure 4 Since $|B| = s$ and $|A \cup B| \geq |C|$, $s \geq |C|/2$. We

have:

$$\frac{1}{8}(\|G\|_f)^2 \geq \sum_{i=1}^s \ell(C_i) = \sum_{i=1}^s |\bar{C}_i| \geq |C|^2/2$$

Rearranging gives Theorem 3.

2.2 Planarizing sets

Repeatedly cutting along non-separating cycles allows us to reduce a surface-embedded graph to a planar graph, while only reducing the face norm:

Lemma 3. *Let G be an embedded graph and let C be a non-separating cycle. Cutting open the graph along C and then contracting each resulting copy of C results in a graph G' such that $\|G'\|_f < \|G\|_f$.*

Proof. Let \mathcal{F} be the set of faces of G and let \mathcal{F}_C be the set of faces of G that have a bounding edge in C . Cutting along a non-separating cycle C of a graph G embedded on surface \mathcal{S} introduces two holes, each bounded by a copy of C . Contracting each hole and each bounding copy of C results in a graph G' with face set \mathcal{F}' . Every face in \mathcal{F} maps to a face in \mathcal{F}' such that the faces in $\mathcal{F} \setminus \mathcal{F}_C$ are the same size as their image in \mathcal{F}' and the faces in \mathcal{F}_C are strictly larger than their counterparts \mathcal{F}'_C in \mathcal{F}' giving:

$$\|G'\|_f = \sqrt{\sum_{f \in \mathcal{F}'} |f|^2} = \sqrt{\sum_{f \in \mathcal{F}' \setminus \mathcal{F}'_C} |f|^2 + \sum_{f \in \mathcal{F}'_C} |f|^2} < \sqrt{\sum_{f \in \mathcal{F} \setminus \mathcal{F}_C} |f|^2 + \sum_{f \in \mathcal{F}_C} |f|^2} = \sqrt{\sum_{f \in \mathcal{F}} |f|^2} = \|G\|_f$$

□

□

Cutting along a non-separating cycle C of a graph G embedded on surface \mathcal{S} reduces the genus of the surface by one and introduces two holes, each bounded by a copy of C . Lemma 3 shows that if we contract the two copies of C (and the corresponding holes), we only reduce the face-norm of the graph. We can repeat this cut-and-contract procedure g times, each time we find a non-separating cycle of length at most $\frac{1}{2}\|G\|_f$, at which point the surface is a sphere and the final graph G' is planar. Of course, applying this method to the dual G^* of the graph, results in a set of *planarizing* edges whose size is measured in terms of the vertex-norm

$$\|G\|_\delta = \sqrt{\sum_{v \in V} \delta(v)^2}$$

of G where $\delta(v)$ is the degree of vertex v . Recall that the dual of a plane graph is given by a vertex for every face of the primal graph, with dual vertices connected when the corresponding primal faces are adjacent. By duality, the degree of a vertex is the size of the face in the dual corresponding to the vertex. We get:

Lemma 4. *There is a set of $\frac{g}{2}\|G\|_\delta$ edges of a genus- g graph whose removal leaves a planar graph.*

3 Pairwise-crossing number of surfaces

There are many measures of how close a graph is to being planar. One measure is the crossing number which is the minimum number of edge crossings in a planar, topological drawing of the graph [24]. A drawing is *topological* if vertices map to distinct points and edges map to simple Jordan arcs connecting the points their endpoints such that (i) no arc passes through a vertex different from its endpoints, (ii) no two arcs meet in more than one point, and (iii) no three arcs share a common interior point. Formally the crossing number of a fixed drawing is number of interior points that are shared by two arcs. The restriction to topological drawings does not increase the crossing number of a graph, see e.g. [6]. Rather than planar drawings, we are interested in drawings on surfaces of genus g and so will refer to *surface topological drawings*. This number has been studied by Shahrokhi, Székely, Sýkora and Vrt'o, who give upper and lower bounds on the crossing

number of complete graphs drawn on compact 2-manifolds [20, 21]; more specific bounds are also known for surfaces such as the torus [10].

We first use the crossing number of a particular drawing of a graph to give bounds on the size of a set of edges whose removal results in a topological drawing in the plane.

Lemma 5. *A graph G admitting a topological drawing on a surface \mathcal{S} of genus g with χ crossings has a subset of at most*

$$\frac{g}{2} \sqrt{16\chi + \|G\|_\delta^2}$$

edges whose removal leaves a graph whose inherited drawing is a planar topological drawing.

Proof. Let H be the graph embedded on \mathcal{S} obtained from G by introducing a vertex at each crossing. Since the drawing is topological, each of these new vertices has degree 4. We have that $\|H\|_\delta^2 = \sum_{v \in H} \delta_H(v)^2 = 16\chi + \sum_{v \in G} \delta_G(v)^2$. By Lemma 4, H has a planarizing edge set S_H with at most $\frac{g}{2} \|H\|_\delta$ edges. Let S_G be the set of edges of G from which S_H are generated. Since $|S_G| \leq |S_H|$, the lemma follows. \square \square

Another class of graphs that is close to being planar are the class of *k-quasi-planar graphs*. A graph is *k-quasi-planar* if it admits a planar, topological drawing in which no subset of $k + 1$ edges pairwise cross; thus a graph that is 1-quasi-planar is planar. Various bounds on the number of edges in such graphs have been given [1, 7, 17], culminating in:

Theorem 4 (Suk and Walczak [22]). *A simple n -vertex graph admitting a topological drawing in the plane in which no subset of $k + 1$ edges pairwise cross has at most $c_k n \log n$ edges where c_k is a constant depending only on k .*

In fact, if one follows the dependence on k through Suk and Walczak's work, one finds that

$$c_k = A^{k^k} \text{ for a fixed constant } A \geq 2 \quad (2)$$

As far as we know, such bounds have not previously been generalized to more general surface topological drawings as we do so here. The proof of Theorem 5 is based on the analysis technique of Pach et al. [17], but here we are able to immediately reduce the genus g topological graph to a planar, topological graph, thus invoking Suk and Walczak's result [22].

Theorem 5. *A simple n -vertex graph admitting a topological drawing on a surface of genus $g > 0$ in which no subset of $k + 1$ edges pairwise cross has at most $(2g^2)^k c_k n \log n$ when $g = O(n)$.*

Proof. Let $\mathcal{G}_{g,k,n}$ be the family of all graphs with at most n vertices and admitting a genus- g topological drawing in which no subset of $k + 1$ edges pairwise cross. Let $m_{g,k,n}$ be the maximum number of edges in any graph in $\mathcal{G}_{g,k,n}$.

We aim to prove the assertion for $\mathcal{G}_{g,k,n}$ that

$$m_{g,k,n} \leq (2g^2)^k c_k n \log n \quad (3)$$

by induction over k . For $k = 1$ (and every g and n), the assertion is true since such graphs are genus- g graphs and have $O(n + g)$ edges which is $O(n)$ for $g = O(n)$. For values of n such that $n \log n \leq (2g^2)^k c_k$, the assertion is true since the right-hand side of Inequality (3) exceeds n^2 for all such values of n . We assume that $m_{g,k-1,n} \leq (2g^2)^{k-1} c_{k-1} n \log n$.

Consider a graph $G \in \mathcal{G}_{g,k,n}$ and fix a genus- g topological drawing of G in which no subset of $k + 1$ edges pairwise cross. Let χ be the number of crossings in this drawing. We first bound χ so we may use Lemma 5.

Consider an edge e of G and let G_e be the subgraph of G consisting of all the edges crossing e . Let G_e inherit its drawing from G . Since the drawing of G has no $k + 1$ pairwise crossing edges, the drawing of G_e has no k pairwise crossing edges for otherwise such a set along with e would witness a set of $k + 1$ pairwise crossing edges in the drawing of G . Therefore $G_e \in \mathcal{G}_{g,k-1,n}$ and so G_e has at most $m_{g,k-1,n}$ edges. The

number of crossings on e is therefore at most $m_{g,k-1,n}$. Summing over all edges of G , $\chi \leq \frac{1}{2}m \cdot m_{g,k-1,n}$ where m is the number of edges in G . By the inductive hypothesis,

$$\chi \leq \frac{1}{2}m \cdot (2g^2)^{k-1}c_{k-1}n \log n. \quad (4)$$

Let S be the set of edges forming a planarizing set for G guaranteed by Lemma 5. By Lemma 5, Equation (4) and the fact that $\|H\|_\delta^2 \leq 2|E(H)| \cdot |V(H)|$ for any graph H ,

$$|S| \leq \frac{g}{2} \sqrt{8m \cdot (2g^2)^{k-1}c_{k-1}n \log n + 2mn} \leq \frac{3g}{2} \sqrt{m \cdot (2g^2)^{k-1}c_{k-1}n \log n} \quad (5)$$

where the last inequality holds for n such that $2 < (2g^2)^{k-1}c_{k-1} \log n$; these coincide with non-base-case values of n . Let G' be the graph obtained by deleting S from G . Then $m \leq E(G') + |S|$. Since G' is a k -quasi-planar graph on at most n vertices, $|E(G')| \leq c_k n \log n$ by Theorem 4. Combining, we get

$$m \leq c_k n \log n + \frac{3g}{2} \sqrt{m \cdot (2g^2)^{k-1}c_{k-1}n \log n}$$

Rearranging:

$$m - \frac{3g}{2} \sqrt{(2g^2)^{k-1}c_{k-1}n \log n \sqrt{m}} \leq c_k n \log n \quad (6)$$

Let $f(m) = m - \frac{3g}{2} \sqrt{(2g^2)^{k-1}c_{k-1}n \log n \sqrt{m}}$. We consider the two cases corresponding to the sign of the left-hand side of (6).

If $f(m) \leq 0$, then

$$m \leq \left(\frac{3g}{2}\right)^2 (2g^2)^{k-1}c_{k-1}n \log n = (2g^2)^k \frac{9}{8}c_{k-1}n \log n \leq (2g^2)^k c_k n \log n,$$

where the last inequality follows from $\frac{9}{8}c_{k-1} < c_k$ (which is clearly true given Equation (2)), thus proving the assertion.

We note that $f(m)$ is an increasing function for all positive values of m such that $f(m) > 0$. We will show that

$$f((2g^2)^k c_k n \log n) > c_k n \log n, \quad (7)$$

implying that $m < (2g^2)^k c_k n \log n$ when $f(m) > 0$, proving the assertion.

$$\begin{aligned} f((2g^2)^k c_k n \log n) &= (2g^2)^k c_k n \log n - \frac{3g}{2} \sqrt{(2g^2)^{k-1}c_{k-1}n \log n} \sqrt{(2g^2)^k c_k n \log n} \\ &= (2g^2)^k c_k n \log n - \frac{3\sqrt{2}}{4} \sqrt{(2g^2)^{2k} c_{k-1}c_k n \log n} \\ &= (2g^2)^k c_k n \log n \left(1 - \frac{3\sqrt{2}}{4} \sqrt{\frac{c_{k-1}}{c_k}}\right) \\ &> (2g^2)^k c_k n \log n \left(1 - \frac{3\sqrt{2}}{4} \frac{1}{\sqrt{2}}\right), \text{ since } c_k > 2c_{k-1}, \text{ by Equation (2)} \\ &= (2g^2)^k c_k n \log n \left(\frac{1}{4}\right) \\ &> c_k n \log n, \text{ for } k \geq 2 \text{ and } g \geq 1 \end{aligned}$$

This proves Equation (7) and so the theorem. □ □




Figure 5: If P and Q are both shortest paths, then P' must also be a shortest path between P 's endpoints.




Figure 6: Making a drawing topological; full details are given by Felsner [6].

4 The (p_g, q_g) -property of genus- g ball systems

The proof of the fact that the ball system of a graph of genus g has the (p_g, q_g) -property is similar to the proof of Proposition 2 in the work of Chepoi, Estellon and Vaxès [2], although we have made efforts to simplify the proof here.

Let G be a graph of diameter at most $2R$ with an embedding on a surface \mathcal{S} of genus g . Let C be a set of p_g vertices; we will define p_g shortly. Consider a set of shortest paths $\mathcal{P} = \{P_{ij} : c_i, c_j \in C\}$ where P_{ij} is the shortest c_i -to- c_j path in G . We can assume, without loss of generality, that the intersection of any two of these paths is *simple*, having at most one component (a path or vertex), for otherwise, one path could be redirected along another without compromising shortness as illustrated in Figure 4.

Taking the image of P_{ij} on the surface for each path $P_{ij} \in \mathcal{P}$, we get a drawing of the complete graph K_{p_g} on \mathcal{S} . We can make this drawing topological by a sequence of simple, local transformations, as illustrated in Figure 4. Since we assumed that path intersections are simple, the first transformation modifies the drawing to achieve the first and third properties of a topological drawing and the second transformation modifies the drawing to achieve the second property of a topological drawing. These transformations *respect intersection* so far as that, in the final drawing of K_{p_g} , the images of two edges of K_{p_g} share a point if and only if the corresponding paths share a vertex in G .

Since the drawing of K_{p_g} is a topological drawing on surface \mathcal{S} of genus g , we can use Theorem 5 to guarantee that, for p_g sufficiently large (and depending only on g), this drawing contains a subset of at least $2q_g - 3$ edges that pairwise cross. Likewise, since the drawing of K_{p_g} respects intersections, there must be a subset \mathcal{P}' of at least $2q_g - 3$ paths of \mathcal{P} that pairwise intersect. We pick the *midpoint* of a c_i -to- c_j path $P_{ij} \in \mathcal{P}$ to be any vertex m_{ij} that is in $B(c_i) \cap B(c_j)$; since diameter of the graph is at most $2R$, the paths \mathcal{P} are shortest and the balls have radius R , such a point always exists.

Claim 1. *For any two paths $P_{ij}, P_{k\ell} \in \mathcal{P}'$, either $m_{ij} \in B(c_k) \cup B(c_\ell)$ or $m_{k\ell} \in B(c_i) \cup B(c_j)$.*

Proof. Let x be a vertex shared by both P_{ij} and $P_{k\ell}$. Assume, w.l.o.g., that c_i is the closest of the endpoints of P_{ij} and $P_{k\ell}$ ($\{c_i, c_j, c_k, c_\ell\}$) to x . Also assume, w.l.o.g., that x is in the c_k -to- $m_{k\ell}$ subpath of $P_{k\ell}$. Since $m_{k\ell} \in B(c_k)$ the distance from $m_{k\ell}$ to x to c_k is at most R and since c_i is closer to x than c_k , then the distance from $m_{k\ell}$ to x to c_i is also at most R , therefore $m_{k\ell} \in B(c_i)$. \square \square

Since this claim holds for every pair of paths, by an averaging argument, there must be some path P_{ij} whose midpoint is contained in the ball centered at the endpoint of at least $\frac{1}{|\mathcal{P}'|} \cdot \binom{|\mathcal{P}'|}{2} = \frac{1}{2}(|\mathcal{P}'| - 1) \geq q_g - 2$ paths. Since m_{ij} is additionally contained in $B(c_i) \cap B(c_j)$, m_{ij} is a point contained in q_g balls, showing that the ball system for G has the (p_g, q_g) -property.

5 Handling apices and toward minor-excluded graphs

The Graph Minor Structure Theorem is one of many results of Robertson and Seymour leading to the Graph Minor theorem. The Graph Minor Structure Theorem shows that for a fixed graph H , any graph excluding H as a minor is composed of graphs that, after the removal of a fixed number of vertices, can be embedded on a surface in which H cannot be embedded with a fixed number of *vortices* (described below). These subgraphs are glued together in a tree-like structure called a tree decomposition.

We are able to show that we can *remove* the apices, so to speak, of one of these graphs (Section 5.1) and that there is one subgraph within which every other vertex is distance R (Section 5.2). Of course, this subgraph may be quite large, but since this subgraph is nearly embeddable on some surface, after removal of the apices, it may be possible to use arguments similar to those in Section 4. We discuss this further in Section 5.3.

5.1 Removing apices

We show something stronger than that of *removing apices from bounded genus graphs*:

Lemma 6. *Let \mathcal{G} be a class of graphs whose ball systems have VC-dimension at most $q - 1$ and satisfy the (p, q) -property. Then there is a constant ρ such that any graph in \mathcal{G} with an additional α apices and diameter at most $2R$ can be covered by at most $\rho + \alpha$ balls of radius R .*

Proof. Let G be a graph such that for a subset of at most α vertices A , $G \setminus A \in \mathcal{G}$. Let \mathcal{B} be the ball system for G and let \mathcal{B}' be the subset of those balls that do not intersect A . The VC-dimension of \mathcal{B}' is at most that of \mathcal{B} , which is at most $q - 1$. Likewise, since \mathcal{B} has the (p, q) -property, so does \mathcal{B}' . By the Fractional Helly Theorem, it follows that \mathcal{B}' has a hitting set of size at most ρ ; this hitting set along with A is a hitting set for \mathcal{B} . \square \square

5.2 The central node of a tree decomposition

A tree decomposition \mathcal{T} of a graph $G = (V, E)$ is a pair (T, \mathcal{X}) where T is a tree and \mathcal{X} is a family of subsets (or *bags*) of V such that:

- Each node a of T has a corresponding subset $X_a \in \mathcal{X}$ and $\cup_{X \in \mathcal{X}} X = V$;
- For every edge $uv \in E$ there is a bag $X \in \mathcal{X}$ such that $u, v \in X$.
- For any three nodes $a, b, c \in T$ such that b is on the a -to- c path in T , $X_a \cap X_c \subseteq X_b$.

We refer to the *nodes* of T and *vertices* of G to avoid confusion. The width of a tree decomposition (T, \mathcal{X}) is $\max_{X \in \mathcal{X}} |X| - 1$. Tree decompositions are not unique. The treewidth of a graph is the minimum possible width of a tree decomposition of the graph.

We show that given a tree decomposition of a graph of diameter $2R$, there is a node a of the tree decomposition such that every vertex in the graph is within distance R of some vertex in X_a . This is similar to Theorem 5 by Gavaille et al. [8], but we are specific about the node of interest in the tree decomposition. We include the proof below for completeness.

Theorem 6 (Central node). *There is a node v of a tree decomposition $\mathcal{T} = (T, \mathcal{X})$ of a graph G with diameter at most $2R$ such that every vertex of G is within distance R of some vertex in X_v ; i.e. $d(x, X_v) \leq R$ for every vertex x of G .*

Consider a node u of T and the corresponding bag $X_u \in \mathcal{X}$. Removing u from T and X_u from G results in $k \geq 1$ subgraphs, each with a tree decomposition derived from \mathcal{T} . Formally, let T_u^1, \dots, T_u^k be the components of $T \setminus \{u\}$. Let \mathcal{X}_u^j be the bags corresponding to nodes of T_u^j with the vertices in X_u removed: $\mathcal{X}_u^j = \{X_v \setminus X_u : v \in T_u^j\}$. Let V_u^j be the vertices in the bags corresponding to nodes of T_u^j with X_u

removed: $V_u^j = \cup \mathcal{X}_u^i$. $\mathcal{T}_u^j = (T_u^j, \mathcal{X}_u^j)$ is a tree decomposition of the subgraph of G induced by V_u^j . Since X_u is a vertex separator, any v -to- w path in G for $v \in V_u^i$ and $w \in V_u^j$ ($i \neq j$) must contain a vertex of X_u .

Let $d(x, y)$ be the shortest-path distance between x and y in G . For a subset of vertices Y , let $d(x, Y)$ be the minimum distance from x to any vertex of Y , so $d(x, Y) = \min_{y \in Y} d(x, y)$. For any two subsets X and Y , let $f(X, Y)$ be the furthest vertex in X from Y ; i.e. $f(X, Y) = \arg \max_{x \in X} d(x, Y)$.

Lemma 7. *If the distance from the furthest vertex in V_u^i to X_u is greater than R for any i , then for every $j \neq i$, the distance from the furthest vertex in V_u^j is strictly less than R .*

Proof. Let $f_i = f(V_u^i, X_u)$ and let $f_j = f(V_u^j, X_u)$ for $i \neq j$.

Let x be a vertex in X_u that is on a shortest path from f_i to f_j . Note also that since $f(V_u^i, X_u)$ and $f(V_u^j, X_u)$ are both vertices in G , $d(f(V_u^i, X_u), f(V_u^j, X_u)) \leq 2R$. So we have:

$$\begin{aligned} 2R &\geq d(f(V_u^i, X_u), f(V_u^j, X_u)) \\ &= d(f(V_u^i, X_u), x) + d(f(V_u^j, X_u), x) \\ &\geq d(f(V_u^i, X_u), X_u) + d(f(V_u^j, X_u), X_u) \\ &> R + d(f(V_u^j, X_u), X_u) \end{aligned}$$

The above then immediately implies that $d(f(V_u^j, X_u), X_u) < R$. □ □

Consider the following procedure for finding the central node, starting at an arbitrary node r :

```

1  SEARCH( $r$ )
2    If  $d(x, X_r) \leq R$  for all  $x \in V(G)$ , return  $r$ .
3    Otherwise:
4      Let  $p$  be a node adjacent to  $r$  in  $T$  such that  $d(f(V_r^i, X_r), X_r) > R$  and  $p \in T_r^i$ .
5      SEARCH( $p$ ).
```

It is clear that if this procedure terminates, then the statement of the lemma is true. It remains to argue that the algorithm must terminate. If we reach line 4, then, by Lemma 7, p is unique. If SEARCH does not terminate, then it is easy to see that SEARCH must oscillate between two adjacent nodes p and q of the tree decomposition: SEARCH(p) calls SEARCH(q) and vice versa. In this case, there must be a vertex $x \in T_q^i$ where i is such that $d(f(V_q^i, X_q), X_q) > R$ and $p \in T_q^i$ and a vertex $y \in T_p^j$ where j is such that $d(f(V_p^j, X_p), X_p) > R$ and $q \in T_p^j$. Let S be a shortest x -to- y path; by definition of p and q , S must visit a vertex $a \in X_p$ and a vertex $b \in X_q$ (possibly $a = b$). Let m be a vertex closest to the middle of S . Since the diameter of G is at most $2R$, $d(x, m)$ and $d(y, m)$ is at most R . Therefore b must come after m along S from x to y and a must come after m along S from y to x . It must be that $m = a = b$, contradicting that $d(x, X_q) > R$ and $d(y, X_p) > R$. This concludes the proof of the Central Node Theorem.

Theorem 5 of Gavioille et al.'s work is an immediate corollary of Theorem 6:

Corollary 1 (Theorem 5 [8]). *For a graph with treewidth tw and diameter $2R$, there is a set S of at most $tw + 1$ vertices such that $d(x, S) \leq R$ for every vertex x in the graph.*

5.3 Minor-free decompositions

Finally, we outline a direction for extending this result to minor-free graph classes and describe the challenges.

Robertson and Seymour showed that for any graph G_H that excludes a fixed minor H , G_H has a well-defined structure [19]. Using the notation and terminology of Demaine et al. [3], the Graph Minor Structure Theorem states that G_H is obtained by h -clique sums of graphs that are h -almost embeddable on surfaces in which H cannot be embedded. A graph G is h -almost-embeddable on a surface S if:

- There is a set A of at most h vertices, called *apex* vertices, such that $G \setminus A$ can be written as a union of graphs $G_0 \cup G_1 \cup \dots \cup G_h$ where G_0 can be cellularly embedded on S .

- For every $i > 0$, G_i is a graph, called a *vortex*, that has a tree-decomposition that is a path with nodes in order x_i^1, x_i^2, \dots and width at most h .
- For every $i > 0$, there is a face F_i such that u_i^1, u_i^2, \dots is a subset of the boundary vertices of F_i in order along the boundary of F_i and $u_i^j \in X_{x_i^j}$ for all j .

Note that since H is fixed, the surfaces in which the components of G_H are almost embeddable have fixed genus.

An h -clique sum between graphs A and B identifies the vertices of a clique on at most h vertices in A and B and then possibly removes some edges of the clique. The clique-sum of graphs provides a natural tree decomposition. Specifically, G_H admits a tree decomposition (T, \mathcal{X}) such that for every $X \in \mathcal{X}$, the subgraph of G_H induced by X is h -almost embeddable and the intersection of any two sets of \mathcal{X} contains at most h vertices. Using this decomposition, we define the *central subgraph* of G_H as the subgraph of G_H induced by the vertices in the central node of this tree decomposition.

Focussing on this central subgraph, we can *remove* the apices by way of Lemma 6. Now, in the efforts to prove the (p, q) -property for the set of balls not intersecting apices of the central subgraph, consider a set of p balls for sufficiently large p . We can assume w.l.o.g. that at most one ball center is in each of the neighboring H -minor-free graphs that are clique-summed to the central subgraph; if a large number of ball centers are in one neighbor, then since the balls must all reach the central subgraph, a large enough number of them must share a vertex, since the clique sums are small.

We can then focus on center-to-center shortest paths, as in Section 4. For this proof technique, we need to show that among a set of center-to-center shortest paths, a sufficiently large number of them share an interior vertex. While these paths must cross the central subgraph and parts of them must be embedded on the surface that the bulk of the central subgraph is embedded on, these paths can use the clique sums and vortices to *hop* over each other, crossing without intersecting. It does not seem possible to bound how much this can happen since the *number* of vortices and clique sums is not bounded. so it is likely that a more global argument, taking into account the balls and not just the shortest paths between ball centers, will be required in order to illustrate the (p, q) -property.

6 Discussion

This paper presents a generalization of the ball-cover property to bounded genus graphs with a constant number of apices. This represents a significant step towards showing this result holds for all minor-free families of graphs. This work leaves open this direct question and several others.

For one, these results, ours and that of Chepoi et al., do not evaluate the explicit number of balls required for coverage, relying as we do, on the Fractional Helly Theorem. Tracing the constant through Matoušek's work reportedly results in a constant in excess of 800 [23] while the best lower bound known is 4 [8]. A direct proof, bypassing the Fractional Helly Theorem, is likely necessary to result in more practical answers. Likewise, an algorithmic result is desirable, particularly if the application to interval routing is to be taken seriously.

Further, since our planarizing set (Lemma 4) reduces a graph of genus g to a planar graph after the removal of $O(g||G||_\delta)$ edges and since Gazit and Miller give an $O(||G||_\delta)$ balanced edge separator for planar graphs, we can combine these results to get an $O(g||G||_\delta)$ edge separator for genus- g graphs. The obvious question is whether an $O(\sqrt{g}||G||_\delta)$, balanced edge separator exists for genus- g graphs. Much like Gazit and Miller's separator is a strictly tighter bound on size than the pre-existing $O(\sqrt{\delta_{\max}n})$ balanced edge separator for planar graphs [4, 14], an $O(\sqrt{g}||G||_\delta)$, balanced edge separator for genus- g graphs would be a strictly tighter bound. Our implied $O(g||G||_\delta)$ edge separator results in a set of planar graphs, since the procedure starts by planarizing the graph; it is likely that a tighter bound of $O(\sqrt{g}||G||_\delta)$ would not result in a set of planar graphs.

Acknowledgements We thank Anastasios Sidiropoulos and Mark Walsh for helpful discussions. This material is based upon work supported by the National Science Foundation under Grant Nos. CCF-0963921

and CCF-1054779.

References

- [1] P. Agarwal, B. Aronov, J. Pach, R. Pollack, and M. Sharir. Quasi-planar graphs have a linear number of edges. *Combinatorica*, 17(1):1–9, 1997.
- [2] Victor Chepoi, Bertrand Estellon, and Yann Vaxes. Covering planar graphs with a fixed number of balls. *Discrete & Computational Geometry*, 37(2):237–244, 2007.
- [3] Erik D. Demaine, Fedor V. Fomin, MohammadTaghi Hajiaghayi, and Dimitrios M. Thilikos. Subexponential parameterized algorithms on graphs of bounded-genus and h -minor-free graphs. *Journal of the ACM*, 52(6):866–893, 2005.
- [4] K. Diks, H.N. Djidjev, O. Sýkora, and I. Vrt’o. Edge separators of planar and outerplanar graphs with applications. *Journal of Algorithms*, 14(2):258 – 279, 1993.
- [5] J. Edmonds. A combinatorial representation for polyhedral surfaces. *Notices of the American Mathematical Society*, 7:646, 1960.
- [6] Stefan Felsner. Topological graphs: Crossing lemma and applications. In *Geometric Graphs and Arrangements*, Advanced Lectures in Mathematics, pages 43–52. Vieweg+Teubner Verlag, 2004.
- [7] Jacob Fox, János Pach, and Andrew Suk. The number of edges in k -quasi-planar graphs. *SIAM J. Discrete Math.*, 27(1):550–561, 2013.
- [8] Cyril Gavoille, David Peleg, Andr Raspaud, and Eric Sopena. Small k -dominating sets in planar graphs with applications. In Andreas Brandstdt and VanBang Le, editors, *Graph-Theoretic Concepts in Computer Science*, volume 2204 of *Lecture Notes in Computer Science*, pages 201–216. Springer Berlin Heidelberg, 2001.
- [9] Hillel Gazit and Gary L. Miller. Planar separators and the Euclidean norm. In *SIGAL International Symposium on Algorithms*, pages 338–347, Tokyo, August 1990. Information Processing Society of Japan, Springer-Verlag.
- [10] Richard K. Guy, Tom Jenkyns, and Jonathan Schaer. The toroidal crossing number of the complete graph. *Journal of Combinatorial Theory*, 4(4):376 – 390, 1968.
- [11] Allen Hatcher. *Algebraic Topology*. Cambridge Univ. Press, 2002.
- [12] E. Helly. Über mengen konvexer körper mit gemeinschaftlichen punkten. *Jahresbericht der Deutschen Mathematiker-Vereinigung*, 32:175–176, 1923.
- [13] Jir Matoušek. Bounded VC-dimension implies a fractional Helly theorem. *Discrete & Computational Geometry*, 31(2):251–255, 2004.
- [14] G. L. Miller. Finding small simple cycle separators for 2-connected planar graphs. *Journal of Compute and System Sciences*, 32(3):265–279, 1986.
- [15] Bojan Mohar and Carsten Thomassen. *Graphs on Surfaces*, chapter 4. Johns Hopkins University Press, Baltimore, 2001.
- [16] James R. Munkres. *Topology*. Prentice-Hall, 2nd edition, 2000.
- [17] János Pach, Farhad Shahrokhi, and Mario Szegedy. Applications of the crossing number. In *Proceedings of the tenth annual Symposium on Computational geometry*, SCG ’94, pages 198–202, New York, NY, USA, 1994. ACM.

- [18] G. Ringel and J. W. T. Youngs. Solution of the Heawood map-coloring problem. *Proceedings of the National Academy of Sciences of the United States of America*, 60:438–445, 1968.
- [19] Neil Robertson and P.D Seymour. Graph minors. XVI. excluding a non-planar graph. *Journal of Combinatorial Theory, Series B*, 89(1):43 – 76, 2003.
- [20] F. Shahrokhi, O. Sýkora, L.A. Székely, and I. Vrt’o. The crossing number of a graph on a compact 2-manifold. *Advances in Mathematics*, 123(2):105 – 119, 1996.
- [21] F. Shahrokhi, L. Székely, O. Sýkora, and I. Vrt’o. Improved bounds for the crossing numbers on surfaces of genus g . In Jan Leeuwen, editor, *Graph-Theoretic Concepts in Computer Science*, volume 790 of *Lecture Notes in Computer Science*, pages 388–395. Springer Berlin Heidelberg, 1994.
- [22] A. Suk and B. Walczak. New bounds on the maximum number of edges in k -quasi-planar graphs. In *Proc. of the Int’l Symp. on Graph Drawing*, 2013.
- [23] S. Thomassé. Invited talk, canaDAM, University of Victoria, May 2011.
- [24] P. Turán. A note of welcome. *J. Graph Theory*, 1:7–9, 1977.
- [25] V. Vapnik and A. Chervonenkis. On the uniform convergence of relative frequencies of events to their probabilities. *Theory of Probability and its Applications*, 16(2):264–280, 1971.
- [26] J. Youngs. Minimal imbeddings and the genus of a graph. *Journal of Mathematical Mechanics*, 12:303–315, 1963.