

A Pacific Northwest Extension Publication  
Oregon State University • University of Idaho • Washington State University

PNW 625 • March 2011

# Weed and Vegetation Management in Christmas Trees

Ed Peachey, Chal Landgren, and Tim Miller

A good weed management plan is essential to produce high quality, marketable Christmas trees. But, complete control of weeds may not be cost effective and is seldom needed. It's important to strike a balance between reducing weed competition with trees and protecting soil and water.

## Management philosophy

An effective weed management program incorporates diverse strategies and practices to keep weed populations off balance. Using the same practice over and over will, predictably, select for weed species with certain traits. For example, using the same herbicide selects for weeds that aren't affected by that herbicide.

Multiple strategies are essential to prevent weed shifts and herbicide resistance from developing. These strategies include proper identification of important weed species, prevention of new introductions, and use of biological, mechanical, and chemical controls. A weed management plan that integrates mechanical and cultural practices with judicious use of herbicides often gives the best long-term result.

Another important strategy is to begin weed management well before trees are planted. This is particularly important if perennial weeds such as bindweed, bracken fern, or blackberry are present at the site. You can effectively challenge them before the Christmas trees are in the ground. Applying broad-spectrum herbicides and tilling several times over the course of a year can substantially reduce weed density. Trees are sensitive to many broad-spectrum herbicides, so once they are planted, fewer options will be available to control weeds.

## Contents

Management philosophy .....	1
The top dozen weeds: identification and biology ...	2
Annual broadleaves .....	2
Perennial broadleaves .....	4
Biennial .....	6
Annual grasses .....	6
Perennial grasses .....	7
Management Strategies .....	8
Prevention .....	8
Mechanical .....	8
Cover crops and vegetated strips .....	9
Flame weeding .....	10
Biological controls .....	10
Chemical control .....	11
Herbicide application timing and efficacy .....	11
Commonly used herbicides and their characteristics .....	12
Herbicide families and resistance management ...	14
Herbicide application and stewardship .....	15
Non-point losses of herbicides .....	15
Strategies to reduce soil and pesticide loss .....	15
Wildlife and Christmas trees .....	18
Other resources .....	18

---

Ed Peachey, weed management, horticulture crops; Chal Landgren, Extension Christmas tree specialist; both of Oregon State University; and Tim Miller, Extension Horticulture weed specialist, Washington State University. Original authors: Ray William, Ph.D., professor emeritus, Oregon State University; and Kassim Al-Khatib, Ph.D., Washington State University

## The top dozen weeds: identification and biology

Weed control efforts begin by determining which species are present, which weeds will compete with Christmas trees for resources or reduce overall quality, and, ultimately, which weeds must be removed to meet management objectives. This requires proper identification in the field and a general knowledge of the biology and characteristics of the important species present. Some species may not overly compete with the trees for water and nutrients. But, misidentification of a perennial weed may lead to dire consequences in later years.

See *Herbicide Effectiveness on Weeds in Christmas Trees* (see “Other resources,” page 18) for information on control of these species.

### Annual broadleaves

**Horseweed** (*Conyza canadensis*), Sunflower family. Grows tall and erect with a unique inflorescence and tiny yellow and white flowers (Figures 1 and 2). Although native to the region, it is becoming more prevalent in some areas of the Pacific Northwest. It is glyphosate resistant in some areas of the U.S.



Figure 1. Horseweed plant




Figure 2. Horseweed flowers

**Prickly lettuce** (*Lactuca serriola*), Sunflower family. An annual (sometimes biennial) weed that has a milky sap in the stem, yellow flowers (Figure 3), and wind-dispersed seeds (Figure 4). Look for a row of prickles along the mid-vein on the back of each leaf (Figure 5). It is resistant to several sulfonylurea herbicides in some areas east of the Cascades. It is commonly found along roadsides west of the Cascades.



Figure 3. Prickly lettuce flowers



Figure 4. Prickly lettuce seeds




Figure 5. Prickly lettuce leaf

**Spotted catsear** (*Hypochaeris radicata*), Sunflower family. A dandelion-like perennial forming a basal rosette of roughly-hairy leaves (Figure 6). Flowers also are very similar to dandelion, although they are borne in midsummer on a many-branched inflorescence from solid stems (Figure 7). It is extremely common and a prolific seed producer (Figure 8).



Figure 6. Spotted catsear rosette



Figure 7. Spotted catsear flowers



Figure 8. Spotted catsear in field

**Hawksbeard**, bristly and smooth (*Crepis setosa* and *C. capillaris*), Sunflower family. These are annuals that have clasping leaves with abrupt, widely separated teeth along their margins (Figure 9). Plants produce a narrow inflorescence of bright yellow flower heads (Figure 10), slightly smaller than dandelion heads.



Figure 9. Hawksbeard leaves



Figure 10. Hawksbeard flowers

### Perennial broadleaves

**Willowherbs** (*Epilobium* spp.), Evening-primrose family. Several species, including annuals (*E. brachycarpum*) and perennials (*E. ciliatum*). Basal leaves and stems are often deep red, particularly in winter and early spring (Figure 11). Flowers are small and pink, and followed by tiny seeds bearing threadlike plumes. Christmas trees can provide a niche for this plant (Figure 12).



Figure 11. Willowherbs, leaves and stems



Figure 12. Niche for willowherbs

**Bindweed**, field and hedge (*Convolvulus arvensis* and *Calystegia sepium*), Morning-glory family. This plant grows twining vines from deep, perennial roots. Leaves are arrowhead- to heart-shape, from about an inch (field bindweed) to 3 inches (hedge bindweed) long. Flowers of both species are funnel-shape. Field bindweed flowers are white to pinkish, and about an inch wide (Figure 13a). Hedge bindweed flowers are larger and white (Figure 14). There are two bracts on the stem below the field bindweed flower (Figure 13b), while hedge bindweed has large clasping bracts that cover the sepals (Figure 14). Both species produce seed that can survive up to 20 years.



Figure 13a.



Figure 13b.



Figure 14.

**Canada thistle** (*Cirsium arvense*), Sunflower family. A perennial weed arising from an extensive root system. Leaves are intensely spiny and shiny with only occasional hairs (Figures 15 and 16 on left). Stems are slender and bear fragrant purple flower heads near the tip. Heads are about the size of a dime and not particularly spiny. Flowers are followed by down-tipped seeds. Bull thistle (an annual) also has purple flowers, but the flower head is much larger and leaves are not shiny (Figure 16 on right). Canada thistle is a perennial, and emerging shoots (Figure 15) are often clumped next to each other. Bull thistle arises from a seed, and plants are often isolated.



Figure 15. Canada thistle



Figure 16. Canada thistle (left); bull thistle (right)

**Blackberry** (*Rubus* spp.), Rose family. These are several species of perennials with biennial stems. The native blackberry is a trailing vine. Introduced species are upright canes with leaves that have five leaflets (Figure 17). All are heavily spiny and easily identified by their large white flowers (Figure 18) and edible black fruits.



Figure 17. Blackberry leaves



Figure 18. Blackberry flower

### Biennial

**Wild carrot** (*Daucus carota*), Carrot family. Biennial plants with lacy foliage (Figure 19) and a distinct carrot odor. First-year plants form a rosette. Second-year plants bear umbels (Figure 20) about 4 inches wide that bear dozens of tiny, five-petal white flowers and often a single dark red or purple flower in the center.



Figure 19. Wild carrot foliage



Figure 20. Wild carrot umbels

### Annual grasses

**Barnyardgrass** (*Echinochloa crus-galli*), Grass family. Seeds usually germinate in late spring after soils warm up. Plants are often prostrate unless forced upward by competition with other plants. Stems are flattened and red near the base (Figure 21). Leaves are hairless, and seeds are borne on multiple racemes that are often brown or red at maturity (Figure 22).



Figure 21. Barnyardgrass stems.



Figure 22. Barnyardgrass seeds

**Rattail fescue** (*Vulpia myuros*), Grass family. Seeds normally germinate after onset of rain in the fall. Leaves are tightly rolled and borne in tight bunches. Plants flower and seed in mid-summer, when they have a feathery appearance (Figures 23 and 24).



Figure 23. Rattail fescue



Figure 24. Rattail fescue

**Witchgrass** (*Panicum capillare*), Grass family. Seeds often germinate in mid-spring. First leaves are arching and hairy, and bear a superficial resemblance to corn, wild proso millet, and barnyardgrass. Plants bear diffuse panicles of very tiny seeds borne at the ends of tiny branches (Figures 25 and 26).



Figure 25. Witchgrass



Figure 26. Witchgrass seed

### Perennial grasses

**Bentgrasses** (*Agrostis* spp.), Grass family. These are several species of perennial grasses that normally spread by stolons on the surface of the soil. Plants usually grow year-round in coastal areas but may be dormant in the interior Pacific Northwest. Plants are prostrate, although flower stalks are often upright. Seeds are very small and borne in diffuse panicles (Figure 27).



Figure 27. Bentgrass

Photo by Brian McDonald, Department of Horticulture. © Oregon State University

## Management Strategies

### Prevention

Regulation of weed seed production is the key to successful prevention. Many annual weeds spend most of their time underground as seeds, emerging only briefly to capture sunlight and make more seeds. Seed dormancy regulates the timing of weed emergence and may keep weed seeds alive in the soil for many years.

Weeds are often moved from site to site on equipment or clothing. We can thank birds, mice, wind, and—all too frequently—ourselves for movement of seeds into our fields and cropland.

Preventing introduction of weed seeds requires:

- Planning, to ensure clean equipment
- Maintaining equipment yards free of weeds
- Controlling weeds along farm roads

Growers are also advised to eliminate new weed species before they have the chance to set seed or establish.

The recent increase in puncturevine (*Tribulus terrestris*) (Figure 28) sightings in the Willamette Valley is a good case study in seed movement. The fruits of puncturevine (Figure 29) stick to tires and



Figure 28. Puncturevine plant



Figure 29. Puncturevine fruit (bur) with 5 attached nutlets that each hold 2–3 seeds

shoes, allowing it to spread easily and sometimes great distances. Once this weed is established, it is very difficult to eradicate because of its very long seed dormancy.

Herbicides can be extremely effective tools, but it is important to remember that herbicides do not influence how long a weed seed survives in the soil or how long it remains dormant. Herbicides only act on germinating seedlings or after plants emerge from the soil.

### Mechanical

Cultivation and mowing can be extremely helpful in preventing a shift to species that are tolerant of herbicides or other practices. Be sure to assess the benefits of cultivation or mowing within existing fields relative to potential root or tree damage. Shields around tires and implements help minimize aboveground damage.

Small seedlings of many weeds are easily controlled with cultivation. But, all practices have a weakness, and overuse of cultivation will increase cultivation-resistant species such as common purslane (*Portulaca oleracea*). Cultivation can also spread perennial weeds such as field bindweed or Canada thistle as small pieces of roots are carried through a field.

Cultivation equipment used in Christmas trees includes narrow rototillers, small disks, harrows, and a wide assortment of tools designed to cut weeds off just under the soil surface. Most weed seeds emerge from the top  $\frac{3}{4}$  inch of soil and seldom emerge from below 2 inches (depending on seed size). It is important not to disturb the soil any deeper than necessary, to avoid aggravation of soil erosion in the fall.

Cultivation (rototilling or light disking) may be a weed control option for the first 2 or 3 years after planting; but, as trees fill the space between rows, cultivation becomes less effective. Moreover, cultivation in later years may damage tree roots.

Timely mowing can also significantly reduce competition with the crop. Mowing prior to weed seed development may reduce the number of seeds produced by a plant. Annual broadleaves are particularly susceptible to mowing; grasses and perennial species are tolerant.



## Cover crops and vegetated strips

Cover crops are used in many cropping systems to suppress weed emergence. Suppression is achieved either through direct competition from a living crop or through mulch layers that remain on the soil surface after the cover crop is killed. A cover crop or permanent grass strip (**perm**) that is well designed and planted at the right time may suppress many winter annuals. The main drawback of cover crops is that they consume water: if they are not carefully regulated, they will reduce tree growth.

### Cover crops

There are many ways to establish cover crops. Some growers have found it easier to establish the cover during site preparation. Before planting the trees, they spray the rows where the trees will be planted to kill the cover crop. To establish cover crops after trees are planted, use narrow seed drills or broadcast seed at high rates and lightly till the soil to cover the seed. This method works best with smaller-seeded cover crops.

In the first 1 to 3 years after tree planting, plant annual cover crops in the fall that are easy to regulate in the spring. Their main function is to stabilize soil and suppress weeds. These covers must be killed with herbicides or cultivation and kept from reseeding before the soil dries in the spring (Figure 30), or they will significantly reduce available water later in the summer. Cereal annual cover crops such as spring type oats, barley, or wheat are preferred because they may winter-kill, grow faster in the fall than winter types, can easily be removed with grass herbicides in the spring if needed (with little risk of injury to trees), and can be killed with mowing after the cereal flowers.



Figure 30. Cover crop kept from reseeding before spring.

Some cover crops, such as spring barley and spring oats, will winter-kill west of the Cascades. However, if planted too late in the fall (after November 1), they may survive the winter without too much damage. Cover crops such as sudangrass will always winter-kill, but they usually need to be planted by the end of August to have enough heat to produce cover. There is seldom enough soil moisture at this time to germinate sudangrass, but depending on rainfall patterns for a particular year, sudangrass may be a viable option.

Planting time of cover crops influences the kind of weed suppression achieved. Spring cereals emerge quickly (compared to winter cereals of rye, winter wheat, or perennial grasses) and can out-compete small-seeded weeds, but only if they are planted before weed seeds germinate and emerge. Cereal cover crops should be drilled before the first rains of fall, so that they get a head start on weeds. Small-seeded perennial grasses emerge very slowly and provide little in the form of weed competition, but they will provide soil cover until harvest.

An important benefit of cover crops is that they reduce the pounding action of rainfall that loosens soil and allows it to erode. The method of seeding cover crops in the fall may have a large impact on whether the cover crop will stabilize soils. Tillage in the fall and conventional seeding are not recommended on slopes of more than 2 or 3 percent, because they may result in excessive erosion. Instead, use direct-seed orchard drills that reduce the amount of soil disturbance during planting.

Typical opener spacings on orchard drills are 7.5 to 8 inches. Removing half of the openers significantly reduces the amount of soil that is loosened during planting. It also reduces the amount of soil coverage provided by the cover crop early on, but an aggressive cover crop soon provides good soil coverage if the crop is planted early enough in the fall.

### Vegetated strips

Permanent grass strips (perms) can be established after 3 or 4 years. These may significantly reduce the number of annual and perennial weeds (Figure 31, page 10). Permanent strips provide a stable soil surface in the winter for harvest.

However, the area dedicated to perennial grasses must be carefully regulated to prevent over-competition. Data from several years of experiments

indicate that even small areas with cover crops will suppress tree growth. Physical or chemical mowing of cover crops can keep aboveground biomass in check and reduce the number of weeds in vegetative strips planted between rows.



Figure 31. Permanent grass strips (perms)

### Flame weeding

Flame weeding is occasionally used in Christmas trees but should be applied with caution. Flaming in late afternoon or when sunlight is bright and air temperatures high provides the most effective results. However, avoid open flame weeding in the summer when senescing weeds and tree trimmings may provide enough fuel to ignite the entire field. Alternatives to open flame weeders are infrared heaters and steam weeders that use propane to generate heat. The heat is applied indirectly to the weeds.

Flame weeding will slow growth or kill some broadleaved weeds, but it only removes the tops of grasses and perennial weeds for a short time. For

that reason, weed populations will shift to grasses rather quickly.

Applicators typically are overzealous when applying flame to weeds and burn the weeds to the ground. This wastes money and time. Only enough flame should be applied so that when a leaf is pressed between thumb and forefinger, the thumbprint is visible. This slight change in the leaf is hard to discern just by looking from a standing position, but it becomes evident within 24 hours. Simple experimentation will tell applicators the amount and intensity of heat that is needed to effectively kill weeds.

A major challenge to flame weeding in Christmas trees is that there are many weeds at different growth stages. Flame weeding is most effective when weeds are very small, usually at the 2- to 3-leaf stage. Moreover, because the time to kill winter annuals is in the spring when air temperatures are low and humidity and soil moisture are high, flame weeding in spring is more energy intensive.

An advantage of flame weeding is that the soil is not disturbed.

### Biological control

There are several biological controls that can be useful in Christmas trees, including a flea beetle and the cinnabar moth for tansy ragwort (*Senecio jacobaea*), the Klamath beetle for St.-Johnswort (*Hypericum perforatum*), two species of puncturevine weevils, and a species of mite that attacks field bindweed. See Table 1 for others. Establishment of the puncturevine weevils and the

Table 1. Target weeds and their biological control agents.

Target weed	Biological control agent	Role	Distribution	Control
St.-Johnswort	<i>Agrilus hyperici</i>	root boring beetle	Low	Good
	<i>Aplocera plagiata</i>	defoliating moth	Wide	Fair
	<i>Chrysolina hyperici</i>	defoliating beetle	Wide	Excellent
	<i>Chrysolina quadrigemina</i>	defoliating beetle	Wide	Excellent
Tansy ragwort	<i>Botanophila seneciella</i>	seed head fly	Wide	Good
	<i>Longitarsus jacobaeae</i>	root/defoliating flea beetle	Wide	Good
	<i>Tyria jacobaeae</i>	defoliating moth	Wide	Good
Puncturevine	<i>Microlarinus lareynii</i>	seed weevil	Low	Poor to good in some areas east of Cascades
	<i>Microlarinus lypriformis</i>	stem boring weevil	Low	Poor to good in some area east of Cascades
Bindweed, field	<i>Aceria malherbae</i>	bud/leaf gall mite	Low	Fair

bindweed mites is unpredictable, as they have not established consistently in the Pacific Northwest. **Inundative or mass releases of these species are important each year.**



© Eric Coombs, Oregon Department of Agriculture

Figure 32. *Chrysolina quadrigemina* on St.-Johnswort

## Chemical control

Herbicides can be **soil active** (killing germinating seedlings) or **foliar active** (killing plants when sprayed on foliage).

**Preemergence (PRE) herbicides** are soil active. They must be applied so that they are present in the soil when seeds begin to germinate or when seedlings are small. PRE herbicides must be activated, usually with rainfall or tillage, so that the herbicide is in the soil where the seeds are germinating—usually in the top inch or two of soil. PRE herbicides do not affect germination itself; they stop growth of the radical or shoot of the seedling as it emerges from the seed. Examples include simazine (Princep, Simazine 4L) and Surflan AS (oryzalin).

In contrast, **postemergence (POST) herbicides** such as Roundup (glyphosate), Finale (glufosinate), 2,4-D, and Fusilade (fluazifop) kill leaves and shoots, have little or no soil activity, and do not affect weed emergence. Postemergence herbicides are primarily absorbed through the treated leaf surface and kill by causing membranes to rupture, by arresting manufacture of essential compounds such as amino acids or fatty acids, or by distorting growth through hormone mimicry.

Postemergence herbicides can be classified as either **contact** or **translocated** herbicides. Some POST herbicides, such as Roundup, 2,4-D, and Stinger, **translocate** or move throughout the plant and can kill underground plant structures. Other

herbicides only kill the tissues that they **contact** above ground.

A third class includes herbicides that have **both PRE (soil) and POST (contact)** activity. These include atrazine, Velpar, and SureGuard. Note that SureGuard's postemergent activity is weak but improves slightly if applied with crop oil concentrate. Herbicides in this class can be applied within a larger time frame and can be effective even as some of the target weeds are emerging.

## Herbicide application timing and efficacy

Weed control usually is poor if PRE herbicides are applied after seed germination and shoots or roots emerging from the seed have grown out of the treated zone of soil. This can happen if herbicides are applied too late in the fall or too early in the spring, depending on the herbicide used.

For example, poor control may result if Velpar is applied too early in the spring. In a recent trial conducted in the Willamette Valley, Velpar was applied at three dates in the spring. Velpar applied too early (April 26) simply ran out of gas compared to the application on May 27 (see Figure 33). This was likely due to the excessive rainfall and cooler temperatures that spring that delayed germination of many weeds, and probably caused the soluble Velpar to leach below the germination zone. Weed control with Westar (hexazinone + sulfometuron) remained consistent whether applied on April 26 or May 27, even though the active ingredient used in Velpar is also present in Westar. Sulfometuron, however, is not prone to leaching, and it persisted long enough provide good weed control. Note that directed spray applications must be used if Velpar is applied after budbreak.




Figure 33. Increasing Velpar activity as application timing is delayed in the spring. Weed control was evaluated mid-June, 2010 ( $\pm$ SE).

The level of weed control resulting from applications of POST herbicides is also affected by stage of growth. Annual weeds typically are controlled best when herbicides are applied to small seedlings. Most perennial species are more easily killed if applications of foliar-active herbicides are delayed until weeds have shoots at least 12 inches long. This allows for greater herbicide uptake (a plant with more or larger leaves receives more herbicide contact) and better translocation of the herbicide from the leaves to the roots. Contact herbicides do not effectively control established perennial species unless applied repeatedly over a long period of time.

Adding surfactants may improve weed control with herbicides (such as Sureguard) that are soil active but also kill foliage. Using the wrong surfactant, however, may expose the Christmas tree to risk of injury. As always, follow labels when using herbicides.

#### Use pesticides safely!

- Wear protective clothing and safety devices as recommended on the label. Bathe or shower after each use.
- Read the pesticide label—even if you've used the pesticide before. Follow closely the instructions on the label (and any other directions you have).
- Be cautious when you apply pesticides. Know your legal responsibility as a pesticide applicator. You may be liable for injury or damage resulting from pesticide use.

### Commonly used herbicides and their characteristics

Refer to Table 2 (page 14) for mode of action of herbicide groups referenced below.

#### Preemergence (PRE) activity

*Dichlobenil* (Casoron), Group 20. Used to control difficult perennials such as Canada thistle and horsetail (*Equisetum*). Apply in midwinter immediately before a cold rain to reduce volatility and enhance weed suppression. Use only around well-established plants; typically, wait 1 year after transplanting. Do not use on light, sandy soils. REI (re-entry interval) 12 hr. Groundwater hazard.

*Indaziflam* (Specticle), Group 29. Apply before weed emergence in the fall or spring. Apply as a directed spray to soil surface. Do not mix into soil. May injure new leaf tissue if applied over the top

of trees. Controls grasses, sedges, and broadleaves. Needs 1/8 to 1/4 inch rain or irrigation water for activation. Does not control emerged weeds. Do not apply to newly transplanted crops until soil settles. This is a new registration (2011), and there is very little experience with this herbicide in the Pacific Northwest.

*Oryzalin* (Surflan A.S.), Group 3. Apply in pine and fir (but not Douglas-fir) Christmas tree plantations, to bare soil surface after transplanting and settling. Activate with 1/2 inch water or shallow cultivation before weeds emerge. REI 24 hr. Toxic to fish.

*Pendimethalin* (Pendulum 3.3 EC), Group 3. Apply at planting or before weed seeds germinate. Soil should be loose and free from all established weeds. Water within a few days to activate herbicide before weeds emerge. REI 24 hr. Toxic to fish.

*Pronamide* (Kerb 50W), Group 3. Make a single application in fall, before freezing weather. Not recommended on trees less than 1 year old. Use higher rates for quackgrass and finer soils. Requires soil moisture to activate. Primarily for grass control. Degraded by microorganisms in warmer weather. REI 24 hr. Restricted use.

*Simazine*, Group 5. Apply in fall or spring. Simazine must be applied before weeds emerge. Avoid applying over actively growing trees unless applied immediately before or during a rain. REI 48 hr. Groundwater hazard.

#### Preemergence and postemergence activity

*Atrazine* (many products), Group 5. To ensure activation, apply in March or April after soil settles from transplanting but before rains stop. Avoid applying over actively growing trees, unless applied immediately before or during a rain. Do not exceed 4 lb ai/A per year. REI 12 hr. Groundwater hazard.

*Flumioxazin* (SureGuard), Group 14. Preemergence and/or postemergence herbicide for control of selected grass and broadleaf weeds. Post-emergence weed control is limited to very small weeds. All over-the-top applications of SureGuard are applied prior to spring budbreak or delayed until conifers have hardened off. REI 12 hr. Toxic to aquatic invertebrates.

*Hexazinone* (Velpar DF), Group 5. Registered for new plantings, but variable tree injury has been observed. Applications in early April may be more effective than March applications due to greater leaching from spring rains. Higher rates are effective on trailing blackberries. REI 48 hr. Groundwater hazard.

*Hexazinone + sulfometuron* (Westar DG), Groups 5 and 2. Apply either before or soon after weeds emerge, when weeds are small and actively growing. Apply broadcast only to dormant trees. If trees have broken dormancy, directed applications are required to avoid contact with new growth. Recommended for Douglas-fir, Fraser fir, grand fir, noble fir, Nordman fir, and Turkish fir. Check current label for reduced rates on some firs. Surfactants are not recommended and may cause excessive injury. REI 48 hr. Groundwater hazard and long soil residual.

*Metribuzin + flufenacet* (Axiom DF), Groups 5 and 15. For use on Douglas-fir and true firs only. Apply pre- or very early postemergence to weeds when trees are completely dormant. Trees must be established at least one growing season. REI 12 hr. Groundwater hazard.

*Oxyfluorfen* (Goal 2XL), Group 14. Apply immediately after transplanting but before budbreak, over tree tops or as directed spray. Controls small, emerged broadleaf weeds with brief soil residual. Perennial emerged annual grasses are not controlled, although fine fescues may be suppressed. REI 6 days. Toxic to aquatic invertebrates.

### **Postemergence activity**

*2,4-D*, Group 4. Apply over-the-top as a broadcast treatment when trees are dormant using a low rate, or direct spot sprays toward actively growing weeds before budbreak or after budset during cool weather. Do not apply over the tops of pine or true firs (*Abies* spp.) REI 48 hr. May be toxic to fish.

*Asulam* (Asulox), Group 18. For bracken fern control. Apply after Christmas tree terminal buds mature and bracken fronds are fully expanded. REI 12 hr. Groundwater hazard.

*Clethodim* (Envoy), Group 1. For Douglas-fir, grand fir, and pines. Apply postemergence to actively

growing annual or perennial grasses as listed on label. REI 24 hr.

*Clopyralid* (Stinger), Group 4. Apply broadcast or in bands during active weed growth up to five-leaf stage. Controls annual weeds of the sunflower family, Canada thistle, clover, and knapweeds. To avoid needle curling, do not apply during first year after transplanting. REI 12 hr. Groundwater hazard.

*Fluazifop* (Fusilade DX), Group 1. Apply to actively growing grasses in early spring, following ample rain, as a directed spray with 0.25 percent nonionic surfactant. Results often are erratic on grasses stressed from lack of vigor, drought, high temperature, or low fertility. More mature grasses and quackgrass can be controlled but may require two applications. Annual bluegrass and all fine fescues resist treatment. REI 12 hr.

*Glufosinate* (Finale), Group 10. Apply to actively growing weeds. Do not let spray or drift contact living tissue or green, thin, or uncalloused bark, as injury may occur. Do not broadcast over Christmas trees. REI 12 hr. Groundwater hazard.

*Glyphosate* (numerous product names), Group 9. Select application equipment to prevent crop injury by directing spray toward base of plants or with selective applicators. Consult labels about rate and time of application (especially for perennial weeds), and to determine which formulations allow over-the-top applications. Avoid spray or mist contacting foliage or green bark of desirable plants. Apply with a wiper at 33 percent solution. REI 4 hr.

*Sethoxydim* (Segment), Group 1. Identify susceptible grasses. Control often is erratic on grasses stunted or stressed from drought, high temperatures, or low fertility. Resistant grasses include annual bluegrass and all fine fescues. Quackgrass can be suppressed. REI 12 hr.

*Triclopyr* (Garlon 3A), Group 4. Apply in late summer or early fall after terminal growth of trees has hardened off, but before target weeds drop leaves. For control of woody plants and perennial and broadleaf weeds. Select application equipment to prevent tree injury by directing spray toward base of plants or with selective applicators. REI 48 hr. Groundwater hazard.

**Table 2. Herbicide mode of action and characteristics.**

Herbicide group number <sup>a</sup>	Mode of action	Common herbicides labeled in Christmas trees	Number of species with documented resistance worldwide <sup>d</sup>
1	ACCCase inhibitor	Poast, Select, Fusilade	39
2	ALS inhibitor	Westar <sup>b</sup>	107
3	Cell division inhibitor	Surflan, Pendulum	10
4	Auxin inhibitor	Stinger, Garlon	28
5	Photosystem II inhibitor	Atrazine, Simazine, Axiom <sup>c</sup> , Westar <sup>b</sup>	68
9	ESPS inhibitor	Roundup	21
10	Glutamine synthesis inhibitors	Finale	0
14	PPO inhibitor	Goal, SureGuard	4
15	Inhibits very long chain fatty acid synthesis	Axiom <sup>c</sup>	1
18	Inhibits DHP synthase step	Asulox	0
20	Inhibits cell wall synthesis	Casoron	1

<sup>a</sup> WSSA herbicide group name designation.

<sup>b</sup> Westar is two products; sulfometuron plus hexazinone. Sulfometuron is in Group 2 and hexazinone in Group 5.

<sup>c</sup> Axiom is two products: metribuzin plus flufenacet. Metribuzin is in Group 5 and flufenacet is in Group 15.

<sup>d</sup> From the International Survey of Resistant Weeds <http://www.weedscience.org/In.asp>.

### Herbicide families and resistance management

Continued use of a single herbicide selects weed species that are tolerant to that herbicide. This is called a **species shift**. Species shift can often be corrected simply by switching to a different herbicide or other control strategy (such as tillage or mowing).

Continued use of a single herbicide over time may also result in a weed population that is **resistant** to that herbicide. This occurs when a species that is usually controlled by a particular herbicide no longer responds to that herbicide, and usually results from genetic differences among individual weeds within the population. Seedlings from resistant plants usually are resistant to herbicides employing the same mode of action (the same group; see Table 2). Herbicide resistance is not reversed if the herbicide is no longer applied—once selected for, the resistant trait remains in the weed population on your farm.

The best strategy to prevent selecting for resistant weeds is to:

1. Integrate herbicides with other weed control practices.
2. Use herbicides that have different modes of action in rotation or in combination/sequence with each other.

Table 2 lists herbicides commonly used in Christmas trees and their herbicide activity group.

Some of these herbicides select for resistant species very quickly (Group 2); others are very slow to select for resistant species.

### Preventing herbicide-resistant weeds

1. Use herbicide rotations. Do not apply the same herbicides year after year. Rotate among herbicide groups with different sites of action.
2. Use short-residual herbicides (herbicides that do not persist in soil for long periods). Using short residual herbicides that are not applied repeatedly within a growing season reduces the selection of herbicide-resistant weeds.
3. Use crop rotation. Each crop has a unique suite of practices used to produce it, including herbicide used. Crop rotation usually diversifies herbicides and other practices used to control weeds. This may not be practical on most Christmas tree farms, however.
4. Use integrated practices and non-chemical methods to manage weeds. Cultivation, for example, can be an effective tool for eliminating weed escapes that may represent the resistant population.
5. Monitor fields and record weed escapes after herbicides are applied. Do not let weeds go to seed or even produce pollen, as pollen often carries a dominant resistance gene.

6. Do not bring resistant weed seeds into your fields via equipment. Clean equipment thoroughly when moving from field to field.
7. If using glyphosate, use the higher labeled rates. In weeds such as annual ryegrass, in which resistance may be polygenic (determined by a number of genes), low glyphosate rates may select for resistance more quickly. Rotate to Finale for postemergence weed control.

### Herbicide application and stewardship

Apply the correct amount of herbicide at the right time and under the right conditions to optimize weed control. Herbicides can be lost through drift during application, through runoff in water, or on soil carried in runoff water. This increases the cost of weed control by reducing efficacy, and may impact non-target plants or animals.

Sprayer calibration is essential to proper use of herbicides. Correct nozzle size and pressure reduce the production of small droplets that are more likely to drift. For details, see the OSU Extension Service video “Calibrating and Using Backpack Sprayers” (see “Other resources,” page 18). The video demonstrates how to use a backpack sprayer safely and effectively, and shows basic sprayer components, calibration methods, pesticide mixing, and helpful operating tips.

Use shielded sprayers to protect small tree seedlings from herbicides (Figure 34).

### Non-point losses of herbicides

A number of pesticides, including herbicides, are routinely identified in the main rivers of the Pacific Northwest. Concerns include drinking water contamination and damage to aquatic ecosystem health, even at very low pesticide concentrations.

Increasingly, buffers are required along many waterways and around field edges to provide some level of



Figure 34. Shielded sprayer

containment for soil and water movement. Herbicides applied in the fall are particularly susceptible to non-point or off-site loss to surface or groundwater during winter rains. Growers should be aware of the toxicity of the products they use and the chances that a pesticide may move off-site. Pesticide labels include information on non-target effects of pesticides.

The propensity of an herbicide to reach surface or groundwater is dependent on several factors. If the herbicide is tightly adsorbed to soil (sticks to soil), it will stay in place unless the soil moves. If the herbicide is not tightly adsorbed (that is, it is easily removed from soil particles), it is at risk of running off the surface with water or percolating into groundwater. Slope of the site, soil type, crop residues on the soil surface, and uncropped buffers around fields, as well as herbicide solubility and application, all play a role in how much pesticide may reach streams and rivers.

### Strategies to reduce soil and pesticide loss

Use sustainable weed management practices to prevent soil loss and reduce contamination of water with herbicides or nutrients. Christmas tree growers use a number of innovative strategies to reduce non-point pesticide loss, including:

- Vegetated filter strips (VFS). If the herbicide is tightly bound to soil, VFS may work well. VFS work best if they are well maintained and designed to prevent channelized flow. Wider filter strips are generally needed to prevent loss




Figure 35. Effect of filter strip width on pesticide load reduction of moderately adsorbed compounds ( $K_{oc} < 1000$ ) in surface runoff. Data summarizes results of 217 studies. Adapted from Reichenberger et al., 1997.

of moderately soluble herbicides such as atrazine (Figure 35, page 15).

- Transient cover crops
- Living mulches
- Catch basins
- Temporary filter strips and straw ropes or barriers to slow water movement

- Managing weedy vegetation by allowing some weeds in strategic places to survive.

Most herbicide labels carry cautions about the potential of herbicides to enter groundwater or streams and rivers and affect fish or other aquatic life. Buffers and setbacks may be recommended for some herbicides but are not required. Buffer strips

**Strategies to stabilize soil and mitigate runoff: native planting of wild blue rye (Figure 36), resident weedy vegetation (Figure 37), fall-planted spring oats suppressed with atrazine and glyphosate in the spring (Figure 38), interrow fescue (Figure 39), and straw ropes to slow runoff and catch soil sediment (Figures 40 and 41).**



Figure 36



Figure 37



Figure 38



Figure 39



Figure 40



Figure 41



are required for atrazine and simazine and should be vegetated if on highly erodible soil. The label for SureGuard recommends installation of filter

strips along rivers, streams, and creeks, and on the downhill side of fields where runoff may occur. See Table 3.

**Table 3. Buffer requirements for currently labeled herbicides in Christmas trees.**

Herbicides	Contamination potential	Buffer requirements or advisories
<b>Preemergence (PRE) with activity</b>		
Dichlobenil (Casoron)	Groundwater hazard	None <sup>a</sup>
Pendimethalin (Pendulum 3.3 EC), Group 3	Toxic to fish	None <sup>a</sup>
Pronamide (Kerb 50W), Group 3	Restricted use	None
Simazine, Group 5	Groundwater hazard	Do not apply within 66 ft of field where surface water runoff enters perennial or intermittent streams and rivers, or within 200 ft of natural or impounded lakes. If applied to highly erodible land, the buffer must be seed planted to crop or seeded with grass.
<b>Preemergence (PRE) and postemergence (POST) activity</b>		
Atrazine (many products), Group 5	Groundwater hazard	Do not apply within 66 ft of field where surface water runoff enters perennial or intermittent streams and rivers, or within 200 ft of natural or impounded lakes. If applied to highly erodible land, the buffer must be seed planted to crop or seeded with grass.
Flufenacet + metribuzin (Axiom DF), Group 15	Groundwater hazard	None
Flumioxazin (SureGuard), Group 14	Toxic to aquatic invertebrates	Label recommends use of filter strips along rivers, streams, and creeks, and on the downhill side of fields where runoff may occur.
Hexazinone (Velpar DF), Group 5	Groundwater hazard	None
Hexazinone + sulfometuron (Westar DG), Groups 5 and 2	Groundwater hazard and long soil residual	None
Oxyfluorfen (Goal 2XL), Group 14	Toxic to aquatic invertebrates	A 25-foot buffer strip is required between treated areas and aquatic habitats such as lakes, reservoirs, rivers, sloughs, ponds, creeks, marshes, streams, and wetlands.
<b>Postemergence (POST) activity</b>		
2,4-D, Group 4	May be toxic to fish. Do not apply to water or where surface water is present.	None <sup>a</sup>
Asulam (Asulox), Group 18	Groundwater hazard	None
Clethodim (Envoy), Group 1	—	None
Clopyralid (Stinger), Group 4	Groundwater hazard	None
Fluazifop (Fusilade DX), Group 1	Toxic to fish and aquatic invertebrates. Groundwater and surface water advisory	A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of fluazifop-p-butyl from runoff water and sediment.
Glufosinate (Finale), Group 10	Groundwater hazard	None
Glyphosate (numerous product names), Group 9	—	None
Sethoxydim (Segment ), Group 1	—	None
Triclopyr (Garlon 3A), Group 4	Groundwater hazard	None <sup>a</sup>

<sup>a</sup> Consult most recent label for buffer requirements for these herbicides. Buffer requirements are likely to change when the National Marine Fisheries Service (NMFS) issues their biological opinion on use of these herbicides in Christmas trees.

## Wildlife and Christmas trees

Implications of weed control for wildlife are highly variable and often site specific. The range of grower responses to individual species runs the gamut from encouraging, to coexisting, to tolerating, to exclusion. Choice of weed control for a farm depends on a host of factors, including:

- The species of wildlife of concern or interest
- The amount and type of habitat change as a result of weed control
- Perspective of the grower relative to wildlife; or, conversely, tolerance of the wild species to the grower
- Interactions between the Christmas tree operation and surrounding landscape
- Damage to trees from deer and/or elk. Damage includes:
  - Antler rubbing along the stem, especially with true fir species. Damage can range from a few injured branches and buds to damaged cambium tissue, which causes most or all of the upper part of the tree to die.
  - Foliage feeding on Douglas-fir and Nordman fir. This is more common in winter and spring.

If your weed control strategy encourages the growth of palatable vegetation (such as clover or herbs), these plants may actually attract elk or deer to the field. In most cases, tree foliage is not a preferred food source; but, if there is little else to eat, then foliage may be “on the menu.” Control typically involves fencing, enclosures, or repellents.

Perching birds may damage newly elongating leaders, especially on taller trees that provide the best vantage points. Weed control levels may influence food sources or habitat for prey. For example, weedy fields provide mice and voles cover from owls and hawks. Similarly, ground-nesting birds are attracted to the weedy cover.

In most situations, growers accept a certain amount of damage from wildlife in the area. Some farms also have ponds, streams, or field edges that encourage some species to move into the field.



**Figure 42. Choice of weed control strategy for wildlife depends on the grower's perspective.**

## Other resources

Calibrating and Using Backpack Sprayers. 1994. Oregon State University Extension Service. <http://oregonstate.edu/media/cchl/vf>

DeFrancesco, J. and K. Murray. 2009. PMSP for Christmas Trees in Oregon, Washington, and Idaho. <http://www.ipmcenters.org/pmsp/pdf/or-wa-idchristmastreespmsp.pdf>

Landgren, C., R. Fletcher, M. Bondi, D. Barney, and R. Mahoney. 2003. Growing Christmas trees in the Pacific Northwest (PNW 6). Oregon State University Extension Service.

Pacific Northwest Weed Management Handbook. 2011. Oregon State University Extension Service. <http://pnwhandbooks.org/weed/>

Peachey, Ed. Herbicide Effectiveness on Weeds in Christmas Trees. [http://hort.oregonstate.edu/files/Herbicide\\_Effectiveness\\_on\\_Weeds\\_of\\_Christmas\\_Trees.pdf](http://hort.oregonstate.edu/files/Herbicide_Effectiveness_on_Weeds_of_Christmas_Trees.pdf)

Reichenberger, S., M. Bach, A. Skitschak, and H. Georg Frede. 1997. Mitigation strategies to reduce pesticide inputs into ground- and surface water and their effectiveness; A review. *Science of the Total Environment* 384: 1–35.

Photos in figures 1, 7–15, 19, 20, 23–26, 28–30, 36, and 38 by Ed Peachey.

Photos in figures 31, 34, 37, and 39–42 by Chal Landgren.

Photos in figures 2–6, 16–18, 21, and 22 by Tim Miller.

---

Trade-name products and services are mentioned as illustrations only. This does not mean that the participating Extension Services endorse these products and services or that they intend to discriminate against products and services not mentioned.

---

© 2011 Oregon State University

Pacific Northwest Extension publications are produced cooperatively by the three Pacific Northwest land-grant universities: Oregon State University, Washington State University, and the University of Idaho. Similar crops, climate, and topography create a natural geographic unit that crosses state lines. Since 1949, the PNW program has published more than 600 titles, preventing duplication of effort, broadening the availability of faculty specialists, and substantially reducing costs for the participating states.

Published and distributed in furtherance of the Acts of Congress of May 8 and June 30, 1914, by the Oregon State University Extension Service, Washington State University Extension, University of Idaho Extension, and the U.S. Department of Agriculture cooperating.

The three participating Extension services offer educational programs, activities, and materials—without regard to race, color, religion, sex, sexual orientation, national origin, age, marital status, disability, and disabled veteran or Vietnam-era veteran status—as required by Title VI of the Civil Rights Act of 1964, Title IX of the Education Amendments of 1972, and Section 504 of the Rehabilitation Act of 1973. The Oregon State University Extension Service, Washington State University Extension, and University of Idaho Extension are Equal Opportunity Employers.

Published March 2011. Replaces *Managing Weeds and Vegetation in Christmas Trees* (PNW 219).