

development of extraordinary materials is not needed—a result of using thick liquid walls of lithium to protect HIF chamber materials from neutron damage.

The classical deposition of ion beams in fusion targets contrasts sharply with the complex plasma physics of laser-matter interaction. The greater effectiveness per beam megajoule amplifies HIF's crucial ability to deliver to fuel targets 10 times the goal of the National Ignition Facility's (NIF's) laser. More, and more effective, driver energy reduces the needed degree of fuel compression substantially. Reduced compression is accompanied by reduced growth of hydrodynamic instabilities, which in turn relaxes tolerances for pellet fabrication.

The compression requirement is reduced to the realm already demonstrated in ICF research by HIF's ability to use the fast-ignition method. Only a small fraction of the pre-compressed fuel is heated to ignition temperature to start propagating fusion burn. If fast ignition could be accomplished with laser beams, NIF's energy gain could be 10 times higher than its official goal. But fast ignition via laser driver holds exquisite challenges. In contrast, 10 years ago in Russia, Boris Sharkov and associates began designing fusion pellets and HIF driver layouts to exploit HIF's ability to achieve fast ignition with classical energy deposition, in fuel pre-compressed to a density already experienced in the lab, with a relatively long-duration ion pulse, in geometries simple to fabricate.<sup>3</sup>

Why has the US not taken advantage of HIF? The oil “shocks” of 1973 and 1979 were not taken seriously enough, nor were warnings of oil's approaching limits. The issue has been leadership. Elements of HIF are spread among the offices of DOE's three undersecretaries. Tellingly, all fusion work to date has been outside the office responsible for civilian energy systems. The National Academy of Sciences, in reviewing military ICF programs in 1985, noted HIF's advantages but averred that HIF was “supported primarily by other programs.”<sup>4</sup> But there are no other programs. HIF has been an orphan—as Burton Richter put it, “starved and virtually ignored.” While energy production is the sole purpose of HIF, its homelessness is shared by all inertial fusion energy work in the US.<sup>5</sup>

We could be much closer to ICF power than we are, but the situation is good overall. While the US has veered from HIF's founders' use of mainstream

accelerator technology, European and Russian HIF programs steadily resolve details and use new facilities. Decades of ICF progress, using the laser driver and various other technologies, have built a formidable technical basis and community. The fast-ignition concept that benefits HIF enormously only came to light in the 1990s. And HIF enjoys the continuous advance of accelerator technology.

Heavy-ion fusion was discovered at DOE labs. DOE now has requested that the National Academy of Sciences assess the need for an office charged to develop civilian power from ICF. The NAS study is expected to take a year, beginning this summer. An unbiased examination will show, again, that through HIF, fusion power is much closer than it appears. Establishing a home for inertial fusion energy would accomplish a lot.

## References

1. R. O. Bangerter et al., eds., *ERDA Summer Study of Heavy Ions for Inertial Fusion*, rep. no. LBL-5543, Lawrence Berkeley Laboratory, Berkeley, CA (1976).
2. B. Y. Sharkov, *Nucl. Instrum. Methods Phys. Res. A* **577**, 14 (2007).
3. J. Foster, oral report to the US Department of Energy's Energy Research Advisory Board, May 1979. The full report, dated 17 October 1979, is still restricted.
4. Committee for the Review of the Department of Energy's Inertial Confinement Fusion Program, *Review of the Department of Energy's Inertial Confinement Fusion Program*, National Academy Press, Washington, DC (1986).
5. R. Betti, *The Next Generation of Fusion Energy Research*, testimony before the House Committee on Science and Technology, Subcommittee on Energy and Environment, Washington, DC, 29 October 2009.

**Robert J. Burke**  
(rjburke@earthlink.net)  
Fusion Power Corp  
Santa Cruz, California

## Of wetting and osmotic transport

In his article “The First Wetting Layer on a Solid,” Peter Feibelman (PHYSICS TODAY, February 2010, page 34) points out that the first layer of water molecules on a solid surface embodies the boundary condition for water transport, pollution, corrosion, and other molecular transport phenomena. That observation and the revealing high-resolution images presented bring to mind a fundamental problem of osmotic water transport.

In 1827 René Dutochet pointed out that osmosis actually involves binary

transport,<sup>1</sup> in which water moves one way and solute moves the other way. In 1855 Adolf Fick took the idea much further,<sup>2</sup> expanding on the work of other experimentalists. He considered a cylindrical pore in a hydrophilic membrane separating either water or a dilute salt solution on one side and a concentrated one on the other.<sup>3</sup> He reasoned that water will preferentially flow along the walls and salt will tend to migrate along the axis of the pore. As a consequence, he expected concentration gradients in the plane of the pore. Under certain conditions, he suggested, salt migration could be completely inhibited even though the pore might be large enough to allow migration of salt molecules. Subsequent contributions by Jacobus van't Hoff<sup>4</sup> and Walther Nernst<sup>5</sup> established that molecular diffusion in aqueous solutions involves the migration of a solute in one direction driven by the gradient of osmotic pressure, and the flow of water in the opposite direction.

Binary transport in aqueous solutions is widely recognized, but the actual mechanisms are not clear. Solute diffusion involves the random migration of free molecules or ions. However, because water is a condensed phase, its migration cannot be visualized in terms of random motion of molecules. It is not clear if such a flow of water can be considered viscous, because viscosity typically involves wall effects and external forces. If it is viscous, what is the nature of that flow?

Work along the lines described in Feibelman's article may throw more light on the nature of binary transport in osmosis and on molecular diffusion in which random motion of unattached molecules in one direction is accompanied by the migration of a condensed phase in the opposite direction.

## References

1. R. J. H. Dutochet, *Ann. Chim. Phys.* **35**, 393 (1827).
2. A. Fick, *Philos. Mag. J. Sci.* **10**, 30 (1855).
3. T. N. Narasimhan, *Eos Trans. Am. Geophys. Union* **85**(47), 499 (2004).
4. J. H. van't Hoff, *Z. Phys. Chem.* **1**, 481 (1887).
5. W. H. Nernst, *Z. Phys. Chem.* **2**, 613 (1888); **2**, 634 (1888).

**T. N. Narasimhan**  
(tnnarasimhan@lbl.gov)  
University of California, Berkeley

## A two-color twist on test taking

Test taking is a humiliating experience for many students, with no perceived direct educational benefit. That need not be

so. When circumstances allow, I use a two-pen method of collaborative testing.

The method is simple. First, schedule approximately 50% more time for the exam than you think would be required to complete the test, and provide about twice the space you think the students might need to answer each question on the paper. Hand out pens of a particular color and brand that are not commonly used (for example, a particular shade of blue), and have the students commence to work on their exams individually. After the expected time of completion has elapsed, collect the first set of pens, and hand out pens of a different color. (I use red as the second color, then grade the exams in purple. Psychologists suggest that grading in red may carry problematic emotional baggage.)

When the students are armed with their new pens, the exam becomes a free-for-all collaborative event. Students can organize themselves as they prefer while they try to get the right answers to the problems. At last, the smartest student is also the most popular! The written answers reveal which errors they knew they made, and then their attempt to provide the right answer, in their own hand. Students don't get to leave early or hide their work.

Grading is easy: They get full credit for work in the first color and half credit

for correct work in the second. As you can imagine, they find most of their mistakes themselves and correct them, so I generally have much less work trying to figure out what they did wrong and how many points it should be worth. Students have told me that they leave the exam having figured out what they didn't understand, filled in the gaps, and strengthened their relationships with their peers. An easy win-win for a busy professor.

**John Selker**

(john.selker@oregonstate.edu)

Oregon State University  
Corvallis

## Building for NCAR's future

We at the National Center for Atmospheric Research (NCAR) enjoyed the article "Laboratory Architecture: Building for an Uncertain Future" (PHYSICS TODAY, April 2010, page 40). However, we would like to correct a factual inaccuracy about our organization.

Although NCAR is planning construction of a new supercomputing center in Cheyenne, the project is still in the design and approval process; our operations remain in Boulder, Colorado, at this time.

The NCAR-Wyoming Supercomputing Center project is a collaborative partnership that is regionally valuable and offers the scientific community crit-

ical computing resources. NCAR weighed several location options and selected the Wyoming site because it combined the greatest increase in scientific benefits for the university community NCAR serves and the best value for taxpayer investment.

The founding vision of collaboration among scientists will remain a reality at the Mesa Lab, even if the supercomputing operations move. The significant changes in the NCAR computing facility will be in the length of the fiber-optic network, the greater efficiency of the new facility, and the 20-fold increase in computing power.

More information is available at <http://www.cisl.ucar.edu/nwsc>.

**Roger Wakimoto**

National Center for Atmospheric Research  
Boulder, Colorado

## Correction

April 2010, page 18—We should have stated that by 2007, mass-independent fractionation of mercury isotopes had been independently observed by several groups.<sup>1</sup>

### Reference

1. See, for example, T. A. Jackson et al., *Appl. Geochem.* **23**, 547 (2008); L. Laffont et al., *Environ. Sci. Technol.* **43**, 8985 (2009); B. A. Bergquist, J. D. Blum, *Elements* **5**, 353 (2009) and references therein. ■

Search for physics conferences at  
[www.physicstoday.org/cal/eventhome.jsp](http://www.physicstoday.org/cal/eventhome.jsp)

SEARCH EVENTS:  
All types  
All subjects  
All countries  
Jan. - Dec. month  
2008 year

conference  
exhibition  
seminar  
workshop

80 subjects from  
Accelerators to  
Wireless


## X-Ray and Gamma Ray Detectors

- Solid State Design
- No Liquid Nitrogen
- Thermoelectric Cooler
- Easy to Use
- USB Controlled
- Low Cost





Your choice of  
Si-PIN Detector or Super Silicon Drift Detector or CdTe-diode Detector

**Si-PIN Spectrum**




149 eV FWHM  
6 mm<sup>2</sup> x 500 μm  
25.6 μs peaking time  
P/B Ratio: 6200/1

**SUPER SDD Spectrum**



127 eV FWHM  
25 mm<sup>2</sup> x 500 μm  
11.2 μs peaking time  
P/B Ratio: 7000/1

**CdTe Spectrum**



850 eV FWHM  
14.4 keV  
122 keV

**APPLICATIONS**

- Fluorescence
- Portable Instruments
- OEM
- X-Ray Tube Beam Monitor
- Process Control
- Semiconductor Processing
- Lead Detectors
- Nuclear Plant Monitors
- RoHS/WEEE Compliance
- Vacuum Applications
- Environmental Monitors
- Plastic & Metal Separation
- Nuclear Medicine
- Sulfur in Oil & Coal Detection
- Teaching and R&D
- Nuclear Safeguards Verification
- Art & Archaeology
- Space & Astronomy

**OEM's #1 Choice for XRF**



**AMPTEK Inc.** 14 DeAngelo Drive, Bedford, MA 01730-2204 USA  
Tel: +1 781 275-2242 E-mail: [sales@amptek.com](mailto:sales@amptek.com) [www.amptek.com](http://www.amptek.com)