

# Pickle Fact Sheet

Rec'd: - 4/23/92  
Total - 310 - 5  
Shelf - 241  
Last invoice - 23697  
Count date: 4/24/92  
Job # Qc

## Four Basic Pickle Products

Brined or fermented pickles require three weeks or longer to cure. Cucumbers change from a bright green to an olive or yellow green. The interior becomes uniformly translucent.

Salt is added at the beginning of the process to interfere with the growth of microorganisms other than the lactic acid bacteria. Vinegar is also used to create an acid environment which discourages growth of microorganisms other than the lactic acid bacteria. The salt in the original brine is quite concentrated and draws liquid out of the cucumbers. The liquid drawn from the cucumbers contains carbohydrates which serve as food for this bacteria.

As the lactic acid bacteria multiply, they produce lactic acid which makes brine even more acidic. This acid reacts with the chlorophyll of the cucumbers causing it to become an olive-green.

A scum usually forms on the surface of the brine as yeast and mold grow. It is important to remove the scum daily.

Fresh pack or quick process pickles are made by combining ingredients (with or without heating) and water bathing right away. As a rule of thumb, recipes should call for at least as much vinegar (5% acid) as water. Use only reliable up-to-date recipes and follow them carefully. Do not alter vinegar, food, or water proportions.

Fruit pickles are made of whole fruits simmered in a spicy, sweet-sour syrup.

Relishes are made from chopped fruits or vegetables with seasonings.

## Utensils

Do not use zinc, copper, brass, galvanized metal, or iron utensils. These metals may react with acids or salts and cause undesirable color changes in the pickles or form undesirable compounds.

## Ingredients Make a Difference

### Vegetables or Fruits

- Select tender vegetables and firm fruits that are free from blemishes.
- For best results, use cucumbers within 24 hours of picking.
- Use a variety of cucumbers grown especially for pickling. Good quality pickles are not made from immature "slicing" cucumbers.
- Use light skinned cucumbers that are attractively warty, not more than 2 inches in diameter and young enough so the seeds have not matured.



OREGON STATE UNIVERSITY EXTENSION SERVICE

Revised July 1992

4-H 93322L

- Commercially waxed cucumbers should not be used for fermented pickles or whole pickles. Pickling liquid cannot penetrate into the waxed cucumbers.
- Wash all vegetables and fruits thoroughly in cold water. Do not use if moldy.
- Remove blossom ends of cucumbers. They may contain enzymes which cause softening in pickles.

#### Salt/Recommended

- Pickling and canning salt are pure, granulated salts containing no anti-caking agents, and no iodine. They are hard to find. In moist climates, problems with caking of the salt leads to consumer dissatisfaction, thus reducing its availability for purchase.
- Table salt can be used. It can cause a cloudy brine, but this does not harm the pickles.

#### Salt/Not recommended

- Sea salt is essentially iodized table salt with a different name for sales purposes. Salt labeled "sea salt" is not suitable for pickling because it contains mineral impurities which may also cause pickle discoloration.
- Ice cream salt or rock salt might contain unsafe impurities so it is not recommended.
- Sour salt is a small tablet containing citric acid and salt that is used in the commercial canning of sea food. At retail level, it is marketed as a water softener. The salt in the tablet removes ions of calcium and magnesium, while the citric acid renders the iron in the water unavailable. This type of salt, due to its tablet form and mixture, is not suitable for pickling purposes.
- Salt substitutes (such as potassium chloride) cannot be used to make satisfactory pickle products. The pickles will be crisp and crunchy, but will not have the characteristic salty pickled taste. Potassium has a bitter taste and the pickles generally taste terrible.
- Flaked salt varies in density, so it is not recommended for use in pickled and fermented foods.

#### Vinegar

- Use vinegar with 5 percent acidity.
- White distilled vinegar has a sharp pungent acid taste and is desirable when light color is important such as for pickled onions, cauliflowers and pears.
- Cider vinegar has a milder flavor but may discolor some pickle products.
- Do not use homemade vinegar or brands with an unknown acidity. The percent acidity must be known for successful pickling.
- If a less sour product is desired, add sugar to the pickling solution rather than reducing the amount of vinegar. Vinegar is important for safety reasons.

## Spices

- Use whole spices. Powdered spices may cause the product to darken and become cloudy.
- Use fresh spices. Last year's supply will have lost much flavor.
- Pick dill as soon as the flowers begin to open.
- Dill seed can be substituted for fresh dill, but may not have as good flavor. (3 dill heads = 1 tablespoon seeds = 1 to 3 drops dill oil)

## Garlic

- Use fresh or mature garlic. Immature or old garlic turns pink or purple as anthocyanins (coloring pigment) react to acid.

## Water

- Moderately soft water makes the best brine for pickles. Hard water may cause cloudiness and iron and sulfur may cause off-colors.
- If only hard water is available, boil and let set for 24 hours and skim off scum. Water should be used from the top of the container without disturbing any sediment on the bottom.

## Sugar

- Either white or brown sugar may be used.
- Brown sugar may give a darker color to the brine.
- Corn syrup and honey may be too strong flavored.

## Firming Agents

- Alum may be safely used to firm fermented cucumbers. However, it is not necessary and is not included in most up-to-date recipes.
- Lime, called slake lime, lime hydrate, or hydrated lime, is really calcium hydroxide. If used, it should be food grade. Soaking fresh cucumber slices in a lime-water solution for 12-24 hours before pickling improves firmness. Excess lime absorbed by the cucumbers must be removed to make safe pickles. Drain, rinse, and then resoak cucumbers in fresh water for one hour. Repeat the rinsing and soaking steps two more times.

## Processing pickles

Both microorganisms and enzymes can cause spoilage of pickled products. Processing will prevent these problems. Follow recommended methods for traditional water bath processing or low temperature pasteurization.

DO NOT PROCESS FERMENTED PICKLES BEFORE THEY DEVELOP AN ACIDIC TASTE.

## Storing Pickles

Cool      Extreme fluctuation of temperature can cause a breakdown of the texture. Pickles stored at warm temperatures could start fermenting again and cause the jars to break.

**Dark** Light often causes products to fade and become less appetizing. This does not necessarily produce spoilage.

**Dry** Continuous moisture may cause closures to rust and eventually produce spoilage.

Remove jar rings. If liquid is lost during processing and rings are not removed, it will cause corrosion and the rings will be hard to remove. This is especially true for pickles due to their high acid and salt content.

After opening, store pickles in the refrigerator with the pickling liquid covering the fruits or vegetables.



Adapted from a University of Idaho Extension System publication. Revised by Nellie Oehler, Extension agent, Lane County, with Carolyn Raab, Extension foods and nutrition specialist, Oregon State University.

---

Extension Service, Oregon State University, Corvallis, O. E. Smith, director. Produced and distributed in furtherance of the Acts of Congress of May 8 and June 30, 1914. Extension work is a cooperative program of Oregon State University, the U.S. Department of Agriculture, and Oregon counties. Oregon State University Extension Service offers educational programs, activities, and materials without regard to race, color, national origin, sex, or disability as required by Title VI of the Civil Rights Act of 1964, Title IX of the Education Amendments of 1972, and Section 504 of the Rehabilitation Act of 1973. Oregon State University Extension Service is an Equal Opportunity Employer.

---