


File Copy



**FOOD OF FIVE SPECIES OF
COOCCURRING FLATFISHES ON
OREGON'S CONTINENTAL SHELF**

by Michael J. Kravitz
William G. Pearcy

Reprinted from **Fish Bull.** 74(4):984-990.



**OREGON STATE
SEA GRANT
COLLEGE PROGRAM**

Reprint no. ORESU-R-76-027

FOOD OF FIVE SPECIES OF
COOCCURRING FLATFISHES ON
OREGON'S CONTINENTAL SHELF

The purpose of this paper is to describe and to compare the food of five flatfish species that actually cooccurred at one specific time and place on the central Oregon continental shelf: English sole, *Parophrys vetulus* Girard; rex sole, *Glyptocephalus zachirus* Lockington; rock sole, *Lepidopsetta bilineata* (Ayres); petrale sole, *Eopsetta jordani* (Lockington); and Pacific sanddab, *Citharichthys sordidus* (Girard). These demersal fishes are common along the west coast of North America, their ranges overlapping between southern California and the Gulf of Alaska (Hart 1973). *Parophrys vetulus*, *C. sordidus*, and *L. bilineata* occur mainly on the inner continental shelf. *Eopsetta jordani* is fished commercially on its feeding grounds (73-128 m), and in deep water (311-457 m) where spawning occurs (Forrester 1969). *Glyptocephalus zachirus* has a broad bathymetric range—it is common off Oregon and Washington from 90 to 550 m (Alverson et al. 1964). Off Oregon it was the second most numerous member of a species association ranging from 119 to 199 m, on an average sediment type of 69% sand, 19% silt, and 12% clay (Day and Percy 1968). In that same study, *C. sordidus* and *P. vetulus* composed 80.3% of a species association of fishes in shallower water (42-73 m) on a sandy bottom. According to Alverson (1960), *L. bilineata* is common on sandy or gravel bottom. The five flatfish species attain maximum sizes ranging from 410 mm for *C. sordidus* to 700 mm for *E. jordani* (Hart 1973).

Percy and Vanderploeg (1973) listed major food items—combined from several locations, seasons, and years—for most of the above species. That

study provided generalized information on food habits, but little insight into possible intra- or interspecific differences in diets resulting from actual interaction among cooccurring fishes. Our study is based on a single collection minimizing temporal and spatial variations associated with sampling. Food items were identified to species whenever possible. Thus, a detailed comparison of food taxa is arrived at with minimal geographic and no seasonal effects.

A trawl haul of 75 min total duration was made beginning at 1345 h Pacific daylight time, on 13 April 1975 with an Atlantic-Western trawl (24-m footrope) from the *Betty-A*, a commercial dragger, at approximately lat. 44°42'N, long. 124°24'W. Depth of water was 95-106 m. The sediment was sand (Byrne and Panshin 1968). Stomachs of fishes were removed and preserved in Formalin¹ at sea (Table 1). Food items were identified and enumerated in laboratories ashore.

TABLE 1.—Fishes captured in an Atlantic-Western trawl on 13 April 1975.

| Species | No. caught | Total length range (mm) | No. examined | No. with stomach contents |
|----------------------------------|------------|-------------------------|--------------|---------------------------|
| <i>Citharichthys sordidus</i> | 181 | 90- 377 | 62 | 26 |
| <i>Parophrys vetulus</i> | 50 | 230- 450 | 50 | 37 |
| <i>Glyptocephalus zachirus</i> | 24 | 240- 360 | 22 | 21 |
| <i>Eopsetta jordani</i> | 22 | 240- 510 | 12 | 7 |
| <i>Lepidopsetta bilineata</i> | 19 | 247- 474 | 19 | 15 |
| <i>Raja binoculata</i> | 9 | 940-1,460 | 8 | 6 |
| <i>Raja rhina</i> | 2 | 800- 890 | 2 | 1 |
| <i>Raja kincaidii</i> | 2 | 560- 570 | 2 | 0 |
| <i>Pleuronichthys verticalis</i> | 2 | 251- 254 | 2 | 1 |
| <i>Ophiodon elongatus</i> | 1 | 850 | 1 | 0 |
| <i>Squalus acanthias</i> | 1 | 1,000 | 0 | — |

Forage Organisms

All the food items identified from five species of flatfishes are listed in Table 2, and the major food taxa (taxa having a frequency of occurrence of 10% or more) are listed for individual fish of three species of flounders in Table 3.

Parophrys vetulus had a diverse diet, feeding primarily on polychaetes and amphipods. Mollusks, ophiuroids, and crustacea were also represented. The amphipod *Ampelisca macrocephala*, the most numerous single prey species, occurred in 60% of fish. The diversity of the diet of *P. vetulus* is

due to the many different types of food consumed by individual fish (represented by the vertical columns in Table 3) rather than by different fish feeding on different prey. *Parophrys vetulus* appears to be an opportunistic feeder. Forrester (1969) reported polychaetes, clams, and ophiuroids as primary food organisms of *P. vetulus*, with incidental occurrences of sandlance, crab, amphipods, shrimp, squid, and small fish. Percy and Vanderploeg (1973) found polychaetes, amphipods, and pelecypods were important prey of *P. vetulus* off Oregon.

Glyptocephalus zachirus fed primarily on four species of amphipods and secondarily on polychaetes. Amphipods occurred in all but one stomach, polychaetes in 71% of the stomachs with food. Nematodes were encountered in 38% of the stomachs but were probably parasitic (Robert Olson, pers. commun.). Percy and Vanderploeg (1973) also found polychaetes and amphipods to be the major food of *G. zachirus* off Oregon.

The principal food of *Lepidopsetta bilineata* was ophiuroids. All but one individual had been feeding on *Ophiura*, which constituted the bulk of the stomach contents. A few polychaetes and mollusks were also present. According to Shubnikov and Lisovenko (1964), the basic items of its diet are polychaetes, mollusks, shrimps, and other crustaceans. Fishes (sandlance) and echinoderms were occasionally found in stomachs. Food items reported for *L. bilineata* in Hecate Strait, British Columbia, by Forrester and Thomson (1969) were clams, polychaetes, crabs, shrimps, sandlance, herring, echinoderms, and amphipods.

Eopsetta jordani preyed on fishes and decapod crustaceans. Polychaetes and amphipods were not present in its diet. Ketchen and Forrester (1966) found euphausiids, herring, sandlance, and shrimp as major food items in stomachs of *E. jordani*. Percy and Vanderploeg (1973) reported shrimps, pelagic fishes, and euphausiids as major food items, indicating that this species feeds largely on pelagic prey.

Citharichthys sordidus had been feeding intensively on the northern anchovy, *Engraulis mordax*. Anchovy were noted in nearly all the sanddab when stomachs were removed, and all intact preserved stomachs contained them. Since anchovy were not caught in the otter trawl, feeding in the net is thought to be unlikely. According to Percy and Vanderploeg (1973), euphausiids, shrimps, amphipods, and crab larvae were common in *C. sordidus* stomachs.

¹Reference to trade names does not imply endorsement by the National Marine Fisheries Service, NOAA.

TABLE 2.—Taxa identified from stomach contents of five species of Pacific Northwest flatfishes.¹

| Taxa | <i>Parophrys vetulus</i> | <i>Glyptocephalus zachirus</i> | <i>Lepidopsetta bilineata</i> | <i>Eopsetta jordani</i> | <i>Citharichthys sordidus</i> |
|-----------------------------------|------------------------------|------------------------------------|-----------------------------------|-----------------------------|-----------------------------------|
| POLYCHAETA | 94.6 | 71.4 | 40.0 | | |
| Aphroditidae | | | | | |
| <i>Aphrodita negligens</i> | | | x | | |
| Capitellidae | | | | | |
| [Capitellidae spp.] | 10.8 | | | | |
| <i>Notomastus</i> spp. | x | | | | |
| Chaetopteridae spp. | x | | | | |
| Cirratulidae | | | | | |
| [Cirratulidae spp.] | 40.5 | | | | |
| <i>Chaetozone setosa</i> | x | | | | |
| <i>Chaetozone</i> spp. | x | | | | |
| <i>Tharyx</i> spp. | x | | | | |
| Goniadidae | | | | | |
| Goniadidae spp. | x | x | | | |
| <i>Glycinde picta</i> (?) | 13.5 | | | | |
| Lumbrineriidae | | | | | |
| <i>Lumbrineris</i> spp. | 18.9 | | | | |
| Magelonidae | | | | | |
| <i>Magelona</i> spp. | x | | | | |
| Maldanidae spp. | x | | | | |
| Nephtyidae | | | | | |
| [<i>Nephtys caecoides</i>] | x | | | | |
| <i>Nephtys</i> spp. | 43.2 | | | | |
| Onuphidae | | | | | |
| <i>Nothria geophiliformis</i> (?) | x | | | | |
| <i>Nothria iridescens</i> (?) | 10.8 | 14.3 | 13.3 | | |
| <i>Nothria</i> spp. | | 19.0 | | | |
| Opheliidae | | | | | |
| Opheliidae spp. | | 9.5 | | | |
| <i>Ammotrypane aulogaster</i> | 8.1 | | | | |
| Orbiniidae | | | | | |
| <i>Haploscoloplos</i> spp. | x | | | | |
| Oweniidae | | | | | |
| <i>Myriochele oculata</i> | 16.2 | | | | |
| <i>Myriochele</i> spp. | 10.8 | | | | |
| <i>Owenia</i> spp. | 16.2 | | | | |
| Paraonidae spp. | 21.6 | | | | |
| Pectinariidae | | | | | |
| <i>Pectinaria</i> spp. | x | x | x | | |
| Phyllodoceidae | | | | | |
| <i>Eteone longa</i> | x | | | | |
| Polynoidae spp. | | | 13.3 | | |
| Sigalionidae | | | | | |
| [<i>Sigalionidae</i> spp.] | 18.9 | | | | |
| <i>Thalenessa spinosa</i> | x | | | | |
| Spionidae spp. | 18.9 | | | | |
| Terebellidae | | | | | |
| [Terebellidae spp.] | 35.1 | | | | |
| <i>Polycirrus</i> spp. | x | | | | |
| Unidentified | x | x | | | |
| GASTROPODA | 13.5 | | | | 11.1 |
| <i>Cylichna attonsa</i> | 10.8 | | | | |
| <i>Mitrella gouldii</i> | x | | | | |
| <i>Mitrella</i> spp. (?) | | | | | 11.1 |
| PELECYPODA | 27.0 | 9.5 | 13.3 | | |
| <i>Acila castrensis</i> | | x | | | |
| <i>Axinopsida serricata</i> | x | | | | |
| <i>Cardiomya oldroydi</i> | x | | | | |
| <i>Nucula tenuis</i> | 16.2 | | | | |
| <i>Macoma</i> spp. | x | | | | |
| <i>Tellina carpenteri</i> (?) | 8.1 | | 13.3 | | |
| Unidentified | x | | | | |
| SCAPHOPODA | 16.2 | 4.8 | | | |
| <i>Dentalium</i> sp. | 13.5 | | | | |
| Unidentified fragment | x | | | | |
| Scaphopoda(?) | | x | | | |
| CEPHALOPODA | | | | | 11.1 |
| Octopoda | | | | | 11.1 |
| Beak of <i>Loligo</i> spp. (?) | | | | | 11.1 |
| CRUSTACEA | 91.9 | 100.0 | 6.7 | 28.6 | 33.3 |
| Cypris larvae (?) | x | | | | |
| Copepoda (calanoid) | 10.8 | | | | |
| Mysidacea | | | | | |
| <i>Neomysis</i> spp. | | | | | 11.1 |
| Unidentified | | | | 14.3 | |
| Ostracoda (?) | x | | | | |
| Cumacea | 10.8 | | | | |
| Tanaidacea | x | | | | |

TABLE 2.—Continued.

| Taxa | <i>Parophrys vetulus</i> | <i>Glyptocephalus zachirus</i> | <i>Lepidopsetta bilineata</i> | <i>Eopsetta jordani</i> | <i>Citharichthys sordidus</i> |
|---------------------------------|------------------------------|------------------------------------|-----------------------------------|-----------------------------|-----------------------------------|
| Euphausiacea | | | | | |
| <i>Euphausia pacifica</i> | | | | | 22.2 |
| Decapoda | | | | | |
| Natantia | | | | | |
| <i>Crangon</i> spp. | | | | | 11.1 |
| <i>Nectocrangon</i> spp. | | | | 14.3 | |
| Unidentified shrimp | | | | | x |
| Reptantia | | | | | |
| <i>Pagurus samuelis</i> | | | x | | |
| <i>Mursia</i> spp. | | | | 14.3 | |
| Crab leg | | x | | | |
| Amphipoda | 83.8 | 95.2 | | | |
| Ampeliscidae | | | | | |
| <i>Ampelisca cristata</i> | | x | | | |
| <i>Ampelisca macrocephala</i> | 59.5 | 33.3 | | | |
| <i>Ampelisca</i> spp. | x | | | | |
| Amphilochidae spp. (?) | x | | | | |
| Aoridae | | | | | |
| <i>Lembos</i> spp. | x | | | | |
| Argissidae | | | | | |
| <i>Argissa hamatipes</i> | x | | | | |
| Isaeidae | | | | | |
| <i>Photis brevipes</i> | x | x | | | |
| <i>Protomedea</i> spp. | x | | | | |
| Lysianassidae | | | | | |
| Lysianassidae spp. | x | | | | |
| <i>Acidostoma</i> spp. | | x | | | |
| <i>Anonyx anivae</i> | x | | | | |
| <i>Hipomedon wecomus</i> | 18.9 | 28.6 | | | |
| Oedicerotidae | | | | | |
| <i>Monoculodes emarginatus</i> | x | | | | |
| <i>Monoculodes</i> sp. #1 | x | x | | | |
| <i>Synchelidium shoemakeri</i> | x | x | | | |
| <i>Westwoodilla caecula</i> | x | | | | |
| Phoxocephalidae | | | | | |
| <i>Paraphoxus abronius</i> | x | | | | |
| <i>Paraphoxus daboius</i> (?) | x | | | | |
| <i>Paraphoxus epistomus</i> (?) | 21.6 | 33.3 | | | |
| <i>Paraphoxus fatigans</i> | 10.8 | 4.8 | | | |
| <i>Paraphoxus lucubrans</i> | x | | | | |
| <i>Paraphoxus milleri</i> | | x | | | |
| <i>Paraphoxus obtusidens</i> | 16.2 | 33.3 | | | |
| <i>Paraphoxus variatus</i> | x | | | | |
| <i>Paraphoxus</i> spp. | 10.8 | | | | |
| Pleustidae | | | | | |
| Pleustidae spp. | | x | | | |
| <i>Pleusymtes coquilla</i> | | x | | | |
| OPHIUROIDEA | 83.8 | | 93.3 | | |
| <i>Amphiodia periercta</i> | 10.8 | | | | |
| <i>Amphiodia urtica</i> | 10.8 | | | | |
| <i>Amphiuridae</i> spp. | 10.8 | | | | |
| <i>Ophiura lutkeni</i> | 35.1 | | 73.3 | | |
| <i>Ophiura sarsii</i> | x | | | | |
| <i>Ophiura</i> spp. | 8.1 | | 20.0 | | |
| PISCES | | 9.5 | | 100.0 | 100.0 |
| Agonidae | | | | 14.3 | |
| <i>Engraulis mordax</i> | | | | 14.3 | 100.0 |
| <i>Glyptocephalus zachirus</i> | | | | 14.3 | |
| <i>Radulinus</i> spp. | | | | 14.3 | |
| Unidentified | | x | | x | |
| Fish scale | x | x | | | |
| NEMERTINEA | 13.5 | | | | |
| NEMATODA | x | x | | | |
| SIPUNCULIDA | 2.7 | | 13.3 | | |
| ECHIURIDA | | x | | | |
| ACANTHOCEPHALA | | x | | | |
| Miscellaneous | | | | | |
| Gastropod egg case | x | | | | |
| Egg mass | x | | | | |
| Lenses | | | | | x |
| Unidentified remains | x | x | | | |

¹Frequency of occurrence is given as a percentage for food taxa whenever these taxa occur in 10% or greater of any of the five species of fishes. Opheliidae were significant on a weight basis in *G. zachirus*, and were combined with *Ammotrypane aulogaster* for calculation of similarity. An "x" denotes any other occurrence.

²Taxa enclosed within brackets were treated as a single group in Table 3.

TABLE 3.—Numbers of food items of major¹ taxa in contents of individual fish stomachs. Each vertical column represents one stomach.

| Fish species and taxa | Number of food items | | | | | | | | | | | | | | | Total no. | | | | | |
|-------------------------------------|----------------------|---|---|---|---|---|---|---|---|---|---|---|----|----|---|-----------|---|---|---|---|----|
| <i>Parophrys vetulus</i> | | | | | | | | | | | | | | | | | | | | | |
| POLYCHAETA: | | | | | | | | | | | | | | | | | | | | | |
| Capitellidae | | | | | | | | | | | | | | | | 3 | 1 | 1 | 1 | 6 | |
| Cirratulidae | 1 | | 1 | 1 | 3 | 1 | 1 | | 1 | 1 | 1 | 1 | 1 | 1 | 3 | | | | 1 | 1 | 29 |
| <i>Glycinde picta</i> (?) | | | 1 | | 1 | | | 1 | | | 1 | | | | | | | | | | 5 |
| <i>Lumbrineris</i> spp. | | | | | * | * | | | | 2 | 3 | | 1 | * | 1 | | | | | | 10 |
| <i>Nephtys</i> spp. | 1 | | 4 | | 7 | 3 | | 4 | | * | * | 3 | | 10 | 1 | 3 | 1 | 3 | | 1 | 47 |
| <i>Nothria iridescens</i> (?) | | | | 1 | | 1 | 2 | | | | | | | 2 | | | | | | | 6 |
| 2 [<i>Myriochele oculata</i> | | | | | | | | | | | 3 | 1 | 1 | | | | | | 1 | 1 | 8 |
| <i>Myriochele</i> spp. | | | | | | * | | | * | * | * | | | | | | | | | | 4 |
| <i>Owenia</i> spp. | | | | 1 | | | | | * | 1 | | | | 2 | 1 | | * | | | | 7 |
| Paraonidae | | | | | 2 | 1 | | 1 | 1 | * | | | | 5 | | | | | 1 | 1 | 13 |
| Sigalionidae | | | | 1 | | | | 1 | 2 | 1 | | | | 1 | | | | | | 1 | 8 |
| Sponidae | | | | | 1 | 1 | | | 2 | | | | 2 | 1 | 1 | | | | 1 | | 9 |
| Terebellidae | | * | 2 | 1 | | 1 | 1 | 1 | | | 1 | 5 | 2 | 4 | 1 | | * | | | 2 | 23 |
| AMPHIPODA: | | | | | | | | | | | | | | | | | | | | | |
| <i>Ampelisca macrocephala</i> | | 2 | 1 | 1 | 8 | 3 | 2 | | 2 | 5 | 2 | 1 | 12 | 7 | 1 | 2 | 1 | 4 | 4 | 1 | 64 |
| <i>Hippomedon wecomus</i> | | | | | | | | 1 | | | 1 | 1 | 1 | 2 | 1 | 1 | | | | | 8 |
| <i>Paraphoxus epistomus</i> (?) | | | | | | | | | | 4 | 2 | 1 | 1 | 1 | 2 | | | 1 | | 1 | 13 |
| <i>Paraphoxus fatigans</i> | 1 | | | | | | | | | | | | 1 | | 1 | | | | | 1 | 4 |
| <i>Paraphoxus obtusidens</i> | | | | | 2 | 1 | | 1 | | | 1 | 1 | 1 | | | | | | | | 7 |
| <i>Paraphoxus</i> spp. ³ | | | | | | | | * | | 1 | | | | | | | * | 1 | | | 4 |
| GASTROPODA: | | | | | | | | | | | | | | | | | | | | | |
| <i>Cylichna attonsa</i> | | | | 1 | | | 2 | | | | | 1 | | 1 | | | | | | | 5 |
| PELECYPODA: | | | | | | | | | | | | | | | | | | | | | |
| <i>Nucula tenuis</i> | | | | 1 | | | | | | | | 2 | | 4 | 1 | | | | | 2 | 12 |
| SCAPHOPODA: | | | | | | | | | | | | | | | | | | | | | |
| <i>Dentalium</i> sp. | | | | | 3 | 1 | | | 2 | | | 1 | | | | | | | 1 | | 8 |
| COPEPODA (calanoid) | | | | | | 1 | 1 | 1 | | 2 | | | | | | | | | | | 5 |
| CUMACEA | | | | | 2 | | | | | | 1 | | | 1 | | | | | | | 5 |
| OPHIUROIDEA: | | | | | | | | | | | | | | | | | | | | | |
| 2 [<i>Amphiodia periercta</i> | | | | | 1 | | | | | | 1 | | | | 1 | 1 | | | | | 4 |
| <i>Amphiodia urtica</i> | | | | | | | | | 1 | | 1 | | | | 1 | 1 | | | | | 4 |
| Amphiuridae | | | | | | 1 | | | 1 | | 1 | | | | | | | | | 1 | 4 |
| <i>Ophiura lutkeni</i> | | 2 | | 2 | 1 | | | 3 | 1 | 1 | | 2 | 1 | | 1 | 1 | 1 | 1 | | 1 | 18 |
| NEMERTINEA | | | 1 | | | | | | 1 | | * | | | * | * | | | | | | 5 |
| <i>Glyptocephalus zachirus</i> | | | | | | | | | | | | | | | | | | | | | |
| POLYCHAETA: | | | | | | | | | | | | | | | | | | | | | |
| <i>Nothria iridescens</i> (?) | | | | | | 1 | | | | | 1 | | | 2 | | | | | | | 4 |
| <i>Nothria</i> spp. ³ | | * | | | | | | * | | | 3 | | | | 2 | | | | 1 | | 6 |
| Opheliidae | | | | | | | | 1 | | | | 1 | | | | | | | | | 2 |
| AMPHIPODA: | | | | | | | | | | | | | | | | | | | | | |
| <i>Ampelisca macrocephala</i> | | | 1 | | | | | | | | | 2 | 1 | 1 | 1 | | 2 | 2 | | 1 | 10 |
| <i>Hippomedon wecomus</i> | | | | | 1 | | | | | 3 | 1 | | | 1 | | | | | | | 8 |
| <i>Paraphoxus epistomus</i> (?) | | 1 | | 1 | | | | | | 2 | | | | 2 | 1 | 1 | | | | 1 | 9 |
| <i>Paraphoxus obtusidens</i> | | | | | | | 1 | 2 | 3 | | 2 | | | 1 | 1 | 3 | | | | | 13 |
| <i>Lepidopsetta bilineata</i> | | | | | | | | | | | | | | | | | | | | | |
| POLYCHAETA: | | | | | | | | | | | | | | | | | | | | | |
| Polynoidae | | | | | | | | | | | | * | | | | | | | | | 2 |
| <i>Nothria iridescens</i> (?) | | | | | | | | | | | | | | | 1 | 3 | | | | | 4 |
| SIPUNCULIDA (?) | | | | | | | | | | | 1 | | | | | | | | | 1 | 2 |
| PELECYPODA: | | | | | | | | | | | | | | | | | | | | | |
| <i>Tellina carpenteri</i> (?) | | | | | | | | | | | 1 | 1 | | | | | | | | | 2 |
| OPHIUROIDEA: | | | | | | | | | | | | | | | | | | | | | |
| <i>Ophiura lutkeni</i> | | 1 | | | | 1 | 2 | 2 | 4 | 5 | 4 | 1 | 1 | 2 | 3 | | | | * | | 26 |
| <i>Ophiura</i> spp. ³ | | | | | | * | * | | | | | | | | | | | | | | 3 |

¹Taxa having a frequency of occurrence of at least 10%.

²Taxa within brackets were treated as a single group for calculation of similarity.

³Taxa not used to determine similarity.

*Fragment.

Discussion

The flatfishes examined in this study comprised two distinct feeding types based on the species composition of prey and the frequency of occurrence of major food items. *Parophrys vetulus*, *G. zachirus*, and *L. bilineata* were benthophagous, feeding on benthic infaunal and epifaunal invertebrates, mainly polychaetes, amphipods, and

ophiuroids. *Eopsetta jordani* and *C. sordidus* were piscivorous and fed more on pelagic animals, consuming mainly fishes in addition to shrimp, mysids, euphausiids, and cephalopods. Fishes did not occur in the stomachs of the benthic invertebrate feeders, except for two fishes found in *G. zachirus*.

Differences were sometimes obvious in the food habits of fishes within each feeding type. The

similarity among the food habits of the three fishes that preyed on benthic invertebrates was calculated using commonly occurring prey (Table 3) and Horn's (1966) measure of niche overlap. The overlap was largest between *P. vetulus* and *G. zachirus* ($C_\lambda = 0.40$). This is because both fishes fed on the same species of amphipods and the polychaete *Nothria iridescens*. *Parophrys vetulus* preyed on a very diverse array of invertebrate taxa, while *G. zachirus* appeared to be more selective in its feeding.

The amount of food overlap among the other species pairs was low (0.19 between *P. vetulus* and *L. bilineata* and only 0.03 between *G. zachirus* and *L. bilineata*). These low values are explained by the high occurrence of *Ophiura lutkeni* only in *L. bilineata*. Also, *L. bilineata* fed on members of two scaleworm families, Aphroditidae and Polynoidae, neither of which is represented in the other flatfishes.

The food habits of the flatfishes that we found to be mainly piscivorous were also different. *Eopsetta jordani* preyed on various fishes, including benthic agonids, pleuronectids, and cottids, as well as a benthic shrimp and crab, whereas *C. sordidus* fed almost exclusively on the pelagic *Engraulis mordax*.

Partitioning of the food resources among the five flatfish species is obvious from our data—the different, syntopic species fed upon different organisms. According to MacArthur and Pianka (1966), a more productive environment should lead to a more restricted diet in terms of different species eaten, but in a patchy environment this does not apply to predators that spend most of their time searching. If the bottom occupied by *P. vetulus* is inhabited by patches of invertebrates, then this species might be such a scavenging "generalist" predator. Rae (1969) documented a food interaction similar to the one in this study between the lemon sole, *Microstomus kitt*, and witch, *Glyptocephalus cynoglossus*, off Scotland. The witch, restricted to muddy bottoms, fed on a more restricted fauna than the lemon sole, whose diet included the hard-bottom species typical of its habitat in addition to species from muddy-bottom types.

Differences in time of feeding could also account for differences in the species composition of prey. Diel changes in the habits of prey can serve to increase or decrease their exposure to predators, and hence their availability as food (Hobson 1965;

Jones et al. 1973). More so than the other species, the stomach contents of *G. zachirus* were in a late stage of digestion, suggesting that they had fed a longer time before capture than other species.

The diet of fishes is related not only to their feeding behavior but also to their digestive morphology and mouth structure. The size of the mouth relative to body length correlated with the size of food organisms for bothid flounders in Georgia coastal waters (Stickney et al. 1974). Symmetry of the jaws plays an important role in the mode of feeding, as species with symmetrical jaws generally take free-swimming food, while those with asymmetrical jaws are mainly bottom feeders (Yazdani 1969). Flatfishes that feed on polychaetes and mollusks typically have smaller stomachs, larger intestines, and smaller gill rakers with fewer teeth than flatfishes that feed on other fishes (DeGroot 1971; Tyler 1973). The mouths of *P. vetulus*, *G. zachirus*, and *L. bilineata* are small,² the jaws and dentition are better developed on the blind side (i.e., asymmetrical), the teeth are incisorlike (bluntly conical in *L. bilineata*), and the gill rakers are without teeth. These morphological adaptations correlate with the preponderance of benthic invertebrates in their diets. The piscivores, *E. jordani* and *C. sordidus*, on the other hand, have larger mouths,³ nearly symmetrical jaws with sharp teeth, and long gill rakers with teeth.

Acknowledgments

We thank William Colgate, Allan Fukuyama, and Valerie Hironaka for identifying amphipods, mollusks, and ophiuroids, respectively. Discussions with Michael Richardson and Robert Carney were useful. This work is a result of research sponsored [in part] by the Oregon State University Sea Grant College Program, supported by NOAA Office of Sea Grant, U.S. Department of Commerce, under Grant #04-5-158-2.

Literature Cited

ALVERSON, D. L.
1960. A study of annual and seasonal bathymetric catch

²Length of maxillary into head on ocular side is $4\frac{1}{4}$ - $4\frac{2}{5}$, $4\frac{1}{2}$ - $5\frac{1}{4}$, and $3\frac{1}{2}$ - $4\frac{1}{5}$, respectively (Norman 1934); also see Norman for line drawings depicting the relative mouth size of flatfishes discussed in this paper.

³Length of maxillary into head is about $2\frac{2}{5}$ and $2\frac{3}{4}$ (nearly 3), respectively (Norman 1934).

- patterns for commercially important groundfishes of the Pacific Northwest coast of North America. *Pac. Mar. Fish. Comm. Bull.* 4:1-66.
- ALVERSON, D. L., A. T. PRUTER, AND L. L. RONHOLT.
1964. A study of demersal fishes and fisheries of the north-eastern Pacific Ocean. H. R. MacMillan Lectures in Fisheries, Inst. Fish., Univ. B.C., 190 p.
- BYRNE, J. V., AND D. A. PANSHIN.
1968. Continental shelf sediments off Oregon. *Oreg. State Univ. Sea Grant Ext. Mar. Advis. Program* 8, 4 p.
- DAY, D. S., AND W. G. PEARCY.
1968. Species associations of benthic fishes on the continental shelf and slope off Oregon. *J. Fish. Res. Board Can.* 25:2665-2675.
- DEGROOT, S. J.
1971. On the interrelationships between morphology of the alimentary tract, food and feeding behaviour in flatfishes (Pisces: Pleuronectiformes). *Neth. J. Sea Res.* 5:121-196.
- FORRESTER, C. R.
1969. Life history information on some groundfish species. *Fish. Res. Board Can., Tech. Rep.* 105, [17 p.]
- FORRESTER, C. R., AND J. A. THOMSON.
1969. Population studies on the rock sole (*Lepidopsetta bilineata*) of northern Hecate Strait, British Columbia. *Fish. Res. Board Can., Tech. Rep.* 108, 104 p.
- HART, J. L.
1973. Pacific fishes of Canada. *Fish. Res. Board Can., Bull.* 180, 740 p.
- HOBSON, E. S.
1965. Diurnal-nocturnal activity of some inshore fishes in the Gulf of California. *Copeia* 1965:291-302.
- HORN, H. S.
1966. Measurement of "overlap" in comparative ecological studies. *Am. Nat.* 100:419-424.
- JONES, D. A., N. PEACOCK, AND O. F. M. PHILLIPS.
1973. Studies on the migration of *Tritaeta gibbosa*, a subtidal benthic amphipod. *Neth. J. Sea Res.* 7:135-149.
- KETCHEN, K. S., AND C. R. FORRESTER.
1966. Population dynamics of the petrale sole, *Eopsetta jordani*, in the waters of western Canada. *Fish. Res. Board Can., Bull.* 153, 195 p.
- MACARTHUR, R. H., AND E. R. PIANKA.
1966. On optimal use of a patchy environment. *Am. Nat.* 100:603-609.
- NORMAN, J. R.
1934. A systematic monograph of the flatfishes (Heterosomata). Vol. I. Psettodidae, Bothidae, Pleuronectidae. *Br. Mus. (Nat. Hist.), Lond.*, 459 p.
- PEARCY, W. G., AND H. A. VANDERPLOEG.
1973. Radioecology of benthic fishes off Oregon. In *Radioactive contamination of the marine environment*, p. 245-261. Int. At. Energy Agency, Vienna.
- RAE, B. B.
1969. The food of the witch. *Mar. Res. Dep. Agric. Fish. Scotl.* 2, 23 p.
- SHUBNIKOV, D. A., AND L. A. LISOVENKO.
1964. Data on the biology of rock sole of the southeastern Bering Sea. *Tr. Vses. Nauchno-issled. Inst. Morsk. Rybn. Khoz. Okeanogr.* 49 (Izv. Tikhookean. Nauchno-issled. Inst. Morsk. Rybn. Khoz. Okeanogr. 51): 209-214. (Transl. in *Soviet Fisheries Investigations in the Northeast Pacific, Part II*, p. 220-226, by Israel Program Sci. Transl., 1968, available Natl. Tech. Inf. Serv., Springfield, VA, as TT 67-51204.)
- STICKNEY, R. R., G. L. TAYLOR, AND R. W. HEARD III.
1974. Food habits of Georgia estuarine fishes. I. Four species of flounders (Pleuronectiformes: Bothidae). *Fish. Bull., U.S.* 72:515-525.
- TYLER, A. V.
1973. Alimentary tract morphology of selected North Atlantic fishes in relation to food habits. *Fish. Res. Board Can., Tech. Rep.* 361, 23 p.
- YAZDANI, G. M.
1969. Adaptations in the jaws of flatfish (Pleuronectiformes). *J. Zool. (Lond.)* 159:181-222.

MICHAEL J. KRAVITZ
WILLIAM G. PEARCY
M. P. GUIN

*School of Oceanography
Oregon State University
Corvallis, OR 97331*



The Oregon State Sea Grant College Program is supported cooperatively by the National Oceanic and Atmospheric Administration, U.S. Department of Commerce, by the State of Oregon, and by participating local governments and private industry.