

AN ABSTRACT OF THE THESIS OF

Paul Joseph Alvin Presidente for the DOCTOR OF PHILOSOPHY
(Name of student) (Degree)

in Zoology presented on May 5, 1972
(Major department) (Date)

Title: PATHOGENICITY OF EXPERIMENTALLY INDUCED
CONCURRENT INFECTIONS OF FASCIOLA HEPATICA
AND HAEMONCHUS CONTORTUS IN SHEEP

Abstract approved: Redacted for privacy
Stuart E. Knapp

Twenty yearling sheep were allotted to four groups, each containing five sheep of approximately equal weight. On experimental days 0, 1, 2, 3, and 4, each sheep in each group was given the following inoculations by stomach tube: five sheep in group I (Fasciola hepatica-exposed controls) were given approximately 120 F. hepatica metacercariae each day; five sheep in group II (Haemonchus contortus-exposed controls) were given approximately 10,000 H. contortus infective larvae (L_3) each day; five sheep in group III (F. hepatica and H. contortus-exposed principals) were given approximately 120 F. hepatica metacercariae and 10,000 H. contortus L_3 each day; five sheep in group IV (nonexposed controls) were given sham doses of tap water each day.

Each sheep was weighed at seven-day intervals until experimental day 203 and a fecal sample collected until day 105.

Clinicopathologic changes in blood were analyzed in samples collected from each sheep at seven-day intervals. Sheep that died in groups I and III were necropsied; gross pathology was recorded, helminths recovered, and liver tissue taken for histopathologic examination. Two sheep in group I were killed and examined on experimental days 269 and 1,063.

Clinicopathologic and pathologic changes in F. hepatica-exposed sheep (group I) were characteristic of chronic fascioliasis. Eosinophilia, leukocytosis, hypoalbuminemia, and progressive macrocytic, normochromic anemia occurred before time of death. One sheep each died on experimental days 162, 168, and 265; 412, 414, and 318 mature F. hepatica were recovered in liver bile ducts. Fluke burdens in two sheep that were killed and necropsied on experimental days 269 and 1,063 were 383 and 28, respectively.

Resistance to H. contortus exposure was evident in group II sheep; four of five sheep acquired patent infections and two of these were negative for nematode eggs in feces on experimental day 70. Transitory mild eosinophilia and hypoalbuminemia, and normocytic, hypochromic anemia was found, but each sheep recovered naturally within 203 days.

Enhanced pathogenicity of simultaneous exposure to F. hepatica and H. contortus in sheep was demonstrated by death of three sheep in group III before experimental day 80. The two remaining sheep in group III died two months before two sheep in group I that harbored similar fluke burdens. A macrocytic, slightly hypochromic anemia was found in one sheep that died on experimental day 51; 253 immature F. hepatica and 28,000 H. contortus were recovered at necropsy. Slightly microcytic, hypochromic anemia, characteristic of H. contortus infection, was found in two sheep that died on experimental days 71 and 78. In one sheep, 238 F. hepatica and 25,300 H. contortus were recovered; the other sheep harbored 332 and 18,100 helminths, respectively. The two remaining sheep died on experimental days 190 and 201; anemia was macrocytic and normochromic, characteristic of chronic fascioliasis. Burdens were 346 and 326 mature F. hepatica; 12,600 and 27,600 H. contortus were recovered.

Establishment of both helminths and host response to F. hepatica infection appeared normal, but development of host resistance to H. contortus was inhibited by simultaneous F. hepatica infection. Eosinophilic response in sheep with the concurrent infections was less than that of sheep with F. hepatica infection alone.

© 1972

PAUL JOSEPH ALVIN PRESIDENTE

ALL RIGHTS RESERVED

Pathogenicity of Experimentally Induced Concurrent
Infections of Fasciola hepatica and Haemonchus
contortus in Sheep

by

Paul Joseph Alvin Presidente

A THESIS

submitted to

Oregon State University

in partial fulfillment of
the requirements for the
degree of

Doctor of Philosophy

June 1972

APPROVED:

Redacted for privacy

Professor of Veterinary Parasitology
in charge of major

Redacted for privacy

Chairman of Department of Zoology

Redacted for privacy

Dean of Graduate School

Date thesis is presented May 5, 1972

Typed by Velda D. Mullins for Paul Joseph Alvin Presidente

ACKNOWLEDGMENT

The author expresses his sincere gratitude to Dr. Stuart E. Knapp for his advice and guidance throughout this investigation, and for his constructive criticism of this dissertation.

Cooperation of the following individuals is gratefully acknowledged: Keith D. Nicol, presently a veterinary student at Ohio State University, for assistance with hematologic analyses of blood samples; Dr. Stanley P. Snyder, pathologist, Department of Veterinary Medicine, Oregon State University, for assistance with the histopathologic portion of the study; Norman L. Baldwin, M.S., Oregon State University, for providing Fasciola hepatica metacercariae used in the experiment.

Gratitude is expressed to Dr. J. N. Shaw, Department of Veterinary Medicine, and Dr. Ivan Pratt, Department of Zoology, Oregon State University for their interest and encouragement throughout the course of this investigation.

I also express appreciation to my family for their encouragement and patience throughout this experiment.

TABLE OF CONTENTS

	<u>Page</u>
INTRODUCTION	1
REVIEW OF LITERATURE	3
Acute Ovine Fascioliasis	3
Subacute Ovine Fascioliasis	5
Chronic Ovine Fascioliasis	6
Ovine Haemonchosis	10
Phlebotomy vs Helminth-Caused Anemias	13
MATERIALS AND METHODS	17
Animals	17
Inoculum	17
Metacercariae	17
Nematode larvae	18
Experimental Design	18
Helminth Egg Counts	19
Hematologic and Serologic Procedures	19
Necropsy Procedure	20
RESULTS	22
Body Weight Data	22
Helminth Egg Counts	24
Clinicopathologic Findings	26
Erythrocytes	26
Leukocytes	37
Serum Albumin	41
Postmortem Findings	44
Gross Pathology	44
Helminthologic Data	46
Histopathology	47
DISCUSSION	50
BIBLIOGRAPHY	54

LIST OF TABLES

<u>Table</u>		<u>Page</u>
1	Mean Wintrobe Indices for Sheep	36
2	Final Weight, Packed Cell Volume, Fecal Sample Analysis, and Helminth Recovery Data for each of 5 <u>Fasciola hepatica</u> -exposed sheep (group I), and 5 <u>F. hepatica</u> and <u>Haemonchus contortus</u> -exposed sheep (group III)	45

LIST OF FIGURES

<u>Figure</u>		<u>Page</u>
1	Mean body weights for 5 <u>Fasciola hepatica</u> -exposed sheep (group I), 5 <u>Haemonchus contortus</u> -exposed sheep (group II), and 5 nonexposed sheep (group IV)	23
2	Body weights for each of 5 <u>Fasciola hepatica</u> and <u>Haemonchus contortus</u> -exposed sheep (group III)	25
3	<u>Haemonchus contortus</u> fecal egg counts for each of 5 <u>Fasciola hepatica</u> and <u>H. contortus</u> -exposed sheep (group III), and mean counts for 5 <u>H. contortus</u> -exposed sheep (group II)	27
4	Mean packed cell volumes for 5 <u>Fasciola hepatica</u> -exposed sheep (group I) 5 <u>Haemonchus contortus</u> -exposed sheep (group II), 5 <u>F. hepatica</u> and <u>H. contortus</u> -exposed sheep (group III), and 5 nonexposed sheep (group IV)	28
5	Packed cell volumes for each of 5 <u>Fasciola hepatica</u> and <u>Haemonchus contortus</u> -exposed sheep (group III)	29
6	Mean erythrocyte counts for 5 <u>Fasciola hepatica</u> -exposed sheep (group I), 5 <u>Haemonchus contortus</u> -exposed sheep (group II), 5 <u>F. hepatica</u> and <u>H. contortus</u> -exposed sheep (group III), and 5 non-exposed sheep (group IV)	30
7	Erythrocyte counts for each of 5 <u>Fasciola hepatica</u> and <u>Haemonchus contortus</u> -exposed sheep (group III)	31
8	Mean hemoglobin concentrations for 5 <u>Fasciola hepatica</u> -exposed sheep (group I), 5 <u>Haemonchus contortus</u> -exposed sheep (group II), 5 <u>F. hepatica</u> and <u>H. contortus</u> -exposed sheep (group III) , and 5 nonexposed sheep (group IV)	32

<u>Figure</u>		<u>Page</u>
9	Hemoglobin concentrations for each of 5 <u>Fasciola hepatica</u> and <u>Haemonchus contortus</u> -exposed sheep (group III)	33
10	Mean leukocyte total and eosinophil counts for 5 <u>Fasciola hepatica</u> -exposed sheep (group I) 5 <u>Haemonchus contortus</u> -exposed sheep (group II), 5 <u>F. hepatica</u> and <u>H. contortus</u> -exposed sheep (group III), and 5 nonexposed sheep (group IV)	38
11	Eosinophil counts for each of 5 <u>Fasciola hepatica</u> and <u>Haemonchus contortus</u> -exposed sheep (group III)	40
12	Mean serum albumin concentrations for 5 <u>Fasciola hepatica</u> -exposed sheep (group I), 5 <u>Haemonchus contortus</u> -exposed sheep (group II), 5 <u>F. hepatica</u> and <u>H. contortus</u> -exposed sheep (group III), and 5 nonexposed sheep (group IV)	42
13	Serum albumin concentrations for each of 5 <u>Fasciola hepatica</u> and <u>Haemonchus contortus</u> -exposed sheep (group III)	43

PATHOGENICITY OF EXPERIMENTALLY INDUCED
CONCURRENT INFECTIONS OF FASCIOLA
HEPATICA AND HAEMONCHUS
CONTORTUS IN SHEEP

INTRODUCTION

Pathogenesis and pathology of Fasciola hepatica in experimentally induced chronic infections in sheep have been investigated by several workers (Boray, 1967; Dow, Ross, and Todd, 1968; Furmaga and Gundlach, 1967a, b; Ross, Dow, and Todd, 1967; Rubaj and Furmaga, 1969; Sewell, Hammond, and Dinning, 1968; Sinclair, 1962, 1964; Symons and Boray, 1967, 1968). Naturally occurring epizootics of ovine fascioliasis have also been examined (Hjerpe et al., 1971; Reid et al., 1970; Ross, 1967a, b).

Considerable research on Haemonchus contortus infections in sheep has been done (Andrews, 1942; Charleston, 1965; Fourie, 1931; Malczewski, 1970; Silverman, Mansfield, and Scott, 1970; Stoll, 1943; Veglia, 1915). Clinicopathologic changes in sheep with mixed gastrointestinal nematode burdens have been extensively investigated (Baker and Douglas, 1966; Baker et al., 1952; Campbell and Gardiner, 1960; Holman and Pattison, 1941; Kuttler and Marble, 1960; Shumard, Bolin, and Eveleth, 1957; Whitlock, 1950; Wilson and Turner, 1965). Enhanced pathogenicity was reported when lambs were simultaneously exposed to H. contortus and Nematodirus

spathiger; Nematodirus infection was prolonged in these infections (Kates and Turner, 1960; Turner and Colglazier, 1954). Previous massive exposure to H. contortus larvae caused reduction in numbers recovered and retardation in maturation rate of Nematodirus battus (Mapes and Coop, 1970). Competitive exclusion of H. contortus by simultaneous exposure to Trichostrongylus axei was reported (Turner, Kates, and Wilson, 1962).

Clinicopathology in young calves with concurrent infections of F. hepatica and Ostertagia ostertagi was recently reported (Reid et al., 1967). Pathogenesis of simultaneous F. hepatica and Haemonchus infections in sheep or cattle has not been studied.

The purpose in the present study was to investigate clinicopathologic changes in sheep after simultaneous exposure to F. hepatica and H. contortus.

REVIEW OF LITERATURE

Since the life cycle of F. hepatica was independently described by Leuckart and Thomas in 1883, the common liver fluke has been intensively investigated. Early experiments with F. hepatica were reviewed by Gordon (1955) and Dawes and Hughes (1964). Comprehensive literature reviews on the biology of F. hepatica were provided by Taylor (1964) and Pantelouris (1965). Pathogenesis of liver flukes in domestic and laboratory animals has been summarized by Sewell (1966) and Sinclair (1967a). Boray (1969) compiled a review of data obtained from experimentally induced infections in sheep and cattle in Australia. Recent literature on fascioliasis in cattle and anemia in ovine infections was discussed by Dawes and Hughes (1970). A comprehensive description of clinical symptoms and pathogenesis of human fascioliasis was recently provided by Náquira-Vildoso and Marcial-Rojas (1971).

Due to considerable variation in host response to F. hepatica (Dawes and Hughes, 1964; Dow, Ross, and Todd, 1968; Sinclair, 1969), this review is restricted to a discussion of current knowledge of acute, subacute, and chronic fascioliasis in sheep.

Acute Ovine Fascioliasis

Recent experiments by several workers (Boray, 1967; Pullan,

Sewell, and Hammond, 1970; Roberts, 1968; Ross, Dow, and Todd, 1967) have clarified the pathogenesis of this disease in sheep. In the acute disease, death occurred 42 to 62 days after exposure to 4,000 to 12,000 F. hepatica metacercariae. Before time of death, symptoms included inappetance, marked weight loss, emaciation, constipation, severe ascites, and anemia.

Clinicopathologic changes observed were normocytic or macrocytic, normochromic anemia beginning five to six weeks after exposure. Increased erythropoiesis was indicated by appearance in peripheral blood of erythrocytes showing anisocytosis, polychromasia, and basophilic stippling; hyperplasia of the red bone marrow occurred. Symons and Boray (1967, 1968) used ^{59}Fe -labeled plasma to demonstrate rapid movement of iron from bone marrow to circulating erythrocytes in sheep six and eight weeks after exposure. Appearance of poikilocytes in circulating blood before time of death suggested exhaustion of hematopoietic tissues. The major change in leukocyte differential counts was a marked eosinophilia that began two weeks after exposure; Roberts (1968) reported that neutrophilia occurred before death. Hepatic dysfunction was indicated by retarded bromsulphthalien (BSP) clearance (Roberts, 1968), elevated serum glutamic-oxaloacetic transaminase (SGOT) and glutamic dehydrogenase (SGD) levels (Pullan, Sewell, and Hammond, 1970; Sewell, 1967a), icterus of the plasma (Symons and Boray, 1968),

and hypoalbuminemia. Serum total protein levels increased within two weeks after exposure; hypergammaglobulinemia was primarily involved (Pullan, Sewell, and Hammond, 1970; Sewell, 1970).

Postmortem examination revealed severe ascites with moderate to large quantities of blood; the carcass was wet and anemic. Rupture of the liver capsule with hemorrhage into the peritoneal cavity was given as cause of death (Boray, 1967). A generalized, fibrinous peritonitis and perihepatitis were evident. The liver was enlarged, congested, and soft; numerous subcapsular hemorrhagic foci were observed. Histopathologic changes in parenchyma associated with migration pathways of immature flukes were described by Dow, Ross, and Todd (1968). Dissection of liver parenchyma revealed 1, 000 to 3, 300 immature F. hepatica. Growth rate of these flukes was retarded and only a few had entered the bile ducts by time of death.

Subacute Ovine Fascioliasis

This disease differs from acute fascioliasis in time that death occurs, number of flukes that become established, and their stage of maturation. Death occurred 56 to 110 days after exposure; a majority of the 500 to 2, 000 flukes were recovered in bile ducts (Pullan, Sewell, and Hammond, 1970; Ross, Dow, and Todd, 1967).

Progressive weight loss and anemia began five weeks after exposure. Liver function was improved and eosinophilia less marked than was found during the acute stage.

At necropsy of the sheep, only small amounts of peritoneal fluid were noted and fibrinous peritonitis was less evident. The liver surface was uneven, firm, and biliary cirrhosis had begun (Soulsby, 1965). The reparative tissue reaction prevented fluke entry into bile ducts and extended the migratory period in liver parenchyma (Boray, 1967). Increased size of flukes intensified the mechanical damage causing increased hemorrhage.

In a study of naturally acquired F. hepatica infections in lambs, Reid et al. (1970) reported development of a severe macrocytic, hypochromic anemia. A marked reticulocytosis (8 to 30%) was observed in lambs when packed cell volume (PCV) dropped below 25%. Similar findings were reported by Hjerpe et al. (1971) in an investigation of an epizootic in California among a flock of ewes. Clinicopathologic changes associated with the hypochromic anemia included reductions in serum iron content, bone marrow hemosiderin reserves, and mean corpuscular hemoglobin concentration (MCHC).

Chronic Ovine Fascioliasis

This disease has been investigated by workers who examined sheep with experimentally induced infections (Boray, 1967; Furmaga

and Gundlach, 1967a, b; Ibrović and Gall-Palla, 1959; Ross, Dow, and Todd, 1967; Sinclair, 1962, 1964), and those that acquired F. hepatica by natural exposure (Ross, 1967a, b). The chronic disease resulted from exposure to 100 to 2,000 F. hepatica metacercariae and establishment of 100 to 1,000 flukes in bile ducts. Time of death correlated with fluke burden, and sheep began dying 16 weeks after exposure. Durbin (1952) reported that in one sheep a patent infection lasted 11 years; 15 F. hepatica were observed from the bile ducts. Economic losses due to chronic fascioliasis were discussed by Gordon (1955), and Roseby (1970) demonstrated a deleterious effect on wool production. Symptoms of chronic fascioliasis began 8 to 14 weeks after exposure, and included increased thirst, a progressive anemia, loss of weight and condition; intermandibular edema and ascites were observed. The anemia that developed was severe, and PCV was as low as 4% at time of death (Boray, 1967). This anemia was normocytic or slightly macrocytic, and normochromic (Furmaga and Gundlach, 1967a; Sewell, Hammond, and Dinning, 1968; Sinclair, 1962, 1964, 1965). Compensatory erythropoiesis was indicated by anisocytosis, basophilic stippling, and poikilocytosis in circulating erythrocytes (Sewell, Hammond, and Dinning, 1968), red bone marrow hyperplasia (Sewell, Hammond, and Dinning, 1968; Sinclair, 1964), and increased rate of plasma iron clearance and incorporation into

circulating erythrocytes (Sinclair, 1964; Symons and Boray, 1967, 1968). Sinclair (1965) observed a reduction in hemosiderin reserves in bone marrow beginning 87 days postexposure and decreased serum iron concentration at 147 days. A marked decrease in erythrocyte count the following week indicated iron deficiency was the limiting factor in erythropoiesis. Symons and Boray (1967, 1968) used ^{59}Fe -labeled plasma to demonstrate that newly produced erythrocytes had a six-day half-life in peripheral circulation. Labeled iron was recovered in bile and feces of infected sheep. Blood loss in feces correlated with degree of anemia observed when ^{51}Cr -labeled erythrocytes were injected into infected sheep (Holmes *et al.*, 1967; Sewell, Hammond, and Dinning, 1968). Sewell (1967b) estimated the daily blood loss as 0.5 to 1.0 ml. per fluke. Intestinal reabsorption of iron lost through bile ducts was investigated by simultaneous labeling of erythrocytes with ^{51}Cr and ^{59}Fe (Holmes and MacLean, 1969). When anemia was hypochromic and host iron reserves were depleted, some reabsorption of labeled iron occurred.

High radioactivity was recorded for expelled ingesta of F. hepatica recovered from sheep previously injected with erythrocytes labeled with ^{51}Cr or ^{59}Fe (Pearson, 1963; Symons and Boray, 1967, 1968). The ingesta contained hematin (Stephenson, 1947; Todd and Ross, 1966), and acetylcholinesterase activity characteristic of host erythrocytes (Frady and Knapp, 1967); the majority

of workers have concluded that mature F. hepatica were almost entirely hematophagic. Dawes (1963) and Dawes and Hughes (1964, 1970) criticized some of this work because direct feeding on blood by flukes was physically impossible; the hyperplastic bile duct epithelium separated the fluke from a blood source.

During the chronic stage of fascioliasis, serum total protein decreased because a marked hypoalbuminemia offset the hyperglobulinemia that occurred during the acute stage (Furmaga and Gundlach, 1967b; Ibrović and Gall-Palla, 1959; Sinclair, 1962). Holmes et al. (1968) used ^{131}I -labeled albumin in infected sheep to demonstrate a hypercatabolism of albumin; the shortened half-life was associated with a loss of albumin into the intestinal tract via the bile. They found similar loss of gammaglobulin when ^{125}I -labeled globulin-7s was used. Effect of corticosteroid administration on serum protein in infected sheep was examined by Sinclair (1967b). Hyperglobulinemia did not occur and hypoalbuminemia began earlier and was more marked compared to changes in infected, nonmedicated controls.

Serum calcium levels decreased during the chronic stage of infection; nonsignificant changes in serum magnesium, phosphorous and potassium were recorded (Pinkiewitz and Madej, 1967; Sinclair, 1960). Increased serum alkaline phosphatase activity was reported (Hjerpe et al., 1971; Pinkiewitz and Madej, 1967).

Postmortem examination during chronic infection revealed that liver parenchyma was fibrotic and firm, especially the left lobe; main bile ducts were prominent, thickened, and fibrous (Rubaj and Furmaga, 1969); Soulsby, 1965). Histopathologic changes in liver at chronic stage were described by Dow, Ross and Todd (1968) and Rubaj and Furmaga (1969).

Ovine Haemonchosis

Pathogenesis and pathology of H. contortus infection in sheep were reviewed recently by Soulsby (1965). Symptoms associated with this infection were principally those caused by anemia. Gordon (1950) reported the following: lack of stamina, pale mucus membranes, intermandibular edema, constipation with hard, dark feces; milk yield was reduced in lactating ewes. Variation in response to exposure was primarily related to condition of the host. Factors that affected outcome of exposure to H. contortus larvae included: age of the sheep (Christie, 1970; Lucker, 1952; Poeschel and Todd, 1969; Silverman, Mansfield, and Scott, 1970); host size, weight, and hematologic status (Gordon, 1950); diet of the sheep (Kates, Allen, and Wilson, 1962; Poeschel and Todd, 1969; Shumard et al., 1956; Wier et al., 1948); the hemoglobin type of the sheep (Evans, Blunt, and Southcott, 1963); and number, dosing schedule, and strain of infective larvae used (Conway and Whitlock, 1965; Dineen

et al., 1965; Knapp, 1964a, b). Lambs over eight months of age compensated for blood loss by moderate worm burdens and recovered within 90 days after exposure (Lucker, 1952).

The primary clinicopathologic change in ovine haemonchosis was a marked anemia. Blood was observed in feces beginning 6 to 10 days after exposure; gastric hemorrhage was caused by 4th-stage larvae (Andrews, 1942; Brambell, Charleston, and Tothill, 1964; Clark, Kiesel, and Goby, 1962; Stoll, 1943; Veglia, 1915). A rapid decline in hemoglobin concentration and PCV began six days after exposure; PCV was 6% at time of death 19 to 24 days post-exposure (Lucker, 1952). Blood loss was greater when H. contortus was mature. Brambell, Charleston, and Tothill (1964) used ^{51}Cr -labeled erythrocytes to demonstrate that 30 ml. of blood was lost in feces eight days after exposure; 500 ml. was lost in feces of one lamb on postexposure day 23. Andrews (1942) measured blood loss in feces of infected lambs; one lamb lost 2,380 ml. in a 10-day period. Blood loss was due to capillary hemorrhage from vacated attachment sites, and to ingestion by the mature worms (Andrews, 1942; Boughton and Hardy, 1935; Fourie, 1931; Veglia, 1915). Martin and Clunies Ross (1934) estimated that for egg production alone, 2,000 female worms required 29 ml. of blood each day. Clark, Kiesel, and Goby (1962) used erythrocytes labeled with ^{51}Cr alone or in combination with ^{59}Fe to estimate

mean blood loss as 0.049 ml. per worm per day.

Normocytic or macrocytic, normochromic or microcytic, hypochromic anemia were described in haemonchosis depending on stage of infection (Baker and Douglas, 1966; Fourie, 1931). Increased erythropoiesis was indicated by compensatory myeloid hyperplasia, and anisocytosis, polychromasia, and basophilic stippling in circulating erythrocytes (Baker and Douglas, 1966; Fourie, 1931; Schalm, 1965). Exhaustion of hematopoietic tissues was indicated when poikilocytes and hypochromatic cells appeared in peripheral blood; decreased mean corpuscular volume (MCV) and MCHC suggested iron deficiency (Baker and Douglas, 1966; Fourie, 1931; Schalm, 1965; Silverman, Mansfield, and Scott, 1970).

Changes observed in leukocyte total and differential counts during haemonchosis were leukocytosis (Andrews, 1942), a relative lymphocytosis (Silverman, Mansfield, and Scott, 1970), and a relative neutrophilia at death (Andrews, 1942). A marked eosinophilia was recorded 6 to 14 days after exposure, especially if larvae were given in daily doses (Charleston, 1965; Malczewski, 1970). Fourie (1931) and Andrews (1942) reported increased eosinophil counts during the recovery phase of infection.

In sheep with mixed gastrointestinal nematode burdens (including H. contortus), hyperglobulinemia and hypoalbuminemia have

been reported (Kuttler and Marble, 1960; Shumard, Bolin, and Eveleth, 1957).

Pathogenesis and pathology of haemonchosis have been studied by several workers (Andrews, 1942; Charleston, 1965; Malczewski, 1970; Stoll, 1943; Veglia, 1915). Migration of larvae to the region of abomasal lamina propria to molt to 4th-stage caused an intense lymphoid infiltration into the area (Charleston, 1965; Malczewski, 1970; Stoll, 1943). Migration of 4th-stage larvae to mucosal surface caused extensive damage to gastric glands and the mucosae; pH increased (Christie, 1970; Christie, Brambell, and Mapes, 1967; Malczewski, 1970), and a marked eosinophilic infiltration into the mucosae occurred (Charleston, 1965; Malczewski, 1970). Lesions were primarily restricted to the mucosae; excessive sloughing and mucus secretion occurred. Hypertrophy of the mucus gland portion occurred and numerous petechial hemorrhages associated with former attachment sites of worms were evident.

Phlebotomy vs Helminth-Caused Anemias

Anemia in both haemonchosis and fascioliasis in sheep is apparently posthemorrhagic in nature, and several investigators have attempted to duplicate the condition in infected sheep by repeated phlebotomy. Todd and Ross (1968) reported a macrocytic, normochromic anemia with marginal hypochromasia

occurred in five sheep after removal of 300 ml. of blood for six consecutive days. Significant changes in leukocyte total and differential counts did not occur. Fourie (1931), Andrews (1942), and Richard et al. (1954) found changes in peripheral blood were similar in sheep infected with H. contortus to those observed in phlebotomized sheep. Charleston (1964) studied bone marrow changes in H. contortus-infected sheep and in phlebotomized sheep. He concluded that resulting anemias were not identical; interference in normoblast maturation and hemoglobinization process in infected sheep was indicated. Also, a slight increase in eosinophil counts in infected sheep was not recorded for sheep that were artificially bled.

Kuttler and Marble (1960) reported significant differences in degree of hypoalbuminemia and hyperalphaglobulinemia observed in sheep with mixed gastrointestinal nematode burdens (including H. contortus) when compared to that found in phlebotomized sheep.

Sinclair (1964, 1965) attempted to duplicate the anemia in chronic fascioliasis by repeated phlebotomy. Because of observed differences between infected sheep and phlebotomized sheep, he concluded that a dysfunction of the reticuloendothelial system was the primary cause of anemia. This was characterized by increased erythrocyte destruction and decreased effective erythrocyte production. Other workers (Sewell, Hammond, and Dinning, 1968;

Symons and Boray, 1968, disagreed with this conclusion because they found erythropoiesis was increased, possibly to a maximum. Sinclair investigated the role of the reticuloendothelial system in sheep with experimentally induced F. hepatica infections. Effect of corticosteroid administration (Sinclair, 1968) and splenectomy (Sinclair, 1970a) on the course of primary infection, and corticosteroid administration on secondary infection (Sinclair, 1970b) were investigated. He concluded that hyperactivity of the reticuloendothelial system was an integral part of the resistance mechanism to F. hepatica infection in sheep. Interference with this system resulted in more severe disease because the normal cellular reaction in the liver was suppressed. Sinclair (1970b) then concluded that the anemia in fascioliasis was primarily hemorrhagic in origin.

Clinicopathologic changes associated with conditions discussed are summarized as follows:

	Fascioliasis			Nematodes		
	acute	sub-acute	chronic	haemon- chosis	mixed infec- tion	Phle- botomy
Weight loss	+++	++	++	+, ++	++	++
Edema	+	+	++	++	NC	NC
Ascites	+++	++	+, ++	NC	NC	NC
Anemia						
normocytic	++	+	++	+	++	
macrocytic	+	++	+, ++	+		++
microcytic				++, +++	+	
normochromic	++	++	++	+	++	++
hypochromic		+		++, +++	+	+
Abnormal red cells	+++	+++	+++	+++	++	++
Leukocytosis	NC	++	++	+	+	NC
Eosinophilia	+++	+	+	+, ++	+	NC
Neutrophilia	++	+	+	++	+	NC
Hypoalbuminemia	+	+++	+++	++	++	+
Hyperglobulinemia	+++	++	++	++	++	+
Bromsulphthalein retention	+++	++	+	NC	NC	NC
SGOT, SGD levels (elevated)	+++	++	+	NC	NC	NC
Alkaline phosphatase (elevated)	+++	++	+	NC	NC	NC

NC means no change; + slight change; ++ moderate change; +++ severe change.

MATERIALS AND METHODS

Animals

Twenty sheep, of mixed breeding and 12 to 18 months old, were housed indoors since time of birth in pens with concrete floors. Fecal samples collected from each sheep before the experiment began indicated small numbers of nematode eggs in feces of four sheep; none were found in remaining sheep. The sheep were given a pelleted ration consisting of 62% grass hay, 21% alfalfa, 12% rolled barley, and 5% molasses.

Inoculum

Metacercariae

Fasciola hepatica metacercariae were collected from Lymnaea columella, a suitable intermediate host, that was artificially exposed to miracidia hatched from ova. Fluka ova were removed from gallbladder and liver bile ducts of sheep naturally infected with F. hepatica. The number of viable metacercariae was determined by dilution count, and 50 doses containing approximately 120 cysts were drawn from the total collection. The inoculum of 600 metacercariae was given to exposed sheep by stomach tube in five equal doses given on consecutive days.

Nematode Larvae

Haemonchus contortus infective larvae (L₃) were recovered from moist, fecal cultures after 27°C. incubation for six days. Nematode eggs were recovered in feces from lambs with experimentally induced H. contortus infections. The Kentucky strain of H. contortus was used; an isolate was obtained in 1960 and maintained here by serial passage in susceptible lambs. The number of infective larvae was determined by dilution count, and 50 doses containing approximately 10,000 L₃ were drawn from the total collection. The inoculum of 50,000 L₃ was given to exposed sheep by stomach tube in five equal doses given on consecutive days.

Experimental Design

On May 1, 1969, the sheep were allotted by body weight to four groups of five sheep each by the method of Gardiner and Wehr (1950). Beginning on May 6 (experimental day 0), and on days 1, 2, 3, and 4, each sheep in the groups was given the following inoculations: five F. hepatica-exposed controls (group I) were given 120 metacercariae each day; five H. contortus-exposed controls (group II) were given 10,000 L₃ each day; five F. hepatica and H. contortus-exposed principals (group III) were given 120 metacercariae and then 10,000 L₃ each day; five nonexposed controls (group IV) were

given sham doses of water each day. Each group was kept in separate pens with concrete floors, and each sheep was weighed at seven-day intervals until time of death, or until experimental day 203.

Helminth Egg Counts

Fecal samples were collected every seven days from each sheep from experimental day 0 until day 105. Presence of F. hepatica ova in these samples was demonstrated by sedimentation. Quantitative estimate of nematode egg output in feces was done by the modified McMaster technique; counts were expressed as number of eggs per gram of feces (e. p. g.).

Hematologic and Serologic Procedures

Blood samples were collected at seven-day intervals from the jugular vein of each sheep until time of death, or until experimental day 203. Whole blood (five ml.) was collected for serum separation, and five ml. was collected in ethylenediaminetetraacetic acid. Blood smears were prepared and stained with Wright's solution for examination of erythrocytes and making leukocyte differential counts. Packed cell volume (PCV) percentage was determined by microhematocrit. For the first 19 sets of blood samples, total erythrocyte and leukocyte counts were made; hemoglobin concentration was

determined by the biuret method. Mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), and mean corpuscular hemoglobin concentration (MCHC), were calculated by methods given by Schalm (1965). Serum samples were frozen and albumin concentration was subsequently determined by the biuret method.

Necropsy Procedure

Eight sheep that died and two sheep that were killed were examined for gross pathologic changes; liver tissue was taken for histologic preparation. Sectioned tissues were stained in hematoxylin and eosin for histopathologic examination. Liver from each sheep was collected; gallbladder contents were poured into a beaker and the sediment was examined for fluke ova. Bile ducts were opened and examined for mature flukes. Each liver was then cut in thin strips, placed in a tray, and soaked in warm tap water for approximately two hours. All tissue was examined and washed over a sieve (1.4-mm. openings) before being discarded. Contents in the bottom of the tray were poured over the sieve to recover flukes that emerged from hepatic tissue. Flukes recovered from each liver were placed in petri dishes at 4°C. and allowed to relax overnight. The number of recovered flukes was counted and average length determined by measurement of a 10% random sample. Whole flukes were flattened between glass plates and their lengths measured.

The abomasum was collected from each sheep in group III. Contents and adherent worms were recovered by rubbing the mucosae with the fingertips and washing over a sieve (0.147-mm. openings); material on the sieve was collected and fixed in formalin. Abomasum contents were diluted to a volume of one L. and all worms in a five ml. aliquot were counted. Total H. contortus burden was estimated by multiplying the aliquot count by the dilution factor of 200.

RESULTS

Body Weight Data

On experimental day 0, mean body weight for group I sheep (F. hepatica-exposed controls) was 44.2 kg., for group II sheep (H. contortus-exposed controls), 43.1 kg., for group III sheep (F. hepatica and H. contortus-exposed principals), 42.7 kg., and for group IV sheep (nonexposed controls), 42.7 kg. Weekly mean body weight for sheep in the three control groups are shown (Figure 1); differences between groups were not significant ($P > 0.05$) at any time during the experiment. Each sheep in group IV gained an average 18.3 kg. (minimum and maximum, 15.2 and 23.9) by experimental day 203, and each sheep in group II, 16.3 kg (minimum and maximum, 11.6 and 18.9). Sheep 2 (34.1 kg. on experimental day 0) and sheep 4 (45.0 kg. on day 0) in group I died of chronic fascioliasis on experimental days 162 and 168; both sheep lost approximately 2.5 kg. in the 14-day period before they died (Figure 1). The other three group I sheep averaged 47.5 kg. on day 0, and gained an average 8.9 kg. by day 203. Sheep 1 died on experimental day 265; sheep 3 and 5 were killed and examined on days 269 and 1,063, respectively. These sheep lost or gained little weight in the period between experimental days 140 and 203. Sheep 3 weighed 51.8 kg. on




Figure 1. Mean body weights for 5 *Fasciola hepatica*-exposed sheep (group I; ●—●), 5 *Haemonchus contortus*-exposed sheep (group II; ■—■), and 5 nonexposed sheep (group III; ○—○).

experimental day 203 and 50.0 kg. at necropsy on day 269. Sheep 5 subsequently gained weight and put on considerable fat; it weighed 93.6 kg. at necropsy.

Each sheep in group IV (F. hepatica and H. contortus-exposed principals) died between experimental days 51 and 201; weekly weights are given (Figure 2). Three sheep that died before day 80 showed progressive weight loss for the 28-day period before time of death; two sheep that died on days 190 and 201 lost or gained little weight in the 56-day period before they died.

Helminth Egg Counts

Ova of F. hepatica were recovered in feces from one sheep in each of groups I (F. hepatica-exposed controls) and III (F. hepatica and H. contortus-exposed principals) on experimental day 70. Each of five group I sheep and three of four group III sheep had patent infections on day 77. Each of eight F. hepatica-exposed sheep in groups I and III passed large numbers of fluke ova in feces until time of death or necropsy. Output of fluke ova in sheep 5 (group I) was 16.1 ova per gram of feces on experimental day 1, 063. Fluke ova were not recovered in feces from sheep in groups II and IV on any sample day.

Haemonchus contortus egg counts for each sheep in group III (F. hepatica and H. contortus-exposed principals) and mean count




Figure 2. Body weights for each of 5 *Fasciola hepatica* and *Haemonchus contortus*-exposed sheep (group III).

for group II sheep (H. contortus-exposed controls) are shown (Figure 3). Four sheep in each of groups II and III had egg counts of 100 to 550 e. p. g. on experimental day 21. Sheep 12 (group III) passed 200 e. p. g. on day 28; none were found in feces from sheep 9 (group II) on any sample day. Mean egg output for group II sheep peaked at 2,790 e. p. g. (minimum and maximum, 150 and 6,300) on experimental day 42 (Figure 3). Three of five group II sheep had no eggs in feces on experimental day 70; output in feces of the other two sheep averaged 425 e. p. g. on day 154. Haemonchus egg counts in feces of three group III sheep peaked at 22,500 e. p. g. on experimental day 56; mean output was 14,000 e. p. g. (minimum and maximum, 1,650 and 30,000) for four sheep on day 70. Two surviving group III sheep averaged 4,850 e. p. g. on day 154.

Clinicopathologic Findings

Erythrocytes

For sheep in the three control groups, mean weekly values for PCV percentage (Figure 4), erythrocyte counts (Figure 6), and hemoglobin concentrations (Figure 8), are given. For the five principals (group III), mean values for these data for experimental days 0 to 49 are also shown. Individual data for each of five group III sheep are given for PCV (Figure 5), erythrocyte counts (Figure 7), and hemoglobin concentrations (Figure 9).




Figure 3. *Haemonchus contortus* fecal egg counts for each of 5 *Fasciola hepatica* and *H. contortus*-exposed sheep (group III), and mean counts for 5 *H. contortus*-exposed sheep (group II; □--□).



Figure 4. Mean packed cell volumes for 5 *Fasciola hepatica*-exposed sheep (group I; ●-●), 5 *Haemonchus contortus*-exposed sheep (group II; ■-■), 5 *F. hepatica* and *H. contortus*-exposed sheep (group III, X-X), and 5 nonexposed sheep (group IV; o-o).




Figure 5. Packed cell volumes for each of 5 *Fasciola hepatica* and *Haemonchus contortus*-exposed sheep (group III).




Figure 6. Mean erythrocyte counts for 5 *Fasciola hepatica*-exposed sheep (group I; ●—●), 5 *Haemonchus contortus*-exposed sheep (group II, ■—■), 5 *F. hepatica* and *H. contortus*-exposed sheep (group III; X—X), and 5 nonexposed sheep (group IV; ○—○).




Figure 7. Erythrocyte counts for each of 5 *Fasciola hepatica* and *Haemonchus contortus*-exposed sheep (group III).




Figure 8. Mean hemoglobin concentrations for 5 *Fasciola hepatica*-exposed sheep (group I; ●—●), 5 *Haemonchus contortus*-exposed sheep (group II; ■—■), 5 *F. hepatica* and *H. contortus*-exposed sheep (group III; X—X), and 5 nonexposed sheep (group IV; ○—○).




Figure 9. Hemoglobin concentrations for each of 5 Fasciola hepatica and Haemonchus contortus-exposed sheep (group III).

Statistical analyses of these data indicated that mean PCV (25.6%) for group I sheep (F. hepatica-exposed controls) was significantly lower ($P < 0.01$) than that for group IV sheep (nonexposed controls, 34.3%) on experimental day 70 (Figure 4). Significant reduction in PCV continued for the duration of the experiment; PCV was less than 9% for two sheep at time of death on experimental days 162 and 168. For the two sheep that were killed and examined, PCV was 15.5% on experimental day 169 and 38.0% on day 1,063. Mean erythrocyte count and hemoglobin concentration for group I sheep decreased after exposure; significant reduction ($P < 0.01$) in erythrocyte count (7.06×10^6 /cmm.) was found on experimental day 77 (Figure 6), and for hemoglobin concentration (8.42 Gm./100 ml.) on day 70 (Figure 8).

For group II sheep (H. contortus-exposed controls), mean PCV (30.1%) was significantly lower ($P < 0.05$) than that for nonexposed controls (36.5%) on experimental day 63 only (Figure 4). Reduction in erythrocyte count was not significant ($P > 0.05$) on any sample day (Figure 6); mean hemoglobin concentration on days 63 (9.68 Gm./100 ml.) and 77 (10.52 Gm./100 ml.) was significantly lower ($P < 0.05$) than that for nonexposed controls (12.20 and 12.58 Gm./100 ml., respectively) (Figure 8).

Significant reduction ($P < 0.01$) in mean PCV for group III sheep (F. hepatica and H. contortus-exposed principals) began on

experimental day 28 (Figure 4) for erythrocyte count, on day 42 (Figure 6); for hemoglobin concentration, on day 35 (Figure 8). Before death occurred PCV for each group III sheep was 6.5% to 10.0% (Figure 5).

Significant changes in calculated Wintrobe indices are given (Table 1). These data indicate that a progressive macrocytic, normochromic to slight hypochromic anemia occurred in group I sheep. For group II sheep, a transitory normocytic, hypochromic anemia occurred. The following types of anemia were found in group III sheep before time of death: for sheep 13, anemia was macrocytic and hypochromic two days before it died; for sheep 15, normocytic and hypochromic one day before it died; for sheep 11, slightly microcytic and hypochromic eight days before it died; for sheep 12 and 14, macrocytic and normochromic approximately two months before they died (Table 1).

Pathologic changes in circulating erythrocytes for all sheep during the experiment are summarized as follow: Prevalence of anisocytosis, basophilic stippling, diffuse basophilia, Howell-Jolly bodies, and poikilocytosis intensified in three of five group I sheep (F. hepatica-exposed controls) after experimental day 77; for sheep 1 and 5, they appeared on day 126, but were not severe for sheep 5. Macrocytic and hypochromic erythrocytes were observed in blood smears from sheep 3 from experimental day 147 until necropsy on day 269. For group II sheep (H. contortus-exposed controls), these

Table 1. Mean Wintrobe Indices for Sheep in Group I (Fasciola hepatica-exposed Controls), Group II (Haemonchus contortus-exposed Controls), and Group IV (Nonexposed Controls). These Data Before Time of Death are Given for Each Sheep in Group III (F. hepatica and H. contortus-exposed Principals).

Group	Sheep No.	Experimental day	Mean corpuscular volume (cμ)	Mean corpuscular hemoglobin (μμg.)	Mean corpuscular hemoglobin concentration (%)
I		49	35.1	12.0	31.4
II		49	36.4	12.4	34.0
III	13	49	44.2*	13.7	31.0*
IV		49	33.0	11.4	34.2

I		70	35.2	11.6	32.9
II		70	36.6	12.2	33.6
III	15	70	32.8	8.1	24.6*
	11	70	29.2	8.2	28.0*
IV		70	34.7	11.7	33.8

I		91	39.1**	11.8	30.3*
II		91	37.5	11.1	29.7*
IV		91	33.7	11.2	33.1

I		126	46.6*	14.5	31.3
II		126	33.9	10.9	32.2
III	12	126	49.2*	17.5	35.6
	14	126	42.7*	14.4	33.6
IV		126	35.0	12.0	34.1

* Difference from nonexposed control value was highly significant ($P < 0.01$).

** Difference from nonexposed control value was significant ($P < 0.05$).

abnormalities were found in four of five sheep on experimental day 35; maximum intensity occurred between days 63 and 77. They were not evident in blood smears from sheep 9 and 10 after day 77, sheep 6 after day 98, and sheep 7 and 8 after day 161. Pathologic changes were severe only in sheep 8. For group III sheep (F. hepatica and H. contortus-exposed principals) these abnormalities were first seen in sheep 13 on day 35, and in all sheep on day 42. Hypochromic erythrocytes were evident in sheep, 11, 13, and 15 before they died. Absence of basophilic stippling, diffuse basophilia, and Howell-Jolly bodies was noted in blood from sheep 12 and 14 on experimental days 84 and 91. Prevalence of all abnormalities intensified on day 98 and continued until time of death. For group IV sheep, erythrocytes appeared normal in all blood smears examined during the experiment.

Leukocytes

On experimental day 0, mean total leukocyte count for group I sheep was 8,840/cmm., for group II sheep, 9,840/cmm., for group III sheep, 10,480/cmm., and for group IV sheep, 9,160/cmm. Statistical analysis of leukocyte data indicated significant increase ($P < 0.05$) in total leukocyte counts occurred among group I sheep from experimental day 35 to day 70; mean counts exceeded 11,980/cmm. (Figure 10). On experimental days 21 and 28, total leukocyte




Figure 10. Mean leukocyte total and eosinophil counts for 5 *Fasciola hepatica*-exposed sheep (group I; ●—●), 5 *Haemonchus contortus*-exposed sheep (group II; ■—■), 5 *F. hepatica* and *H. contortus*-exposed sheep (group III; X—X), and 5 nonexposed sheep (group IV; ○—○).

counts for sheep 13 (group III) were 14,000 and 14,500/cmm., respectively; on days 42, 49 and 63, counts for sheep 14 exceeded 13,600/cmm. Changes in total leukocyte counts for other sheep in group III and for group II were not significant ($P > 0.05$).

Analyses of leukocyte differential cell count data indicated significant increase ($P < 0.01$) in eosinophil counts occurred in group I sheep (F. hepatica-exposed controls) from experimental day 14 to day 84, and began among group III sheep on day 21. Mean leukocyte total and eosinophil counts for sheep in the three control groups are given (Figure 10). For the five principals (group III), mean eosinophil counts for experimental days 0 to 49 are also shown (Figure 10); data for each of these sheep are given (Figure 11). Peak eosinophilia for sheep in groups I (3,476/cmm.) and III (2,723/cmm.) occurred on experimental day 42. A slight eosinophilic response among group II sheep (H. contortus-exposed controls) peaked on experimental day 21 (884/cmm.) and later, on day 49 (713/cmm.); these counts were not significantly different ($P > 0.05$) from that for nonexposed controls on these days (281 and 191/cmm., respectively) (Figure 10). Decrease in eosinophil counts for sheep 11, 13, and 15 (group III) (Figure 11) was similar to that found among group II sheep (Figure 10). For sheep 12 and 14 (group III) (Figure 11), eosinophil counts paralleled that of group I sheep (Figure 10).




Figure 11. Eosinophil counts for each of 5 Fasciola hepatica and Haemonchus contortus-exposed sheep (Group III).

Among group I sheep, significant increase ($P < 0.05$) in mean neutrophil counts occurred between experimental days 42 and 70; on day 63, maximum count was 3,973/cmm. compared to 2,371/cmm. for nonexposed controls. A significant decrease ($P < 0.05$) in mean neutrophil counts among group I sheep occurred on experimental days 119 and 126 and among group II sheep on day 126. Significant change in mean neutrophil counts for sheep in groups II and III did not occur. Analyses of mean basophil, monocyte, and band counts indicated that significant differences between groups did not occur during the experiment.

Serum albumin

On experimental day 0, mean serum albumin concentration for group I sheep was 3.81 Gm./100 ml., for group II sheep, 3.87 Gm./ml., for group III sheep, 3.78 Gm./100 ml., and for group IV sheep, 3.87 Gm./100 ml. For sheep in the three control groups, mean weekly serum albumin concentrations are given (Figure 12). For the five principals, (group III), mean serum albumin values for experimental days 0 to 49 are also shown (Figure 12); data for each of these sheep are given (Figure 13). Statistical analysis of data indicated the following: For group I sheep (F. hepatica-exposed controls), significant decrease ($P < 0.01$) in albumin content began on experimental day 21 and continued through day 203, maximum reduction




Figure 12. Mean serum albumin concentrations for 5 *Fasciola hepatica*-exposed sheep (group I; ●—●), 5 *Haemonchus contortus*-exposed sheep (group II; ■—■), 5 *F. hepatica* and *H. contortus*-exposed sheep (group III; X—X), and 5 nonexposed sheep (group IV; ○—○).




Figure 13. Serum albumin concentrations for each of 5 *Fasciola hepatica* and *Haemonchus contortus*-exposed sheep (group III).

(1.31 Gm./100 ml.) occurred on day 70 (Figure 12); for group II sheep (H. contortus-exposed controls), it was found in the period from experimental day 21 to 63, maximum reduction (3.08 Gm./100 ml.) occurred on day 56 and concentration was 4.04 Gm./100 ml. on day 112 (Figure 12); for group III sheep (F. hepatica and H. contortus-exposed principals), it began on day 14 and values decreased until time of death (Figures 12, and 13). Serum albumin concentration at time of death for two group I sheep was 1.36 Gm./100 ml. (sheep 2, day 154) and 1.6 Gm./100 ml. (sheep 4, day 161) (Figure 12); for five group III sheep, range in concentration was 1.26 to 1.80 Gm./100 ml. (Figure 13).

Postmortem Findings

Day of necropsy, and final weight and PCV determined before death are given for eight sheep (three in group I; five in group III) that died, and two sheep (group I) that were killed and necropsied (Table 2).

Gross Pathology

Postmortem examination of eight sheep that died revealed intermandibular and subcutaneous edema, and moderate to severe ascites and hydrothorax. Carcasses appeared anemic and blood was thin and watery. Jaundice of skin and mesenteries was noted

Table 2. Final Weight, Packed Cell Volume, Fecal Sample Analysis, and Helminth Recovery Data for each of 5 *Fasciola hepatica*-exposed sheep (group I), and 5 *F. hepatica* and *Haemonchus contortus*-exposed sheep (group III).

Sheep No.	Group	Day of death	Exposure		Helminth burdens							
			No. of <i>F. hepatica</i>	No. of <i>H. contortus</i>	Ova in feces*	<i>Fasciola hepatica</i>			<i>H. contortus</i>		Final weight (kg.)*	Final PCV (%)*
						No. of flukes	Mean length (mm.)	No. measured	Eggs per gram of feces*	No. of worms		
2	I	162	600	none	pos.	412	29.6 (23-35)	50	---	---	35.2	7.5
4	I	168	600	none	pos.	414	31.0 (25-36)	50	---	---	42.3	8.0
1	I	265	600	none	pos.	318	27.0 (22-32)	32	---	---	---	---
3	I	269**	600	none	pos.	383	33.0 (27-41)	38	---	---	50.0	15.5
5	I	1,063**	600	none	pos.	28	28.5 (24-33)	26	---	---	93.6	38.0
13	III	51	600	50,000	neg.	253	6.2 (4-9)	25	27,900	28,000	25.7	10.0
15	III	71	600	50,000	neg.	238	15.4 (9-22)	24	30,000	25,300	43.2	6.5
11	III	78	600	50,000	pos.	332	18.9 (14-25)	33	18,200	18,100	38.6	7.0
12	III	190	600	50,000	pos.	346	30.9 (25-35)	35	15,500	12,600	36.8	7.0
14	III	201	600	50,000	pos.	326	27.8 (22-35)	33	13,800	27,600	57.3	8.0

* The last sample taken before the sheep died.

** The sheep was killed and necropsied. Numbers in parentheses are minimal and maximal values.

for one sheep (13, group III). Perforation of the gallbladder (sheep 12 and 14) or upper duodenum (region of common bile duct, sheep 1) was considered the immediate cause of death for three sheep; numerous mature F. hepatica were recovered in peritoneal fluid from these sheep.

The liver was enlarged in each sheep and gallbladder distension was observed in three sheep (2, 4, and 11). Fluke ova were recovered in bile from eight sheep that were examined after experiment day 71. Numerous subcapsular, hemorrhagic tracts caused by immature F. hepatica were evident on the visceral surface of liver from one sheep (13, experimental day 51). Liver texture changed from soft to firm as the fibrotic reparative process advanced. The "hobnailed appearance" of the visceral surface was observed in sheep (15, 11) that died on experimental days 71 and 78. In sheep that died or were killed after this time, mild fibrosis of liver, especially the left lobe, was evident. Bile ducts on the parietal surface were prominent as duct walls became thickened, distended, and raised. A marked thickening and folding of epithelium of the main bile duct was found in the sheep (5) necropsied on experimental day 1, 063.

Helminthologic Data

Final fecal examination results, number of H. contortus recovered in abomasums from each group III sheep, and number and

mean length of F. hepatica recovered in liver from each sheep in groups I and III are given (Table 2). Fluke ova were recovered in feces of eight sheep that were examined after experimental day 71. Haemonchus contortus e. p. g. counts and worm burdens in group III sheep at time of death were high, even in two sheep (12, 14) that died on experimental days 190 and 201.

The first three sheep (13, 15, 11) that died harbored moderate burdens of F. hepatica (238 to 253) and heavy burdens of H. contortus (18, 100 to 28, 000). Order in time that deaths occurred correlates with the number of H. contortus recovered.

Larger F. hepatica burdens were recovered in two group I sheep (2, 412 flukes; 4, 414 flukes) that died before death of the two remaining group III sheep (12, 346 flukes; 14, 326 flukes). Two group I sheep (1, 3) lived at least two months longer than two group III sheep (12, 14) harboring similar fluke burdens. Assuming that a similar number of F. hepatica became established in sheep 5 (group I), significant reduction in its fluke burden occurred in the 2.2-year period after sheep 3 was killed and examined. Total recovery was 383 F. hepatica in sheep 3 and only 28 in sheep 5.

Histopathology

Examination of sectioned liver from four sheep (13, 15, 11, and 5) revealed the following: Fresh migration tracts in parenchyma

were seen only in one sheep (13) that died on experimental day 51. Section through most recently disrupted tissue indicated cellular debris, lysed erythrocytes, and leukocytic infiltration primarily of neutrophils; some eosinophils and macrophages were evident. In older regions of tracts, eosinophils and macrophages containing hemosiderin characterized the infiltrate; early fibroblastic invasion was observed. In sheep (15, 11) that died on experimental days 71 and 78, migratory tracts contained fibroblasts, collagen, macrophages, and plasma cells. By day 1,063, resolution of tissue damage was complete since no evidence of migration tracts were seen in sections from sheep 5.

On experimental day 51, necrotic foci adjacent to fresh tracts were found; these contained hepatocytes showing degenerative nuclear changes of pyknosis, karyorrhexis, and karyolysis. Mononuclear leukocytic infiltration to portal areas adjacent to migratory tracts was evident, and bile ductules showed early proliferative changes within the portal region. By experimental days 71 and 78, plasma cells, macrophages, and some neutrophils characterized the infiltrate in portal areas; bile ductule proliferation and early stages of portal fibrosis were found. Collagen deposition around proliferating bile ductules caused separation of portal triad components. At this time, hyperplastic changes in bile ductule epithelium were evident. Number and height of epithelial cells increased, and in

larger ducts, invagination of the epithelium occurred. By day 1, 063, portal fibrosis was extensive and marked adenomatous thickening of the main bile duct epithelium, with a dense underlying collagen layer, were prominent features. Erythrocytes were evident on eroded surfaces of this epithelium, and an intense infiltration of eosinophils, plasma cells, and lymphocytes was found in these areas. In hyperplastic bile ductules and adenomatous structure of the main bile duct, epithelial cells in several areas, had eosinophilic intranuclear inclusion bodies of unknown etiology.

Numerous refractile, crystalline structures in parenchyma were evident in liver sections from sheep examined on experimental days 71 and 78. Zonal fatty degeneration around central lobular veins was found in sheep 11 that died on day 78.

DISCUSSION

Clinicopathologic, gross and histopathologic data for five F. hepatica-infected sheep (group I) were characteristic of sheep with chronic fascioliasis. Weight loss, ascites, time of death, and gross liver pathology agree with data reported by Boray (1967, 1969) and Ross, Dow, and Todd (1967). Onset of significant reduction in PCV, erythrocyte count, hemoglobin and serum albumin concentrations, and pathologic erythrocyte condition was as reported by others (Furmaga and Gundlach, 1967a, b; Hjerpe et al., 1971; Ibrović and Gall-Palla, 1959; Sewell, Hammond, and Dinning, 1968; Sinclair, 1962, 1964, 1965; Symons and Boray, 1967, 1968). Macrocytic, normochromic anemia in subacute and chronic fascioliasis was reported by Hjerpe et al. (1971), Reid et al. (1970), Ross (1967a, b). Eosinophilia and leukocytosis observed in the present experiment agree with findings of most workers. Histopathologic changes in liver were similar to descriptions given by others (Dawes, 1963; Dow, Ross, and Todd, 1968; Rubaj and Furmaga, 1969). Presence of eosinophilic intranuclear inclusion bodies in hyperplastic bile duct epithelial cells has not been reported in chronic ovine fascioliasis.

Age resistance to H. contortus infection was demonstrated by sheep in group II. One of five sheep did not acquire a patent

infection, and three of them had no eggs in feces by experimental day 70. Increased erythropoiesis was sufficient to compensate for blood loss, and the normocytic, hypochromic anemia was of short duration. Data similar to these have been reported by others (Baker and Douglas, 1966; Fourie, 1931; Silverman, Mansfield, and Scott, 1970). Eosinophilia and hypoalbuminemia observed in the present experiment agree with data given by Charleston (1965), Kuttler and Marble (1960), Malczewski (1970), and Shumard, Bolin, and Eveleth (1957).

Response of the five F. hepatica and H. contortus-infected sheep indicated several significant changes when compared to that of the three control groups. Death of three sheep before experimental day 80, and of the two remaining sheep two months before other sheep (group I) with similar F. hepatica burdens, indicated enhanced pathogenicity in sheep with concurrent infections. Significant decrease in PCV, erythrocyte count, hemoglobin and serum albumin concentration occurred by experimental day 42, much earlier than was found in group I sheep. Microcytic, hypochromic anemia at time of death of two sheep (15, 11) suggested that the primary causative agent was H. contortus. Macrocytic, normochromic anemia at time of death of two sheep (12, 14) indicated that F. hepatica infection was responsible.

Establishment of both helminths and host response to F.

hepatica appeared normal. Reparative tissue response in sheep with concurrent infections occurred at the time reported for F. hepatica-infected sheep by Dawes (1963) and Dow, Ross, and Todd (1968). Inhibition of development of host resistance to H. contortus by presence of F. hepatica was indicated. Fecal egg counts were higher and the patent period lasted longer in sheep with concurrent infections when compared to that for sheep with H. contortus alone. Examination of data for body weights, erythrocyte counts, hemoglobin and serum albumin concentrations, and prevalence of abnormal erythrocytes in blood smears for two sheep in group III (12, 14) indicated improvement in condition, corresponding to time of recovery to H. contortus infection by sheep in group II. A marked increase in e. p. g. counts for two sheep (12, 14) from experimental day 154 to time of death suggested that superinfection occurred. This indicates that resistance resulting from primary infection did not occur, and is in marked contrast to data given by Silverman, Mansfield, and Scott (1970). Supportive evidence for impaired response in concurrent infections was found in comparison of mean eosinophil counts for F. hepatica-infected sheep and for those with F. hepatica and H. contortus. Since eosinophilia is characteristic of each helminth infection when studied separately, significant increase over that of sheep with F. hepatica alone was anticipated. Results in the present experiment indicate that eosinophilic response in sheep with

simultaneous F. hepatica and H. contortus infections is less than that for sheep with F. hepatica infection alone.

Interference with host response to one nematode species in concurrent infection with two gastrointestinal nematodes was reported by Kates and Turner (1960). They found an extended patent period for N. spathiger in lambs when simultaneous H. contortus infection was experimentally induced.

BIBLIOGRAPHY

- Andrews, J. S. 1942. Stomach worm (Haemonchus contorus) infection in lambs and its relation to gastric hemorrhage and general pathology. *Journal of Agricultural Research* 65:1-18.
- Baker, N. F., E. F. Cook, J. R. Douglas and C. E. Cornelius. 1952. The pathogenesis of trichostrongyloid parasites. III. Some physiological observations in lambs suffering from acute parasitic gastroenteritis. *Journal of Parasitology* 45:643-651.
- Baker, N. F. and J. R. Douglas. 1966. Blood alterations in helminth infections. In: *Biology of parasites, emphasis on veterinary parasites*, ed. by E. J. L. Soulsby. New York, Academic Press. p. 155-183.
- Boray, J. C. 1967. Studies on experimental infections with Fasciola hepatica, with particular reference to acute fascioliasis in sheep. *Annals of Tropical Medicine and Parasitology* 61:439-450.
- _____ 1969. Experimental fascioliasis in Australia. In: *Advances in parasitology*, ed. by Ben Dawes. Vol. 7. New York, Academic Press. p. 95-210.
- Boughton, I. B. and W. T. Hardy. 1935. Stomach worms (Haemonchus contortus) of sheep and goats. College Station. p. 236-239. (Texas Agricultural Experiment Station. Annual Report 48).
- Brambell, M. R., W.A.G. Charleston and P. Tohill. 1964. Abomasal bleeding caused by immature stages of Haemonchus contortus in sheep showing "age resistance." *Journal of Comparative Pathology and Therapeutics* 74:338-345.
- Campbell, J. A. and A. C. Gardiner. 1960. Anaemia in trichostrongylid infestations. *The Veterinary Record* 72: 1006-1011.
- Charleston, W.A.G. 1964. Comparison of blood and bone marrow changes in lambs with haemonchosis and in lambs repeatedly bled. *Journal of Comparative Pathology and Therapeutics* 74:224-240.

- Charleston, W.A.G. 1965. Pathogenesis of experimental haemonchosis in sheep with special reference to the development of resistance. *Journal of Comparative Pathology* 75: 55-67.
- Christie, M. G. 1970. The fate of very large doses of Haemonchus contortus and their effect on conditions in the ovine abomasum. *Journal of Comparative Pathology* 80:89-100.
- Christie, M. G., M. R. Brambell and C. J. Mapes. 1967. Effect of young Haemonchus contortus on abomasal pH in sheep. *The Veterinary Record* 80:207-208.
- Clark, C.H., G. K. Kiesel and C. H. Goby. 1962. Measurements of blood loss caused by Haemonchus contortus infection in sheep. *American Journal of Veterinary Research* 23:977-980.
- Conway, D. P. and J. H. Whitlock. 1965. A study of the variables influencing artificial infections with Haemonchus contortus. *Cornell Veterinarian* 55:19-54.
- Dawes, B. 1963. Hyperplasia of the bile duct in fascioliasis and its relation to the problem of nutrition in the liver fluke. *Parasitology* 53:123-134.
- Dawes, B. and D. L. Hughes. 1964. Fascioliasis: the invasive stages of Fasciola hepatica in mammalian hosts. In: *Advances in parasitology*, ed. by Ben Dawes. Vol. 2. New York, Academic Press. p. 97-168.
- _____ 1970. Fascioliasis: the invasive stages in mammals. In: *Advances in parasitology*, ed. by Ben Dawes. Vol. 8. New York, Academic Press. p. 259-274.
- Dineen, J. K., A. D. Donald, B. M. Wagland and J. Offner. 1965. The dynamics of the host-parasite relationship. III. The response of sheep to primary infection with Haemonchus contortus. *Parasitology* 55:514-525.
- Dow, C., J. G. Ross and J. R. Todd. 1968. The histopathology of Fasciola hepatica infections in sheep. *Parasitology* 58:129-135.
- Durbin, C. G. 1952. Longevity of the liver fluke, Fasciola sp. in sheep. *Proceedings of the Helminthological Society of Washington* 19:120.

- Evans, J. V., M. H. Blunt and W. H. Southcott. 1963. The effects of infection with Haemonchus contortus on the sodium and potassium concentrations in the erythrocytes and plasma, in sheep of different haemoglobin types. *Australian Journal of Agricultural Research* 14:549-558.
- Fourie, P. J. J. 1931. The haematology and pathology of heamonchosis in sheep. 17th Report of the Director of Veterinary Services and Animal Industry, Union of South Africa. p. 495-572.
- Fraday, C. H. and S. E. Knapp. 1967. A radioisotopic assay of acetylcholinesterase in Fasciola hepatica. *Journal of Parasitology* 53:298-302.
- Furmaga, S. and J. L. Gundyach. 1967a. Parasitologic and hematologic investigations in the course of experimental fascioliasis in sheep. *Acta Parasitologica Polonica* 14:313-325.
- _____ 1967b. Studies on the serum proteins in the course of experimental fascioliasis in sheep. *Acta Parasitologica Polonica* 15:93-102.
- Gardiner, J. L. and E. E. Wehr. 1950. Selecting experimental groups of chickens by weight. *Proceedings of the Helminthological Society of Washington* 17:25-26.
- Gordon, H. McL. 1950. Some aspects of parasitic gastroenteritis of sheep. *Australian Veterinary Journal* 26:14-28.
- _____ 1955. Some aspects of fascioliasis. *Australian Veterinary Journal* 31:182-189.
- Hjerpe, C. A., B. C. Tennant, G. L. Crenshaw and N. F. Baker. 1971. Ovine fascioliasis in California. *Journal of the American Veterinary Medical Association* 159:1266-1271.
- Holman, H. H. and I. H. Pattison. 1941. A study of the blood picture of lambs suffering from parasitic gastritis. *The Veterinary Record* 53:491-498.
- Holmes, P. H., J. D. Dargie, J. M. MacLean and W. Mulligan. 1968. Albumin and globulin turnover in chronic ovine fascioliasis. *The Veterinary Record* 83:227-228.

- Holmes, P. H. and J. M. MacLean. 1969. Intestinal loss of iron and its possible reabsorption in chronic ovine fascioliasis. *Research in Veterinary Science* 10:488-489.
- Holmes, P. H., J. M. MacLean, J. D. Dargie, F. W. Jennings and W. Mulligan. 1967. The anaemia in fascioliasis: experiments with ^{51}Cr -labelled red cells. *The Veterinary Record* 80:285-286.
- Ibrović, M. and V. Gall-Palla. 1959. Neka klinička, patološko-anatomoska i biohemiska zapažanja u vezi s akutnom masovnom metiljavošću ovaca. *Veterinaria. Sarajevo* 8:531-537. (Abstracted in *Helminthological Abstracts* 30:109. 1961.)
- Kates, K. C., R. W. Allen and G. I. Wilson. 1962. Effects of two diets on experimental haemonchosis in lambs. *Journal of Parasitology* 48:865-870.
- Kates, K. C., and J. H. Turner. 1960. An experiment on the combined pathogenic effects of Haemonchus contortus and Nematodirus spathiger on lambs. *Proceedings of the Helminthological Society of Washington* 27:62-67.
- Knapp, S. E. 1964a. Response of lambs to normal and irradiated Haemonchus contortus larvae. *Proceedings of the First International Congress of Parasitology. Vol. 1:108-109. Rome, Italy.*
- _____ 1964b. Immune responses by domestic ruminants to irradiated gastrointestinal nematodes. 50 p. (Termination report to United States Atomic Energy Commission. September, 1964.)
- Kuttler, K. L. and D. W. Marble. 1960. Serum protein changes in lambs with naturally acquired nematode infections. *American Journal of Veterinary Research* 21:445-448.
- Lucker, J. T. 1952. Variability in effect of stomach worm, Haemonchus contortus, infections in lambs. *Proceedings of the 89th Meeting of the American Veterinary Medical Association:171-178.*
- Malczewski, A. 1970. Gastro-intestinal helminths of ruminants in Poland. II. Pathogenesis and pathology in experimental Haemonchus contortus infection in lambs. *Acta Parasitologica Polonica* 18:399-415.

- Mapes, C. J. and R. L. Coop. 1970. The interaction of infections of Haemonchus contortus and Nematodirus battus in lambs. I. The effect of massive infections of Haemonchus on subsequent infections of Nematodirus. *Journal of Comparative Pathology* 80:123-136.
- Martin, C. J. and I. Clunies Ross. 1934. A minimal computation of the amount of blood removed daily by Haemonchus contortus. *Journal of Helminthology* 12:137-142.
- Náquira-Vildoso, F. and R. A. Marcial-Rojas. 1971. Fascioliasis. In: *Pathology of protozoal and helminthic diseases with clinical correlations*, ed. by R. A. Marcial-Rojas. Baltimore, Williams and Wilkins Company. p. 477-489.
- Pantelouris, E. M. 1965. The common liver fluke, Fasciola hepatica L. Oxford, Pergamon Press. 259 p.
- Pearson, I. G. 1963. Use of the chromium radioisotope ^{51}Cr to estimate blood loss through ingestion by Fasciola hepatica. *Experimental Parasitology* 13:186-193.
- Pinkiewicz, E. and E. Madej. 1967. Changes in the peripheral blood of Ca, P, K, Na, Mg and AP in the course of experimental fascioliasis in sheep. *Acta Parasitologica Polonica* 15:225-229.
- Poeschel, G. P. and A. C. Todd. 1969. Relationship of the host's diet to the parasitic development of Haemonchus contortus. *American Journal of Veterinary Research* 30:1223-1228.
- Pullan, N. B., M. M. H. Sewell and J. A. Hammond. 1970. Studies on the pathogenicity of massive infections of Fasciola hepatica L. in lambs. *British Veterinary Journal* 126:548-557.
- Reid, J.F.S., J. Armour, F. W. Jennings, K.S. Kirkpatrick and G. M. Urquhart. 1967. The fascioliasis/ostertagiasis complex in young cattle. A guide to diagnosis and therapy. *The Veterinary Record* 80:371-374.
- Reid, J.F.S., J. Armour, G. M. Urquhart and F. W. Jennings. 1970. Studies on ovine fascioliasis. I. Observations on the sequential development of the naturally-acquired disease. *The Veterinary Record* 86:242-249.

- Richard, R. M., R. F. Shumard, A. L. Pope, P. H. Phillips and C. A. Herrick. 1954. The effect of phlebotomy versus stomach worm, Haemonchus contortus, infection on the growth and certain blood constituents of lambs. *Journal of Animal Science* 13:274-282.
- Roberts, H. E. 1968. Observations on experimental acute fascioliasis in sheep. *British Veterinary Journal* 124:433-449.
- Roseby, F. B. 1970. The effect of fasciolosis on the wool production of Merino sheep. *Australian Veterinary Journal* 46:361-365.
- Ross, J. G. 1967a. An epidemiological study of fascioliasis in sheep. *The Veterinary Record* 80:214-217.
- _____ 1967b. A further season of epidemiological studies of Fasciola hepatica infections in sheep. *The Veterinary Record* 80:368-371 .
- Ross, J. G., C. Dow and J. R. Todd. 1967. A study of Fasciola hepatica infections in sheep. *The Veterinary Record* 80:543-546.
- Rubaj, B. and S. Furmaga. 1969. Pathomorphological and histochemical studies of livers of sheep experimentally infected with the liver fluke. *Acta Parasitologica Polonica* 16:77-80.
- Schalm, O. W. 1965. *Veterinary hematology*. 2nd ed. Philadelphia, Lea and Febiger. 664 p.
- Sewell, M.M.H. 1966. The pathogenesis of fascioliasis. *The Veterinary Record* 78:98-109.
- _____ 1967a. Serum enzyme activities in acute ovine fascioliasis. *The Veterinary Record* 80:577-578.
- _____ 1967b. The aetiology of anaemia in chronic fascioliasis: experiments with radioactive tracers. *The Veterinary Record* 80:579-580.
- _____ 1970. Studies on ovine fascioliasis. *The Veterinary Record* 86:634.

- Sewell, M.M.H., J. A. Hammond and D. C. Dinning. 1968. Studies on the aetiology of anaemia in chronic fascioliasis in sheep. *British Veterinary Journal* 124:160-170.
- Shumard, R. F., D. W. Bolin and D. F. Eveleth. 1957. Physiological and nutritional changes in lambs infected with the nematodes, Haemonchus contortus, Trichostrongylus colubriformis, and Nematodirus spathiger. *American Journal of Veterinary Research* 18:330-337.
- Shumard, R. F., R. J. Emerick, W. E. Bemrick, C. A. Herrick, A. L. Pope and P. H. Phillips. 1956. Effects of trace minerals, dicalcium phosphate, phenothiazine and combinations of these on the resistance of lambs to Haemonchus contortus and other nematodes. *American Journal of Veterinary Research* 17:252-255.
- Silverman, P. H., M. E. Mansfield and H. L. Scott. 1970. Haemonchus contortus infection in sheep: effects of various levels of primary infections on nontreated lambs. *American Journal of Veterinary Research* 31:841-857.
- Sinclair, K. B. 1960. Serum calcium and magnesium levels in sheep infected with Fasciola hepatica. *The Veterinary Record* 72:506.
- _____ 1962. Observations on the clinical pathology of ovine fascioliasis. *British Veterinary Journal* 118:37-53.
- _____ 1964. Studies on the anaemia of ovine fascioliasis. *British Veterinary Journal* 120:212-222.
- _____ 1965. Iron metabolism in ovine fascioliasis. *British Veterinary Journal* 121:451-461.
- _____ 1967a. Pathogenesis of Fasciola and other liver flukes. *Helminthological Abstracts* 36:115-134.
- _____ 1967b. The effect of corticosteroid on the plasma protein of lambs infected with Fasciola hepatica. *Research in Veterinary Science* 9:181-183.

- Sinclair, K. B. 1968. The effect of corticosteroid on the pathogenicity and development of Fasciola hepatica in lambs. *British Veterinary Journal* 124:133-139.
- _____ 1969. Some aspects of the pathogenesis and treatment of fascioliasis. *The Veterinary Record* 84:544-547.
- _____ 1970a. The effect of splenectomy on the pathogenicity of Fasciola hepatica in the sheep. *British Veterinary Journal* 126:15-29.
- _____ 1970b. The pathogenicity of Fasciola hepatica in previously infected, corticosteroid-treated lambs. *Research in Veterinary Science* 11:209-216.
- Soulsby, E. J. L. 1965. Textbook of veterinary clinical parasitology. Vol. I. Helminths. Pennsylvania, F. A. Davis Company. 1120 p.
- Stephenson, W. 1947. Physiological and histochemical observations on the adult liver fluke, Fasciola hepatica L. II. Feeding. *Parasitology* 38:123-127.
- Stoll, N. R. 1943. The wandering of Haemonchus in the sheep host. *Journal of Parasitology* 29:407-418.
- Symons, L. E. A. and J. C. Boray. 1967. Anaemia of ovine fascioliasis. *The Veterinary Record* 8:130.
- _____ 1968. The anaemia of acute and chronic ovine fascioliasis. *Zeitschrift für Tropenmedizin und Parasitologie* 19:451-472.
- Taylor, E. L. 1964. Fascioliasis and the liver fluke. Rome. 234 p. (Food and Agriculture Organization of the United States. F.A.O. Agricultural Studies no. 64)
- Todd, J. R. and J. G. Ross. 1966. Origin of hemoglobin in the cecal contents of Fasciola hepatica. *Experimental Parasitology* 19:151-154.
- _____ 1968. Biochemical and haematological changes in the blood of normal sheep following repeated phlebotomy. *British Veterinary Journal* 124:352-362.

- Turner, J. H. and M. L. Colglazier. 1954. Control of pasture-acquired infections of Nematodirus \$pathiger and Haemonchus contortus in lambs with phenothiazine-salt mixture. *American Journal of Veterinary Research* 15:564-573.
- Turner, J. H., K. C. Kates and G. I. Wilson. 1962. The interaction of concurrent infections of the abomasal nematodes, Haemonchus contortus, Ostertagia circumcincta, and Trichostrongylus axei (Trichostrongylidae), in lambs. *Proceedings of the Helminthological Society of Washington* 29:210-216.
- Veglia, F. 1915. The anatomy and life history of Haemonchus contortus (Rud.), 3rd. and 4th. Reports of the Director of Veterinary Research, Dept. of Agriculture, Union of South Africa. p. 347-500.
- Weir, W. C., T. L. Bahler, A. L. Pope, P. H. Phillips, C. A. Herrick and G. Bohstedt. 1948. The effect of hemopoietic dietary factors on the resistance of lambs to parasitism with the stomach worm, Haemonchus contortus. *Journal of Animal Science* 7:566-574.
- Whitlock, J. H. 1950. The anemias in the trichostrongylidoses. *Cornell Veterinarian* 40:288-299.
- Wilson, G. I. and J. H. Turner. 1965. Serum protein studies on sheep and goats: studies on Targhee lambs raised under varied exposures to parasitism. *American Journal of Veterinary Research* 26:645-650.