



Open Access Articles

Introduction of the exotic tick *Hyalomma truncatum* on a human with travel to Ethiopia: A case report

The Faculty of Oregon State University has made this article openly available.
Please share how this access benefits you. Your story matters.

Citation	Mathison, B. A., Gerth, W. J., Pritt, B. S., & Baugh, S. (2015). Introduction of the exotic tick <i>Hyalomma truncatum</i> on a human with travel to Ethiopia: A case report. <i>Ticks and Tick-borne Diseases</i> , 6(2), 152-154. doi:10.1016/j.ttbdis.2014.11.005
DOI	10.1016/j.ttbdis.2014.11.005
Publisher	Elsevier
Version	Version of Record
Terms of Use	http://cdss.library.oregonstate.edu/sa-termsofuse



Mini Review

Introduction of the exotic tick *Hyalomma truncatum* on a human with travel to Ethiopia: A case reportBlaine A. Mathison^{a,*}, William J. Gerth^b, Bobbi S. Pritt^c, Stephen Baugh^d^a Division of Parasitic Diseases and Malaria, Centers for Disease Control and Prevention, Atlanta, GA, United States^b Oregon State University Plant Clinic, Corvallis, OR, United States^c Mayo Clinic, Rochester, MN, United States^d Legacy Medical Group Northwest, Portland, OR, United States

ARTICLE INFO

Article history:

Received 1 August 2014

Received in revised form

10 November 2014

Accepted 11 November 2014

Available online 20 November 2014

ABSTRACT

An Oregon resident returned from a photography trip to Ethiopia with a male *Hyalomma truncatum* tick attached to the skin on his lower back. The tick was identified morphologically and deposited in the U.S. National Tick Collection housed at Georgia Southern University, Statesboro, Georgia. The public health importance of *Hyalomma* species of ticks and diagnostic dilemmas with identifying exotic ticks imported into the U.S. are discussed.

Published by Elsevier GmbH.

Keywords:

Hyalomma

Ticks

Exotic

Vector-borne

Case report

A 59-year-old previously healthy male presented to his primary care clinic on December 9, 2013, after discovering a tick on his lower back the preceding day. He had initially mistaken the tick for a skin tag and bluntly removed it. After identifying it as a possible tick, he placed it in a plastic bag with moistened napkin and placed it in the refrigerator. The patient had returned from a photography trip to Ethiopia where he had visited several rural towns and villages approximately 5 days prior to presentation. After returning to the U.S., he had spent time in California but did not hike nor camp. The patient had reported a self-limiting headache the previous day and chills during the flight home. He was otherwise asymptomatic and had not experienced arthralgias or myalgias. On examination, the patient was afebrile and appeared well. The area surrounding the attachment site was raised, firm, and tender, surrounded by a reddened, concentric rash that was three inches in diameter [Fig. 1]. The case was discussed with the local infectious disease specialist via telephone. Given the patient's mild symptoms, serologies were not recommended unless he developed worsening symptoms. However, because of

the risk of acquiring rickettsial diseases with this species of tick, the patient was started on 100 mg doxycycline twice daily for a total of 21 days.

The tick was sent to the Oregon State University Plant Clinic for identification, where it was determined to be a genus or species of hard tick not commonly occurring in the Pacific Northwest. Digital images of the tick were shared with the first author, who made the tentative identification of an adult stage *Hyalomma*. Diagnostic features supporting the genus-level identification included the presence of eyes, irregular festoons, elongate mouthparts, an inornate dorsal shield, and a characteristic banding pattern seen on the legs of many *Hyalomma* species [Fig. 2], as well as having the spurs on the forecoxae subequal in length [Fig. 3] (Mathison and Pritt, 2014). Digital images were shared with Dr. Dmitry Apanaskevich at the U.S. National Tick Collection (USNTC) at Georgia Southern University who suggested an identification of *H. truncatum*. The specimen was forwarded to the Centers for Disease Control and Prevention, Division of Parasitic Diseases and Malaria, where the identification of *H. truncatum* was confirmed based on the following morphologic criteria: a dorsal shield irregularly punctate, denser posteriorly; lateral grooves [Fig. 2] deep and reaching the eyes; subanal plates in-line with the adanal plates [Fig. 3] (Apanaskevich and Horak, 2008c). The specimen (a male) was deposited in the USNTC for archiving.

* Corresponding author. Tel.: +1 404 718 4103.

E-mail address: gqa4@cdc.gov (B.A. Mathison).



Fig. 1. Skin lesion observed at the bite site.

Discussion

Hyalomma is one of the most medically important tick genera in the Old World. Species in this genus have been reported to transmit a variety of viral, bacterial, and parasitic diseases of medical and veterinary importance (Bakheit et al., 2012). One of the most important human viral agents transmitted by *Hyalomma* spp. is Crimean-Congo hemorrhagic fever (CCHF) virus. Sexually and transovarially transmission of CCHF virus was observed experimentally in *Hyalomma truncatum* (Gonzalez et al., 1992). In addition, extensive documentation exist showing that several tick species transmit this virus in different geographical areas of the world: *H. marginatum* in southern Russia, Turkey, and the Balkan and Crimean Peninsulas, *H. anatolicum* in Iran, Pakistan, Turkmenistan, and Tajikistan, *H. asiaticum* from central Asia to China, and *H. rufipes* in Africa (Hoogstraal, 1979; Bakheit et al., 2012; Goddard, 2012).

Other important disease agents of humans and their documented vectors include *Rickettsia conorii* (*H. rufipes* and *H.*



Fig. 2. Dorsal image of the *Hyalomma* specimen (male) (EY, eye; LG, lateral groove; FS, festoons).




Fig. 3. Ventral image of the *Hyalomma* specimen (male) (FC, spurs on the forecoxa; AN, adanal plate; SA, subanal plate).

truncatum), *R. aeschlimannii* (*H. marginatum*, *H. truncatum*, and *H. scupense*), *R. sibirica* (*H. asiaticum*, *H. excavatum*, and *H. truncatum*), *Anaplasma phagocytophilum* (*H. lusitanicum*), and *Coxiella burnetii* (*H. aegyptium* and *H. scupense*) (Beati et al., 1997; Bakheit et al., 2012; Goddard, 2012), and Dugbe virus (Hoogstraal et al., 1981). Kumsa et al. (2014) detected *R. aeschlimannii* in *H. marginatum* and *H. truncatum* in Ethiopia. Paštiu et al. (2012) found *H. aegyptium* naturally infected with *Ehrlichia canis* and *A. phagocytophilum* in Romania. *Hyalomma truncatum* has been suggested as a possible vector of Rift Valley fever virus (Linthicum et al., 1989; Nchu and Rand, 2013). *Hyalomma* species have also been implicated in tick paralysis in humans (Edussuriya and Weilgama, 2003; Gürbüz et al., 2010; Doğan et al., 2012). In general, prophylaxis is not recommended for prevention of rickettsial or viral tickborne infections (Wormser et al., 2006; Bakken et al., 2006). However, a course of ribavirin may be considered for prophylaxis of CCHF in individuals at high risk of severe disease (Appannanavar and Mishra, 2011).

There are scattered reports in the literature of *Hyalomma* species being imported into the United States, most-commonly on animals and animal products. Mertins and Schlater (1991) documented five species of *Hyalomma* on ostriches imported from Africa and Europe. Burrige and Simmons (2003) documented *H. aegyptium* imported on Greek tortoises to New York, Florida, and North Carolina. Becklund (1968) and Keirans and Durden (2001) presented comprehensive overviews of imported ticks into the U.S. Most of the records for *Hyalomma* spp. are from animals or products made from animals (e.g. trophy hides), but Keirans and Durden (2001) did report one case of *H. marginatum* being found on a human with travel history to Greece. In the USNTC, there is a specimen of *H. truncatum* collected from a human in Illinois following travel to Botswana (Dmitry Apanaskevich, pers. comm. 2013). *Hyalomma truncatum* has a wide host range; Apanaskevich and Horak (2008c)

documented seven mammalian hosts for the immature stages and 18 mammalian hosts (including humans) for adults in South Africa. As such, the risk for a person to acquire this tick in an endemic area would be greater than for those species whose normal hosts are tortoises and fossorial or other wild animals that have less human contact.

Because *Hyalomma* species are not native to North America, medical practitioners and diagnostic microbiologists and parasitologists may not consider them in the differential diagnosis when identifying ticks removed from patients treated in the U.S. Given their morphologic features (elongate mouthparts, eyes present, festoons present), *Hyalomma* spp. might be misidentified as *Amblyomma* in most keys to North American ticks. *Hyalomma* can best be separated from *Amblyomma* by having festoons of varying size, an inornate dorsal shield (scutum), and spurs on the coxae I roughly equal in length (Mathison and Pritt, 2014). Many *Hyalomma* species also have a characteristic banding pattern on their legs whereby the legs are dark with white maculae at the joints (Fig. 2). Regional keys for *Hyalomma* are available for the United Kingdom (Arthur, 1963), Russia [former U.S.S.R.] (Pomerantzev, 1950), Sudan (Hoogstraal, 1956), Uganda (Matthysse and Colbo, 1987), Saudi Arabia (Hoogstraal et al., 1981), Pakistan and the Indian subcontinent (Kaiser and Hoogstraal, 1964), and Iran (Hosseini-Chegeni et al., 2013). Recent systematic treatments are available for the subgenera *Euhyalomma* (Apanaskevich and Horak, 2005a,b, 2007, 2008a,b,c,d, 2009, 2010a,b) and *Hyalomma* (Apanaskevich et al., 2009). As with other ectoparasites, detailed travel history can be important for obtaining a definitive diagnosis of ticks, as well as to help assess the risk of vector-borne diseases.

Acknowledgements

We would like to thank Dmitry Apanaskevich (USNTC) for assisting with the identification. Rosmarie Kelly (Georgia Department of Health, Atlanta, GA) and William Nicholson (Rickettsial Zoonoses Branch, Division of Vector-Borne Diseases, CDC, Atlanta, GA) kindly reviewed the manuscript.

References

- Apanaskevich, D.A., Horak, I.G., 2005a. The genus *Hyalomma* Koch, 1844. I. Reinstatement of *Hyalomma* (*Euhyalomma*) *glabrum* Delpy, 1949 (Acari: Ixodidae) as a valid species with a redescription of the adults, the first description of its immature stages and notes on its biology. *Onderstepoort. J. Vet. Res.* 73, 1–12.
- Apanaskevich, D.A., Horak, I.G., 2005b. The genus *Hyalomma* Koch, 1844. II. Taxonomic status of *H. (Euhyalomma) anatolicum* Koch, 1844 and *H. (E.) excavatum* Koch, 1844 (Acari: Ixodidae) with redescription of all stages. *Acarina* 13, 181–197.
- Apanaskevich, D.A., Horak, I.G., 2007. The genus *Hyalomma* Koch, 1844. III. Redescription of the adults and larvae of *H. (Euhyalomma) impressum* Koch, 1844 (Acari: Ixodidae) with a first description of its nymph and notes on its biology. *Folia Parasitol.* 54, 51–58.
- Apanaskevich, D.A., Horak, I.G., 2008a. The genus *Hyalomma* Koch, 1844. IV. Redescription of all parasitic stages of *H. (Euhyalomma) lusitanicum* Koch, 1844 and the adults of *H. (E.) franchinii* Tonelli Rondelli, 1932 (Acari: Ixodidae) with a first description of its immature stages. *Folia Parasitol.* 55, 61–74.
- Apanaskevich, D.A., Horak, I.G., 2008b. The genus *Hyalomma* Koch, 1844. V. Re-evaluation of the taxonomic rank of taxa comprising the *H. (Euhyalomma) marginatum* complex of species (Acari: Ixodidae) with redescription of parasitic stages and notes on biology. *Int. J. Acarol.* 34, 13–42.
- Apanaskevich, D.A., Horak, I.G., 2008c. The genus *Hyalomma* Koch, 1844. VI. Systematics of *H. (Euhyalomma) truncatum* and the closely related species, *H. (E.) albiparvum* and *H. (E.) nitidum* (Acari: Ixodidae). *Exp. Appl. Acarol.* 44, 115–136.
- Apanaskevich, D.A., Horak, I.G., 2008d. The genus *Hyalomma* Koch, 1844. VII. Redescription of all parasitic stages of *H. (E.) dromedarii* and *H. (E.) schulzei* (Acari: Ixodidae). *J. Med. Entomol.* 45, 817–831.
- Apanaskevich, D.A., Horak, I.G., 2009. The genus *Hyalomma* Koch, 1844–2009. IX. Redescription of all parasitic stages of *H. (E.) impeltatum* Schulze & Schlottke, 1930 and *H. (E.) somalicum* Tonelli Rondelli, 1935 (Acari: Ixodidae). *Syst. Parasitol.* 73, 199–218.
- Apanaskevich, D.A., Horak, I.G., 2010a. The genus *Hyalomma* Koch, 1844. X. Redescription of the parasitic stages of *H. (Euhyalomma) scupese* Schulze, 1919 (*H. detritum* Schulze) (Acari: Ixodidae) and notes on its biology. *Folia Parasitol.* 57, 69–78.
- Apanaskevich, D.A., Horak, I.G., 2010b. The genus *Hyalomma* Koch, 1844. XI. Redescription of all stages of *H. (Euhyalomma) asiaticum* (Acari: Ixodidae) and notes on its biology. *Exp. Appl. Acarol.* 52, 207–220.
- Apanaskevich, D.A., Horak, I.G., Geevarghese, G., 2009. The genus *Hyalomma* Koch, 1844. VIII. Redescription of three *Hyalomma* Schulze, 1919 species (Acari: Ixodidae) from South Asia with notes on their biology. *Zootaxa* 2050, 31–55.
- Appannanavar, S.B., Mishra, B., 2011. An update on Crimean-Congo hemorrhagic fever. *J. Global Infect. Dis.* 3, 285–292.
- Arthur, D.R., 1963. *British Ticks*. Butterworths, London, United Kingdom.
- Bakheit, M.A., Latif, A.A., Vatansever, Z., Seitzer, U., Ahmed, J., 2012. The huge risks due to *Hyalomma* ticks. In: Mehlhorn, H. (Ed.), *Arthropods as Vectors of Emerging Diseases*. Springer-Verlag, Berlin (Chapter 8).
- Bakken, J.S., Folk, S.M., Paddock, Ch.D., Bloch, K.C., Krusell, A., Sexton, D.J., Buckingham, S.C., Marshall, G.S., Storch, G.A., Dasch, G.A., McQuiston, J.H., Swerdlow, D.L., Dumler, J.S., Nicholson, W.L., Walker, D.H., Eremeeva, M.E., Ohl, C.A., 2006. Diagnosis and management of tickborne rickettsial diseases: rocky mountain spotted fever, ehrlichiosis, and anaplasmosis – United States. *Morb. Mortal. Wkly. Rep.* 55, 1–27.
- Beati, L., Meskini, M., Thiers, B., Raoult, D., 1997. *Rickettsia aeschlimannii* sp. nov., a new spotted fever group of rickettsia associated with *Hyalomma marginatum* ticks. *Int. J. Syst. Bacteriol.* 47, 548–554.
- Becklund, W.W., 1968. Ticks of veterinary significance found on imports in the United States. *J. Parasitol.* 54, 622–628.
- Burridge, M.J., Simmons, L.A., 2003. Exotic ticks introduced into the United States on imported reptiles from 1962 to 2001 and their potential roles in the international dissemination of diseases. *Vet. Parasitol.* 113, 289–320.
- Doğan, M., Devge, C., Tanrıöver Pata, Y.S., Sönmezoglu, M., 2012. Case report: facial nerve paralysis due to intra-aural *Hyalomma* tick infestation. *Turkiye Parazitol. Derg.* 36, 254–257.
- Edussuriya, B.D., Weilgama, D.J., 2003. Case reports: intra-aural tick infestations in humans in Sri Lanka. *Trans. R. Soc. Trop. Med. Hyg.* 97, 412–413.
- Goddard, J., 2012. *Physician's Guide to Arthropods of Medical Importance*, 6th ed. CRC Press, Boca Raton, LA.
- Gonzalez, J.P., Camicas, J.L., Cornet, J.P., Faye, O., Wilson, M.L., 1992. Sexual and transovarian transmission of Crimean-Congo haemorrhagic fever virus in *Hyalomma truncatum* ticks. *Res. Virol.* 143, 23–28.
- Gürbüz, M.K., Erdoğan, M., Doğan, N., Birdane, L., Cingi, C., Cingi, E., 2010. Case report: isolated facial paralysis with a tick. *Turkiye Parazitol. Derg.* 34, 61–64.
- Hoogstraal, H., 1956. African Ixodoidea. I. Ticks of the Sudan (with special reference to Equatoria province and with preliminary reviews of the genera *Boophilus*, *Magaropus*, and *Hyalomma*). U.S. Naval Medical Research Report NM 005 050.29.07.
- Hoogstraal, H., 1979. The epidemiology of tick-borne Crimean-Congo hemorrhagic fever in Asia, Europe, and Africa. *J. Med. Entomol.* 15, 307–417.
- Hoogstraal, H., Wassef, H.Y., Büttiker, W., 1981. Ticks (Acarina) of Saudi Arabia. *Fam. Argasidae, Ixodidae*. Fauna of Saudi Arabia, vol. 3, pp. 25–110.
- Hosseini-Chegeni, A., Hosseini, R., Tavakoli, M., Telmadarrayi, Z., Abdigoudarzi, M., 2013. The Iranian *Hyalomma* (Acari: Ixodidae) with a key to the identification of male species. *Persian J. Acarol.* 2, 503–529.
- Kaiser, M.N., Hoogstraal, H., 1964. The *Hyalomma* ticks (Ixodoidea, Ixodidae) of Pakistan, India, and Ceylon, with keys to subgenera and species. *Acarologia* 6, 257–286.
- Keirans, J.E., Durden, L.A., 2001. Invasion: exotic ticks (Acari: Argasidae, Ixodidae) imported into the United States. A review and new records. *J. Med. Entomol.* 38, 850–861.
- Kumsa, B., Socolovschi, C., Raoult, D., Parola, P., 2014. Spotted fever group rickettsiae in ixodid ticks in Oromia, Ethiopia. *Ticks Tick Borne Dis.*, <http://dx.doi.org/10.1016/j.ttbdis.2014.08.001> (ePub 2014 Sep 25), pii:S1877-959X(14)00170-8.
- Linthicum, K.J., Logan, T.M., Bailey, C.L., Dohm, D.J., Moulton, J.R., 1989. Transstadial and horizontal transmission of Rift Valley fever virus in *Hyalomma truncatum*. *Am. J. Trop. Med. Hyg.* 41, 491–496.
- Mathison, B.A., Pritt, B.S., 2014. Laboratory identification of arthropod ectoparasites. *Concepts Magn. Reson.* 27, 48–67.
- Matthysse, J.G., Colbo, M.H., 1987. Ixodid Ticks of Uganda. Entomological Society of America, Lanham, MD.
- Mertins, J.W., Schlatter, J.L., 1991. Exotic ectoparasites of ostriches recently imported into the United States. *J. Wildl. Dis.* 27, 180–182.
- Nchu, F., Rand, A., 2013. Rift Valley fever outbreaks: possible implication of *Hyalomma truncatum* (Acari: Ixodidae). *Afr. J. Microbiol. Res.* 7, 3891–3894.
- Paştıu, A.L., Matei, I.A., Mihalca, A.D., D'Amico, G., Dumitrache, M.O., Kalmár, Z., Sándor, A., Lefkaditis, M., Gherman, C.M., Cozma, V., 2012. Zoonotic pathogens associated with *Hyalomma aegyptium* in endangered tortoise: evidence for host-switching behavior in ticks? *Parasites Vectors* 5, 301–306.
- Pomerantzev, B.I., 1950. Ixodid ticks (Ixodidae). Fauna of the USSR. Arachnida, vol. IV. Zoological Institute of the Academy of Sciences USSR, Moscow, Russia.
- Wormser, G.P., Dattwyler, R.J., Shapiro, E.D., Halperin, J.J., Steere, A.C., Klempner, M.S., Krause, P.J., Bakken, J.S., Strle, F., Stanek, G., Bockenstedt, L., Fish, D., Dumler, J.S., Nadelman, R.B., 2006. The clinical assessment, treatment, and prevention of Lyme disease, human granulocytic anaplasmosis, and babesiosis: clinical practice guidelines by the Infectious Diseases Society of America. *Clin. Infect. Dis.* 43, 1089–1134.