

# Hazelnut Cultivar

'Lewis' (OSU243.002)

A.N. Azarenko, S. Mehlenbacher, D. Smith, and R. McCluskey

'Lewis' is a hazelnut cultivar developed by Oregon State University and released in January 1997. This cultivar is precocious, and it is more productive and has a smaller tree size than 'Barcelona.' Harvest is earlier than for 'Barcelona,' and the nuts have fewer defects.

#### **Horticultural characteristics**

Tree growth habit. 'Lewis' is a "grower-friendly" tree, with a good balance between vegetative and reproductive growth (Figure 1). It can be managed much like a 'Barcelona' tree. Tree size is approximately 75 percent of 'Barcelona,' as measured by the trunk cross-sectional area (Figure 4).

Flowering characteristics. Female flowers bloom at the same time as 'Barcelona.' The incompatibility alleles are  $S_3S_8$ . The recommended pollinizers are 'Tonda di Giffoni' and 'Hall's Giant.'

Yield and yield efficiency. This selection has out-yielded 'Barcelona' in the first 4 bearing years, and yields were similar in 1997 (Figure 5). Also, 'Lewis' is a more efficient nut producer than 'Barcelona' because yields are greater and tree size is smaller (Figure 6). Therefore, more of the tree's resources are used to produce nuts, and less are used to develop leaves and wood.

Harvest time. Nut clusters contain 3–4 nuts (Figure 2). Nuts are free-husking and ready to harvest 5–7 days earlier than 'Barcelona.'

Nut and kernel quality. 'Lewis' nuts (Figure 3) are smaller than 'Barcelona.' Only 17 percent of the nuts had defects in 1997, compared to 28 percent for 'Barcelona.' Most of the defects of 'Lewis' are shriveled, poorly filled, or moldy nuts. In 1997, only 1 percent of the nuts were blanks, in contrast to 8 percent for 'Barcelona.' The percent kernel is 4–5 percent greater than 'Barcelona.' Raw kernels have very little fiber. The kernels blanch slightly better than 'Barcelona' and have good flavor and texture.

*Propagation.* 'Lewis' has been easy to propagate by the "tie-off' method. Layers are well rooted and vigorous.

Pest tolerance. One of the major strengths of this cultivar is its tolerance to eastern filbert blight. Its tolerance to the pathogen is better than 'Willamette' and 'Hall's Giant.' Big bud mite tolerance is comparable to 'Casina.'

#### Development

'Lewis' was selected from progeny of a cross of OSU17.028 ('Barcelona' x 'Tombul Ghiaghli') and 'Willamette.' The cross was made in 1981 by Dr. Maxine Thompson at the Smith Vegetable and Hazelnut Research Farm. Preliminary analysis of the performance of the genotype began in the seedling row, and in 1988 it was identified as a promising selection. An advanced selection trial was established in 1991 at the Smith Farm. The data presented in this report were obtained from that trial.



Figure 1.—Tree growth habit.



Figure 2.—Nut cluster.

Anita Nina Azarenko, professor of horticulture; Shawn Mehlenbacher, professor of horticulture; David Smith, senior faculty research assistant in horticulture; and Rebecca McCluskey, former research assistant in horticulture; Oregon State University.





Figure 3.—Nuts, raw kernels, and blanched kernels.

## 'Lewis'

### Flowering characteristics

Incompatibility alleles: S<sub>3</sub>S<sub>8</sub>

Time of female flower bloom: similar to 'Barcelona'

#### **Pollinizer recommendations**

'Hall's Giant'  $(S_5S_{15})$ —half of the pollinizer trees 'Tonda di Giffoni'  $(S_2S_{23})$ —half of the pollinizer trees

#### **Estimated time of harvest**

5-7 days earlier than 'Barcelona'

#### Nut and kernel quality (1996 and 1997)

	'Barcelona'		'Lewis'	
	1996	1997	1996	1997
Nut weight (g)	3.8	3.5	2.9	2.7
Kernel weight (g)	1.7	1.4	1.4	1.2
Kernel percentage	43	41	47	46
Blanching rating	4.1	3.6	3.3	2.8
(1-7; 1=100%  removal)				
of pellicle)	0.5	72	00	02
Nuts free of defects (%)	85	72	90	83

#### **Pest tolerance**

Eastern filbert blight: more tolerant than 'Willamette' and 'Hall's Giant'

Big bud mite: comparable to 'Casina'

#### **Released January 1997**




Figure 4.—Trunk cross-sectional area.




Figure 5.—Nut yield.





Figure 6.—Yield efficiency.



© 2002 Oregon State University. This publication may be photocopied or reprinted in its entirety for noncommercial purposes.

This publication was produced and distributed in furtherance of the Acts of Congress of May 8 and June 30, 1914. Extension work is a cooperative program of Oregon State University, the U.S. Department of Agriculture, and Oregon counties. Oregon State University Extension Service offers educational programs, activities, and materials—without regard to race, color, religion, sex, sexual orientation, national origin, age, marital status, disability, and disabled veteran or Vietnam-era veteran status—as required by Title VI of the Civil Rights Act of 1964, Title IX of the Education Amendments of 1972, and Section 504 of the Rehabilitation Act of 1973. Oregon State University Extension Service is an Equal Opportunity Employer.