

077

ASS

Analytica

Canada Thistle

Scientific name: *Cirsium arvense*

By Gordon Cook, *Sherman County Extension Agent*
 Martin Zimmerman, *Gilliam County Extension Agent*
 Ronald Burr, *Extension Agronomist, Oregon State University*



Perennial

Flowers purple to white


Mature plant 1 to 4 feet

Leaves usually long and narrow with short, sharp spines

Plants spreading by creeping underground roots

Canada thistle is most commonly found on range lands and in non-cultivated areas. However, it does occur in grain fields.

Control: Best control obtained at flower bud stage of growth. Spot treatment on non-cropland 1 to 1½ pounds picloram (Tordon) per acre. Spot treatment on cropland 4 to 6 pounds dicamba (Banvel). 10 pounds 2,4-D Amine at bud stage. 2 pounds 2,4-D or 2 pounds Amitrol (non-cropland only) give limited control.



OREGON STATE UNIVERSITY
EXTENSION SERVICE

Extension Service, Oregon State University, Corvallis, Joseph R. Cox, director. This publication was produced and distributed in furtherance of the Acts of Congress of May 8 and June 30, 1914. Extension work is a cooperative program of Oregon State University, the U. S. Department of Agriculture, and Oregon counties.