

# Apple

## 2014 Pest Management Guide for the Willamette Valley

EM 8418 • Revised March 2014

Jay W. Pscheidt, Ed Peachey, and Steve Castagnoli

The chemicals, formulations, and rates listed for insect, mite, and disease control are among the best recommendations based on label directions, research, and orchard use experience. Only a thorough knowledge of the orchard, its variety, tree size and density, canopy characteristics, pest complex, and past pest problems will enable you to correctly select chemicals, rates, amount of water used per acre, and method of application for optimum pest control. Occasionally, different formulations of a product or like formulations containing a different amount of active ingredient also are registered and effective for use on the pests listed. These products also may be used; we do not intend to discriminate against them. You may wish to consult their labels and determine whether their use confers advantages over the products listed in this guide.

Always refer to the pesticide label for use instructions. It is the legal document regarding use patterns. Two questions frequently are asked about the chemical control of insects and diseases: “How much chemical do I use per acre?” and “What is the least amount of water I need per acre to apply in my concentrate sprayer?” Notice that the schedule below suggests an amount of formulated product (not active ingredient) to use per acre. This amount is based on a “typical” middle age and density orchard with moderate pest pressure. Common sense indicates that less material may be needed (than that given) for 1- to 4-year-old orchards. Conversely, more chemical (within label limits) may be required for large, mature trees experiencing heavy pest pressure from multiple pests.

Many insecticide labels today indicate the minimum amount of water needed per acre to apply concentrate sprays of insecticides, as well as how to calculate the amount of chemical needed per acre in a concentrate sprayer. **CHECK LABEL BEFORE SPRAYING!** Some label directions indicate dilute

applications only, such as the dimethoate labels for cherry fruit fly control.

Also:

1. Make sure any tank-mixes of pesticides are compatible. For example, the elevated pH of some boron spray solutions weakens many insecticides. Boron also is incompatible with water-soluble packets.
2. Use adjuvants and spreader stickers with caution.

### Important information

1. Be aware of worker protection standards (WPS). All new pesticide labels will provide orchard reentry intervals and personal protection equipment information.
2. Diazinon is now classified as a restricted use pesticide due to bird toxicity. Maximum per-acre application rates have been reduced to 4 lb 50W, and the preharvest interval extended to 21 days.
3. Endosulfan: Preharvest intervals have changed. Maximum per-acre application rates are reduced.
4. *Orchard Pest Management, a Resource Book for the Pacific Northwest*, 1993 (edited by Beers, Brunner, Willet, and Warner, published by the Good Fruit Grower, Yakima, WA) provides a comprehensive list of the tree fruit insect and mite pests of orchards. Life histories, damage, detection, monitoring, and management of the pests are covered. It is one of our primary sources of information in developing this pest management guide and the most complete reference on orchard use of the principles of integrated pest management.

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*


## Stages

Delayed Dormant (Stages 1–2)  
 Prepink or Green Bud (Stages 3–4)  
 Pink or Preblossom (Stages 5–6)

### Not shown

Calyx; Cover Sprays; Pre- or Postharvest

Illustration courtesy of Washington State University Extension.



## Apple Pest Control Recommendations

Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.

Delayed Dormant (Stages 1–2)		
Pest or disease/ Material	Amount of product per acre	Comments/Reentry interval/Preharvest interval (PHI)
<b>European red mite eggs, scale, aphid eggs, and apple rust mite</b>		
<i>Note:</i> Delayed Dormant Stage is the best time to control San Jose scale. See footnote 1, page 9.		
Horticultural mineral oil (HMO) + one of the following:	4–8 gal	6-hour reentry.
diazinon 50W	4 lb	Limited to 2 applications per season. 24-hour reentry.
lime sulfur (Ca polysulfides 29%)	5–10 gal	2-day reentry.
Lorsban 4EC	4 pt	Do not use the EC formulation of Lorsban after the delayed dormant period. 24-hour reentry.
Supracide 2E	8 pt	Detrimental to predatory mites with this timing. Do not use Supracide after the delayed dormant period. 2- to 14-day reentry.
<b>Crown and collar rot</b>		
<i>Note:</i> Aliette, Agri-Fos, Fosphite, and Phostrol also registered but may be more useful in the fall.		
Ridomil Gold SL	0.5 pt/ 100 gal water	Rates are based on tree size. Have rain or irrigation move material into root zone. 48-hour reentry.

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

**Prepink or Green Bud (Stages 3–4—little leaves separating just enough to expose blossom bud cluster)**

<b>Pest or disease/ Material</b>	<b>Amount of product per acre</b>	<b>Comments/Reentry interval/Preharvest interval (PHI)</b>
<b>Scab (see footnote 5 and footnote 6, page 9, and Table 1, page 10)</b>		
Adament 50WG	4–5 oz	Group 3 + 11 fungicide. 12-hour reentry. <b>75-day PHI.</b>
Cabrio EG	9.5 oz	May be used at harvest. 12-hr reentry.
Captan 80WDG	2.5–5 lb	See footnote 3, page 9. 24-hour reentry. <b>0-day PHI.</b>
Flint 50WG	2–2.5 oz	12-hour reentry. <b>14-day PHI.</b>
Indar 2F	6–8 fl oz	Add a wetting agent. 12-hour reentry. <b>14-day PHI.</b>
Inspire Super	12 fl oz	Group 3 + 9 fungicide. 12-hour reentry. <b>14-day PHI.</b>
Luna Sensation	4–5.8 fl oz	Group 7 + 11 fungicide. 12-hour reentry. <b>14-day PHI.</b>
Luna Tranquility	11.2–16 fl oz	Group 7 + 9 fungicide. 12-hour reentry. <b>72-day PHI.</b>
mancozeb	6 lb	Do not use this rate beyond bloom. 24-hour reentry. <b>77-day PHI.</b>
Merivon	4–5.5 fl oz	Group 7 + 11 fungicide. 12-hour reentry. <b>0-day PHI.</b>
Omega 500 F	10–13.8 fl oz	12-hr reentry. <b>28-day PHI.</b>
Polyram 80DF	6 lb	Do not use this rate beyond bloom. 24-hour reentry. <b>77-day PHI.</b>
Pristine	14.5–18.5 oz	Group 7 + 11 fungicide. 12-hour reentry. <b>0-day PHI.</b>
Procure	8–16 fl oz	See footnote 5, page 9. Should be tank-mixed with a product that has good protection activity. 12-hour reentry. <b>14-day PHI.</b>
Rally 40WSP	5–8 oz	Do not apply more than 5 lb/A per season. Should be tank-mixed with a product that has good protection activity. 24-hour reentry. <b>14-day PHI.</b>
Sovran	3.2–6.4 oz	See footnote 10, page 9. 12-hour reentry. <b>30-day PHI.</b>
Sulforix	2 qt/100 gal water	See footnote 2, page 9.
Syllit FL	1.5–3 pt	See footnote 4, page 9. 2-day reentry. <b>7-day PHI.</b>
Tebuzol 45DF	4–8 oz	Group 3 fungicide. 5-day reentry. <b>75-day PHI.</b>
TopGuard	8–12 fl oz	Group 3 fungicide. 12-hour reentry. <b>14-day PHI.</b>

*Prepink or Green Bud continues on next page*

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

**CONTINUED—Prepink or Green Bud (Stages 3–4—little leaves separating just enough to expose blossom bud cluster)**

<b>Pest or disease/ Material</b>	<b>Amount of product per acre</b>	<b>Comments/Reentry interval/Preharvest interval (PHI)</b>
<b>Powdery mildew</b>		
Adament 50WG	4–5 oz	Group 3 + 11 fungicide. 12-hour reentry. <b>75-day PHI.</b>
Cabrio EG	9.5 oz	May be used at harvest. 12-hr reentry.
Flint 50WG	2–2.5 oz	12-hour reentry. <b>14-day PHI.</b>
Fontelis	16–20 fl oz	12-hour reentry. <b>28-day PHI.</b>
HMO such as JMS Stylet oil	1–2 gal/ 100 gal water	Do not use past second cover or near sulfur sprays or on wet foliage. 4-hour reentry.
Indar 2F	6–8 fl oz	Add a wetting agent. 12-hour reentry. <b>14-day PHI.</b>
Inspire Super	12 fl oz	Group 3 + 9 fungicide. 12-hour reentry. <b>14-day PHI.</b>
Luna Privilege	2.4–6.84 fl oz	Tank mix with another fungicide. 12-hour reentry. <b>7-day PHI.</b>
Luna Sensation	5–5.8 fl oz	Group 7 + 11 fungicide. 12-hour reentry. <b>14-day PHI.</b>
Luna Tranquility	11.2–16 fl oz	Group 7 + 9 fungicide. 12-hour reentry. <b>72-day PHI.</b>
Merivon	4–5.5 fl oz	Group 7 + 11 fungicide. 12-hour reentry. <b>0-day PHI.</b>
Oso SC	3.75–13 fl oz	4-hour reentry. <b>0-day PHI.</b>
Ph-D WDG	6.2 oz	4-hour reentry. <b>0-day PHI.</b>
Pristine	14.5–18.5 oz	Group 7 + 11 fungicide. 12-hour reentry. <b>0-day PHI.</b>
Procure	8–16 fl oz	12-hour reentry. <b>14-day PHI.</b>
Rally 40WSP	5–10 oz	Do not exceed 5 lb/A per season. 24-hour reentry. <b>14-day PHI.</b>
Sovran	4–6.4 oz	See footnote 10, page 9. 12-hour reentry. <b>30-day PHI.</b>
Sulforix	2 qt/100 gal water	See footnote 2, page 9.
Tebuzol 45DF	4–8 oz	Group 3 fungicide. 5-day reentry. <b>75-day PHI.</b>
TopGuard	8–12 fl oz	Group 3 fungicide. 12-hour reentry. <b>14-day PHI.</b>
<b>Green fruit worm, leafrollers, aphids, plant bugs</b>		
Delegate	4.5–7 oz	<b>7-day PHI.</b>
endosulfan	4–5 lb	2-day reentry. <b>21-day PHI. Endosulfan use will be canceled on apples on July 30, 2015.</b>
<b>Green fruitworm, leafrollers, aphids</b>		
diazinon 50WP	4 lb	Limited to 2 applications per season. 24-hour reentry. <b>21-day PHI.</b>
Lorsban 50W	3 lb	24-hour reentry. <b>28-day PHI.</b>
<b>Green fruitworm, leafrollers, tentiform leafminer</b>		
<i>Note: Tentiform leafminer is a concern only if it was a problem the previous season and low levels of parasitism were noticed.</i>		
Altacor	2.5–4 oz	<b>14-day PHI.</b>
Delegate	4.5–7 oz	<b>7-day PHI.</b>
Imidan 70WP	3.5–5 lb	24-hour reentry. <b>7-day PHI.</b>
Proclaim 5SG	3.2–4.8 oz	<b>14-day PHI.</b>
Success 2L	6–10 oz	Do not apply more than 29 oz/A per season. <b>7-day PHI.</b>

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

**Pink or Preblossom (Stages 5–6—just before blossoms open)**

<b>Pest or disease/ Material</b>	<b>Amount of product per acre</b>	<b>Comments/Reentry interval/Preharvest interval (PHI)</b>
<b>Apple rust mite</b>		
Vendex 50WP	1–1.5 lb	2-day reentry. <b>14-day PHI.</b>

**Scab and powdery mildew**

See materials listed for Prepink or Green Bud Stage.

**Codling moth (mating disruption)**

*Note:* Other products are available, but experience is limited with those products. If pest pressure is high, combine with one or more insecticides against the first generation. Treat with insecticides against the second generation if pressure remains high.

Checkmate	200 ties	—
Isomate C+	200–400 ties	—
Isomate CTT	200 ties	—
No mate	200–400 ties	—

**Calyx (when three-fourths of petals have fallen; apply before calyx closes on central fruit cluster)**

<b>Pest or disease/ Material</b>	<b>Amount of product per acre</b>	<b>Comments/Reentry interval/Preharvest interval (PHI)</b>
<b>Scab (See footnote 5 and footnote 6, page 9 and Table 1, page 10)</b>		
Adament 50WG	4–5 oz	Group 3 + 11 fungicide. 12-hour reentry. <b>75-day PHI.</b>
Cabrio EG	9.5 oz	May be used at harvest. 12-hr reentry.
Captan 80WDG	2.5–5 lb	See footnote 3, page 9. 24-hour reentry. <b>0-day PHI.</b>
Flint 50WG	2–2.5 oz	12-hour reentry. <b>14-day PHI.</b>
Fontelis	16–20 fl oz	Tank-mix with another fungicide and use after bloom. 12-hour reentry. <b>28-day PHI.</b>
Indar 2F	6–8 fl oz	Add a wetting agent. 12-hour reentry. <b>14-day PHI.</b>
Inspire Super	12 fl oz	Group 3 + 9 fungicide. 12-hour reentry. <b>14-day PHI.</b>
Luna Sensation	4–5.8 fl oz	Group 7 + 11 fungicide. 12-hour reentry. <b>14-day PHI.</b>
Luna Tranquility	11.2–16 fl oz	Group 7 + 9 fungicide. 12-hour reentry. <b>72-day PHI.</b>
mancozeb	3 lb	24-hour reentry. <b>77-day PHI.</b>
Merivon	4–5.5 fl oz	Group 7 + 11 fungicide. 12-hour reentry. <b>0-day PHI.</b>
Omega 500 F	10–13.8 fl oz	12-hr reentry. <b>28-day PHI.</b>
Polyram 80DF	3 lb	24-hour reentry. <b>77-day PHI.</b>
Pristine	14.5–18.5 oz	Group 7 + 11 fungicide. 12-hour reentry. <b>0-day PHI.</b>
Procure	8–16 fl oz	12-hour reentry. <b>14-day PHI.</b>
Rally 40WSP	5–8 oz	Do not apply more than 5 lb/A per season. Should be tank-mixed with a product that has good protection activity. 24-hour reentry. <b>14-day PHI.</b>
Scala SC	5–10 oz	Tank-mix with another fungicide and use after bloom. 12-hour reentry. <b>72-day PHI.</b>
Sovran	3.2–6.4 oz	See footnote 10, page 9. 12-hour reentry. <b>30-day PHI.</b>
Sulforix	2 qt/100 gal water	See footnote 2, page 9.
Syllit FL	1.5–3 pt	See footnote 4, page 9. 2-day reentry. <b>7-day PHI.</b>

*Calyx continues on next page*

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

**CONTINUED—Calyx (when three-fourths of petals have fallen; apply before calyx closes on central fruit cluster)**

<b>Pest or disease/ Material</b>	<b>Amount of product per acre</b>	<b>Comments/Reentry interval/Preharvest interval (PHI)</b>
<b>CONTINUED—Scab (See footnote 5 and footnote 6, page 9 and Table 1, page 10)</b>		
Tebuzol 45DF	4–8 oz	Group 3 fungicide. 5-day reentry. <b>75-day PHI.</b>
TopGuard	8–12 fl oz	Group 3 fungicide. 12-hour reentry. <b>14-day PHI.</b>
Ziram 76DF	6 lb	2-day reentry. <b>14-day PHI.</b>
<b>Powdery mildew</b>		
Adament 50WG	4–5 oz	Group 3 + 11 fungicide. 12-hour reentry. <b>75-day PHI.</b>
Cabrio EG	9.5 oz	May be used at harvest. 12-hr reentry.
Flint 50WG	2–2.5 oz	12-hour reentry. <b>14-day PHI.</b>
Fontelis	16–20 fl oz	Tank mix with another fungicide. 12-hour reentry. <b>28-day PHI.</b>
Indar 2F	6–8 fl oz	Add a wetting agent. 12-hour reentry. <b>14-day PHI.</b>
Inspire Super	12 fl oz	Group 3 + 9 fungicide. 12-hour reentry. <b>14-day PHI.</b>
JMS Stylet oil	1–2 gal/ 100 gal water	Do not use past second cover or near sulfur sprays or on wet foliage. 4-hour reentry.
Luna Sensation	5–5.8 fl oz	Group 7 + 11 fungicide. 12-hour reentry. <b>14-day PHI.</b>
Luna Tranquility	11.2–16 fl oz	Group 7 + 9 fungicide. 12-hour reentry. <b>72-day PHI.</b>
Merivon	4–5.5 fl oz	Group 7 + 11 fungicide. 12-hour reentry. <b>0-day PHI.</b>
Oso SC	3.75–13 fl oz	4-hour reentry. <b>0-day PHI.</b>
Ph-D WDG	6.2 oz	4-hour reentry. <b>0-day PHI.</b>
Pristine	14.5–18.5 oz	Group 7 + 11 fungicide. 12-hour reentry. <b>0-day PHI.</b>
Procure	8–16 fl oz	12-hour reentry. <b>14-day PHI.</b>
Rally 40WSP	5–10 oz	Do not apply more than 5 lb/A per season. 24-hour reentry. <b>14-day PHI.</b>
Sovran	4–6.4 oz	See footnote 10, page 9. 12-hour reentry. <b>30-day PHI.</b>
Sulforix	2 qt/100 gal water	See footnote 2, page 9.
Tebuzol 45DF	4–8 oz	Group 3 fungicide. 5-day reentry. <b>75-day PHI.</b>
TopGuard	8–12 fl oz	Group 3 fungicide. 12-hour reentry. <b>14-day PHI.</b>

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

**Cover Sprays (1–4 cover sprays may be needed)**

<b>Pest or disease/ Material</b>	<b>Amount of product per acre</b>	<b>Comments/Reentry interval/Preharvest interval (PHI)</b>
<b>Codling moth, leafrollers</b>		
Altacor	2.5–4 oz	<b>5-day PHI.</b>
Assail 70W	3.4 oz	<b>7-day PHI.</b>
Avaunt 30WG	6 oz	<b>28-day PHI.</b>
Belt SC	5 oz	<b>14-day PHI.</b>
Calypso 4F	2–4 oz	<b>30-day PHI.</b>
Danitol 2.4EC	16–21.3 oz	<b>14-day PHI.</b>
Delegate	6–7 oz	<b>7-day PHI.</b>
diazinon 50WP	4 lb	Do not apply more than 4 lb per application or more than 12 lb per season. 24-hour reentry. <b>21-day PHI.</b>
Esteem 35WP	4–5 lb	The addition of 1% horticultural mineral oil has been shown to increase codling moth control of Esteem. <b>45-day PHI.</b>
Imidan 70WP	4–5 lb	A water-soluble bag formulation (70WSB) also is available. 24-hour reentry. <b>7-day PHI.</b>
Intrepid 2F	16 oz	For use against low to moderate pest pressure situations. <b>14-day PHI.</b>
Proclaim 5SG	3.2–4.8 oz	For codling moth, provides suppression only. <b>14-day PHI.</b>
Rimon 0.83EC	30–50 oz	See label for timing. <b>14-day PHI.</b>
<b>Codling moth, aphids, leafrollers, scale crawlers</b>		
diazinon 50WP	4 lb	24-hour reentry. <b>21-day PHI.</b>
<b>White apple leafhopper</b>		
Actara	2–2.75 oz	<b>35-day PHI.</b>
Assail 70WP	1.1–1.7 oz	<b>7-day PHI.</b>
endosulfan 50WP	4–5 lb	Do not exceed 2 applications per year or a maximum of 6 lb/A per season. See footnote 8, page 9. 2-day reentry. <b>21-day PHI. Endosulfan use will be canceled on apples on July 30, 2015.</b>
Provado 1.6F	4–8 oz	<b>7-day PHI.</b>
<b>Mites</b>		
Acramite 50WS	0.75–1 lb	12-hour reentry. <b>7-day PHI.</b>
Apollo 50SC	4–8 oz	Apply only once per season. Will not control rust mites. <b>45-day PHI.</b>
Envidor 2SC	16–18 oz	Apply only once per season. <b>7-day PHI.</b>
FujiMite 5EC	32 oz	Do not apply more than twice per season. <b>14-day PHI.</b>
Kanemite 50WSP	4 lb	<b>7-day PHI.</b>
Onager	16–24 oz	<b>28-day PHI.</b>
Vendex 50WP	1–2 lb	2-day reentry. <b>14-day PHI.</b>
Zeal 72WDG	2–3 oz	Apply only once per season. <b>28-day PHI.</b>

*Cover Sprays continues on next page*

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

**CONTINUED—Cover Sprays (1–4 cover sprays may be needed)**

<b>Pest or disease/ Material</b>	<b>Amount of product per acre</b>	<b>Comments/Reentry interval/Preharvest interval (PHI)</b>
<b>Bull's eye rot and scab</b>		
Captan 80WDG	2.5–5 lb	24-hour reentry. <b>0-day PHI.</b>
mancozeb	3 lb	24-hour reentry. <b>77-day PHI.</b>
Ziram 76DF	6 lb	2-day reentry. <b>14-day PHI.</b>

**Scab and powdery mildew**

*Note:* See materials listed for Calyx Stage. Apply scab sprays before wet weather is expected to occur and stop when dry weather prevails. Powdery mildew sprays can be stopped when terminal growth stops.

**Anthracnose**

*Note:* Scout for cankers in trees. Remove and destroy cankers during dry weather.

**Apple maggot**

*Note:* Sprays used for codling moth will control apple maggot. However, 1 or 2 additional sprays for apple maggot may be required later in the season.

Assail 70WP	3.4 oz	<b>7-day PHI.</b>
Belay 50WDG	3 oz	<b>7-day PHI.</b>
Delegate 25WG	6–7 oz	<b>7-day PHI.</b>
Imidan 70WP	3–5 lb	A water-soluble bag formulation (70WSB) also is available. 24-hour reentry. <b>7-day PHI.</b>

**Pre- or Postharvest (before fall rains—see footnote 9, page 9)**

<b>Pest or disease/ Material</b>	<b>Amount of product per acre</b>	<b>Comments/Reentry interval/Preharvest interval (PHI)</b>
<b>Anthracnose, Nectria canker, Bull's eye rot</b>		
bordeaux 6-6-100	—	Do not use on yellow-colored cultivars before harvest.
Captan 80WDG	3.75 lb	Do not apply more than 64 lb/A per year. 24-hour reentry. <b>0-day PHI.</b>
Copper-Count-N	10 qt	Postharvest only. 48-hour reentry.
Cuprofix Ultra 40 Disperss	8–20 lb	Postharvest only. 48-hour reentry.
Kocide 3000	5.25–7 lb	Do not use on yellow-colored cultivars before harvest. 48-hour reentry.
Nu-Cop 50DF	12–16 lb	Do not use on yellow-colored cultivars before harvest. 48-hour reentry.
Ziram 76DF	6 lb	48-hour reentry. <b>14-day PHI.</b>

**Crown and collar rot**

*Note:* Ridomil is also registered but may be more useful in the spring.

Agri-Fos	1.25–2.5 qt	Do not use with copper materials. 4-hour reentry.
Aliette WDG	2.5–5 lb	Do not use with copper materials. 12-hour reentry. <b>14-day PHI.</b>
Fosphite	1–3 qt	Do not use with copper materials. 4-hour reentry.
Phostrol	2.5–5 pt	4-hour reentry.

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*



## Footnotes

1. Use oil emulsion, 3.2% actual oil, plus bordeaux 6-6-100. This spray will control all other pests listed except blister mite. Bordeaux is not compatible with lime sulfur or polysulfide.
2. Lime sulfur may injure Delicious and Delicious strains during hot weather and causes yellow foliage on Braeburn. Lime sulfur will help control apple rust mite.
3. Captan may cause minor leaf spotting to Delicious under certain conditions.
4. Syllit is not compatible with lime and should not be combined with oils or oil emulsions.
5. Apple scab forecasting is useful when spring rains become less frequent and drier weather prevails. Several materials can be applied within a certain time limit after the start of an infection period. Keep to a protection schedule throughout the bloom period. All ascospores will have matured and be ready for dispersal once 865 degree days (base 32°F) have accumulated since bud break. Group 11 materials such as Flint and Sovran claim long kickback activity. These claims are doubtful, and kickback activity may be much shorter. These materials are best used **prior** to infection periods.
6. To delay or prevent the development of resistant strains of apple scab or powdery mildew, alternate or tank-mix materials with different modes of activity (or from different fungicide groups).
7. Codling moth: spray timing. **CALENDAR APPROACH:** First spray at 15 to 21 days after petal fall followed by another in about 3 weeks. Third spray for second generation usually is made in early July followed by another in about 3 weeks. **PHEROMONE TRAPS TO TIME SPRAYS:** Mid-May place one trap for every 3 acres in upper one-third of the tree canopy. Inspect once weekly or more frequently. Make first spray when two or more moths are caught in one or more of the traps for 2 weeks in a row.  
  
Repeat spray when first application has weathered off (usually 3 weeks for azinphos methyl and Guthion) and two or more moths are caught in one or more of the traps. Spot treatments may be sufficient in parts of blocks. Continue trapping through September. **DEGREE DAY ACCUMULATION:** first spray at 250 degree days following first consistent catch of codling moths in pheromone traps (50°F lower threshold).
8. White apple leafhopper has become a serious problem for some growers in the Willamette Valley. It is best controlled during the first generation after egg hatch is complete but before there are a large number of mature, winged adults. Larger nymphs and adults are difficult to control. Use Thiodan 50WP around petal fall or shortly after. Note that timing of the first cover spray for codling moth may be too late to control leafhoppers. Also the commonly used codling moth insecticides are not that effective on leafhoppers. An application of Sevin (carbaryl) directed at the second-generation nymphs, which should be present in August, usually provides sufficient control of leafhoppers to prevent picker annoyance problems. **DO NOT USE CARBARYL (SEVIN) DURING PETAL FALL (FIRST LEAFHOPPER SPRAY) AS FRUIT THINNING WILL OCCUR. USE THIODAN.** Endosulfan use will be canceled on apples on July 30, 2015.
9. Use Captan or Ziram preharvest for control of Bull's eye rot. Focus on early- and mid-leaf fall for control of Nectria canker. Do not use Topsin as it is toxic to earthworms, which help decompose scab-infected leaves.
10. Sovran drift may injure some sweet cherry cultivars such as Van. Please be extra careful when spraying near cherry orchards.

---

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

**Table 1. Approximate Hours of Wetness  
at Indicated Temperatures Required for Leaf Scab Infection  
and Days Required for Lesions to Appear**

Average temperature (°F)	Hours of wetness required for infection* From primary and secondary inoculum			Days required for lesions to appear**
	Light	Moderate	Heavy	
78	13	17	26	—
77	11	14	21	—
76	9.5	12	19	—
63–75	9	12	18	9
62	9	12	19	10
61	9	13	20	10
60	9.5	13	20	11
59	10	13	21	12
58	10	14	21	12
57	10	14	22	13
56	11	15	22	13
55	11	16	24	14
54	11.5	16	24	14
53	12	17	25	15
52	12	18	26	15
51	13	18	27	16
50	14	19	29	16
49	14.5	20	30	17
48	15	20	30	17
47	15	23	35	—
46	16	24	37	—
45	17	26	40	—
44	19	28	43	—
43	21	30	47	—
42	23	33	50	—
41	26	37	53	—
40	29	41	56	—
39	33	45	60	—
38	37	50	64	—
37	41	55	68	—
33–36	48	72	96	—

From W.D. Mills, Cornell University.

\*Leaves remain wet for varying lengths of time after the rain stops, depending on conditions. Add together wetting periods from intermittent showers. Add together any wet periods with less than 8 hours dry time between them. Determine average temperature for the period from hourly readings. Lesions may not be apparent for 2–4 weeks.

\*\*Days required for conidia to appear once infection has been established. No further wetting is required. For this column, daily maximum and minimum temperatures are adequate for determining the average.

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

**Table 2. Effectiveness of Fungicides for Apple Disease Management\***

Fungicide	Fungicide group #	Apple scab	Powdery mildew	Bull's eye rot
Cabrio	11	Good**	Excellent**	?? (good)
Captan	M4	Excellent	None	Good
Flint	11	Excellent**	Good–Excellent**	Slight–Fair
Focus	3	Good**	Excellent**	??
Fontelis	7	Fair–Good**	Good**	??
HMO***	Not classified	??	Good	??
Indar	3	Good**	Excellent**	??
Kaligreen	Not classified	None	Slight–Fair	??
Lime sulfur	M2	Excellent	Good	??
Luna Privilege	7	Fair–Good**	Good**	??
Mancozeb	M3	Good	None	Slight–Fair
Omega	29	Good	None–poor	??
Oso	19	??	Fair–Good	??
Ph-D	19	??	Fair–Good	??
Polyram	M3	Good	None	??
Procure	3	Good**	Excellent**	Slight–Fair
Rally	3	Good**	Fair–Good	??
Scala	9	Fair	None	??
Sovran	11	Excellent**	Good–Excellent**	??
Sulfur	M2	Fair	Good	??
Syllit	M7	Good**	None	??
Tebuzol	3	Good**	Excellent**	??
TopGuard	3	Good**	Excellent**	??
Topsin	1	Fair**	Fair–Good**	Excellent**
Vanguard	9	Fair**	None	??
Ziram	M3	Fair	None	Fair–Good
<b>Combination products</b>				
Adament	3 + 11	Excellent	Excellent	??
Inspire Super	3 + 9	Good	Excellent	??
Luna Sensation	7 + 11	Excellent	Excellent	??
Luna Tranquility	7 + 9	Good–Excellent	Excellent	??
Merivon	7 + 11	Excellent	Excellent	??
Pristine	7 + 11	Good–Excellent**	Excellent**	Good

\*These ratings are relative rankings based on labeled application rates, good spray coverage, and proper spray timing. Actual levels of disease control will be influenced by these factors in addition to cultivar susceptibility, disease pressure, and weather conditions.

\*\*Resistant pathogens will lower the effectiveness of these fungicides.

\*\*\*Horticultural mineral oil.

### Follow the “Rules” for fungicide stewardship:

Rotate or mix fungicides of different chemical groups.  
 Use labeled rates.  
 Limit total number of applications.  
 Educate yourself about fungicide activity, mode of action, and class—as well as resistance management practices.  
 Start a fungicide program with multisite mode of action materials.

For more information about fungicides registered for use on apples and their specific modes of action, consult OSU Extension publication EM 8950: *How to Reduce the Risk of Pesticide Resistance in Apple Pests in Oregon*.

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

## Table 3. Quick Reference Guide for Herbicides Labeled for Use in Fruit and Nut Crops

- Shaded boxes indicate the herbicide is labeled for use in that crop.
- Nonbearing (NB) indicates the herbicide is labeled only for crops that will not be harvested for 1 year. It has a 365-day preharvest interval.
- Herbicides in **bold, italic** type are recommended for new plantings.
- For more complete information, please refer to the *PNW Weed Management Handbook*: <http://pnwhandbooks.org/weed/>.

Ingredient common name (herbicide mode of action)	Product name example	Nuts			Pome fruits		Stone fruits						Rates
		Chestnut	Hazelnut	Walnut	Apple	Pear	Apricot	Cherry	Nectarine	Peach	Plum	Prune	
<b>Applications that persist in soil and are soil active</b>													
diuron (7)	Karmex												See label for crop-specific application rates.
dichlobenil (20)	Casoron												4–6 lb ai/A (100–150 lb/A Casoron). Apply in cold, wet weather.
<i>isoxaben</i> (21)	<i>Trellis, Gallery</i>				NB	NB	NB	NB	NB	NB	NB	NB	0.5–1 lb ai/A (0.66–1.33 lb/A product)
indaziflam (29)	Alion												0.065–0.085 lb ai/A (5–6.5 oz/A product) depending on soil texture
<i>napropamide</i> (3)	<i>Devrinol</i>												5 lb ai/A (50 lb/A Devrinol 10-G)
norflurazon (12)	Solicam												1.97–7.8 lb ai/A (2.5–10 lb/A Solicam)
<i>oryzalin</i> (3)	<i>Surflan</i>												2–6 lb ai/A (2–6 qt/A Surflan)
<i>pendimethalin</i> (3)	<i>Prowl</i>												Prowl H <sub>2</sub> O: 1.9–6 lb ai/A (2–6.3 qt/A) depending on desired length of weed control and crop
<i>pronamide</i> (3)	<i>Kerb</i>												1–4 lb ai/A (2–8 lb/A). Rate depends on species present and soil texture.
simazine (5)	Princep												See product labels for rates. Princep Caliber 90 is a Special Local Needs label (OR-080038) for sweet cherries only.
terbacil (5)	Sinbar					NB	NB	NB				NB	0.4–0.8 lb ai/A (0.5–1 lb/A), newly established; 2–4 lb/A Sinbar, bearing, depending on soil type
<i>trifluralin</i> (3)	<i>Treflan</i>												0.5–1 lb ai/A (1–2 pt/A Treflan 4EC)
trifluralin (3) + isoxaben (21) + oxyfluorfen (14)	Showcase	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	2.5–5 lb ai/A (100–200 lb/A Showcase)
<b>Applications that persist in soil and have both soil and foliar activity</b>													
<i>flumioxazin</i> (14)	<i>Chateau</i>												0.188–0.38 lb ai/A (6–12 oz/A Chateau WDG). Note differences in rates and uses in SW and WDG labels. Avoid contact with green bark on small trees.
oxyfluorfen (14)	Goal												1.25–2 lb ai/A (5–8 pt/A Goal 2XL)

*Table continues on next page*

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

Ingredient common name (herbicide mode of action)	Product name example	Nuts			Pome fruits		Stone fruits						Rates
		Chestnut	Hazelnut	Walnut	Apple	Pear	Apricot	Cherry	Nectarine	Peach	Plum	Prune	
<b>CONTINUED—Applications that persist in soil and have both soil and foliar activity</b>													
penoxsulam (2)	Pindar GT												(1.5–3.0 pints/A)
rimsulfuron (2)	Matrix												0.063 lb ai/A (4 oz/A Matrix FNV per year)
saflufenacil (14)	Treevix												0.045 lb ai/A (1 oz/A Treevix)
<b>Postemergence contact and translocated herbicides</b>													
acetic acid	WeedPharm												
carfentrazone (14)	Aim												Green sucker control in hazelnuts. 0.031 lb ai/A (2 fl oz/A Aim EC)
clethodim (1)	Select Max		NB	NB	NB	NB	NB	NB		NB		NB	0.068–0.121 lb ai/A (9–16 fl oz/A Select Max)
clopyralid (4)	Stinger												Apples: 0.094–0.25 lb ae/A (0.25–0.66 pints/A Stinger) Others: 0.12–0.25 lbs ae/A (0.33–0.66 pints/A Stinger)
diquat (22)	Reglone		NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	0.375–0.5 lbs ai/A (1.5–2 pints)
fluazifop (1)	Fusilade DX		NB	NB	NB	NB							0.25–0.375 lb ai/A (16–24 oz/A Fusilade DX). Refer to specific grassy weeds listed on label.
glufosinate (10)	Rely 280												Sucker control. 0.88–1.46 lb ai/A (1.5 to 2.5 quarts/A Rely)
glyphosate (9)	Roundup												General weed control and grass suppression in row middles. Read label carefully for crops listed and geographic location.
halosulfuron (2)	Sandea												Apples: 0.035–0.094 lb ai/A (0.75–2 oz/A) Nut crops: 0.031–0.063 lb ai/A (2/3–1 1/2 oz/A)
paraquat (22)	Gramoxone												Green sucker control in hazelnuts. 0.625–1 lb cation/A (2.5–4 pt/A Gramoxone; 1.7–2.7 pt/A Firestorm)
pyraflufen (14)	Venue												0.001–0.005 lb ai/A (0.7–4 fl oz product/A)
sethoxydim (1)	Poast										NB	NB	Grass suppression in row middles. 0.28–0.47 lb ai/A (1.5–2.5 pt/A product)
2,4-D (4)	2,4-D												Green sucker control in hazelnuts. 0.7–0.95 lb ai/A (1.5–2 pt/A Saber)

Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.

## OSU Internet Resources for Plant Protection

Information regarding plant protection is available from several sources at OSU. The following listings are excellent examples:

- OSU Integrated Plant Protection Center. Online weather data and degree day information for insect pests and diseases (<http://uspest.org/wea/>)
- Codling moth development information (<http://ippc2.orst.edu/cgi-bin/ddmodel.pl?clm>)
- Apple scab infection season information (<http://ippc2.orst.edu/cgi-bin/ddmodel.pl?spp=asc>)
- Fire blight risk information (<http://ippc2.orst.edu/cgi-bin/ddmodel.pl?fbl>)  
Directions for the use of each model are available at each site.
- Pacific Northwest Plant Disease Management Handbook (<http://pnwhandbooks.org/plantdisease>)
- Pacific Northwest Insect Management Handbook (<http://pnwhandbooks.org/insect>)
- Pacific Northwest Weed Management Handbook (<http://pnwhandbooks.org/weed>)

### Oregon Poison Center

The Oregon Health & Science University  
3181 S.W. Sam Jackson Park Road  
Portland, OR 97239  
Phone: 1-800-222-1222

**If a person has collapsed or is not breathing, dial 911.**

## Basic Elements of Safe Pesticide Use

- Always read the label with care. This is the first step in selecting the right material for the job. Never rely on your memory. Before opening the container, pay strict attention to warnings and cautions printed on the label.
- Keep all pesticide and spray materials out of the reach of children, pets, and irresponsible persons. Storage outside of the home, away from food and feed, and under lock and key is the safest method.
- Store only in the original container and keep tightly closed.
- NEVER smoke, eat, or drink while applying pesticides.
- Avoid inhalation or direct contact. Always wear protective clothing and safety devices as recommended on the label.
- Avoid spills. If spills occur, take immediate action to remove contaminated clothing and wash thoroughly.
- After each application, bathe and change to clean clothing. Wash clothing after each use. Always use fresh clothing when starting new application.
- Avoid contamination of fish ponds and water supplies. Cover feed and water containers when treating around livestock or pet areas.
- Keep separate equipment for use with hormone-type herbicides to avoid accidental injury to susceptible plants. Also avoid applications under wind conditions that could create drift to nontarget areas.
- Rinse empty containers three times before disposing of them. Add the rinse to the spray tank and dispose of containers according to local regulations to avoid hazard to humans, animals, and the environment.
- Follow label directions for mixing and application to keep residues within the limits prescribed by law.
- Plan ahead. Discuss with your physician the materials you will be using during the season so that he or she can be prepared to provide the appropriate treatment in case of accidental exposure. If symptoms of illness occur, call the physician or get the patient to a hospital immediately. Always provide the medical personnel with as much information as possible.
- Be cautious when you apply pesticides. Know your legal responsibility as a pesticide applicator. You may be liable for injury or damage resulting from pesticide use.

---

Prepared by Jay W. Pscheidt, Professor and Extension plant pathology specialist, Department of Botany and Plant Pathology; Ed Peachey, assistant professor of weed science, Department of Horticulture; and Steve Castagnoli, Associate Professor, Tree Fruits Research & Extension Horticulturalist for Hood River County, MCAREC; all of Oregon State University. The information in this pest management guide is valid for 2014. Trade-name products and services are mentioned as illustrations only. This does not mean that the Oregon State University Extension Service either endorses these products and services or intends to discriminate against products and services not mentioned. Due to constantly changing laws and regulations, the Oregon State University Extension Service can assume no liability for the suggested use of chemicals contained in this guide. Pesticides should be applied according to the label directions on the pesticide container.

---

Dedicated to the memory of Jeff Olsen, Extension horticulturist, Yamhill County.

---

© 2014 Oregon State University. This publication was produced and distributed in furtherance of the Acts of Congress of May 8 and June 30, 1914. Extension work is a cooperative program of Oregon State University, the U.S. Department of Agriculture, and Oregon counties. Oregon State University Extension Service offers educational programs, activities, and materials without discrimination based on age, color, disability, gender identity or expression, genetic information, marital status, national origin, race, religion, sex, sexual orientation, or veteran's status. Oregon State University Extension Service is an Equal Opportunity Employer.

---

Revised March 2014.

---

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*