

Use 1 quart of wetting agent per 100 gallons of water with amitrole, paraquat, and 2,4-D amine formulations. Use 40 to 50 gallons of water per acre. If you use amitrole, for example, apply 40 to 50 gallons of spray per acre, containing 1 pound active material and a half-quart of wetting agent.

- *Soil-active herbicides.* These materials give season-long weed control. They should not be applied where roots of desirable plants may be present. Rainfall is necessary to leach these chemicals into the soil before the weeds are up.

Use atrazine (AAtrex ®), simazine (Princep ®), bromacil (Hyvar X ®), diuron (Karmex ®), karbutilate (Tandex ®), or monuron (Telvar

®) at 2.4 to 4 pounds active ingredient per acre. This will be 3 to 5 pounds of 80 percent formulation.

Pesticide warning

The user of any pesticide is responsible for residues in a crop or for problems caused by drift from one property to other property or crops. Always follow the label instructions.

Drawings, and some of the information included in this Fact Sheet, appear courtesy of University of California Extension Leaflet 2741.

Revised by Martin Zimmerman, Gilliam County Extension agent, and Homer M. Hepworth, Extension agronomist, Oregon State University.

THIS PUBLICATION IS OUT OF DATE.
For most current information:
<http://extension.oregonstate.edu/catalog>