

INTERNAL REPORT 129

LITERATURE REVIEW OF FOODS OF AQUATIC INSECTS

E. Grafius and N. H. Anderson
Department of Entomology
Oregon State University

NOTICE: This internal report contains information of a preliminary nature, prepared primarily for internal use in the US/IBP Coniferous Forest Biome program. This information is not for use prior to publication unless permission is obtained in writing from the authors.

Introduction

The following list is from published feeding records up to 1972, of selected aquatic insects with some interpretive comments. Food types are listed as 1. diatoms and algae (a); 2. detritus (d); and 3. animal (c). Percentages of each food, when given, are listed in the same order as the respective food categories. Where one category comprised more than 75% of the diet, only the dominant food type is listed. An attempt was made to find references at the species level, but where these were not available, more general references at the generic or familial level are included. Data reported by Coffman (1967) are also shown in Coffman, Cummins and Wuycheck (1971) and those of Minshall (1965) are repeated in Minshall (1967). The original theses are used since data are presented in a more quantitative manner.

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Ephemeroptera			
Baetidae			
<u>Baetis bicaudatus</u>	ad(41, 57) ³	Gilpin & Brusven, 1970	38 specimens, 26 empty
<u>B. herodes</u>	d(96) ²	Minshall, 1965	measured monthly
<u>B. parvus</u>	ad(41, 53) ³	Gilpin & Grusven, 1970	10 specimens, 1 empty
<u>B. phoebus</u>	d(78) ²	Minshall, 1965	measured monthly
<u>B. rhodani</u>	ad	Jones, 1950	fed on red algae
<u>B. tricaudatus</u>	ad(45, 52) ³	Gilpin & Brusven, 1970	74 specimens, 31 empty
<u>B. vagans</u>	ad(35, 65)	Minckley, 1963	data reported graphically
<u>Baetis</u> sp.	ad(50, 50) ²	Muttkowski & Smith, 1929	2 specimens
<u>Baetis</u> sp.	a	Ivanova, 1958	in lab preference study
<u>Baetis</u> sp.	ad(70, 30) ³	Chapman & Demory, 1963	Deer Cr. 290 specimens
<u>Baetis</u> sp.	ad(55, 45) ³	Chapman & Demory, 1963	Needle Br. 220 specimens
<u>Callibaetis fluctuans</u>	d	Morgan, 1911	fed on artificial diets of ground plant material

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Ephemeroptera			
Baetidae			
<u>Callibaetis</u> sp.	a	Morgan, 1911	---
<u>Centroptilum album</u>	a(88) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>C. luteolum</u>	a	Ivanova, 1958	lab preference study
<u>C. rufostrigatum</u>	d(95) ²	Minshall, 1965	measured monthly
<u>Centroptilum</u> sp.	ad(40, 58) ³	Gilpin & Brusven, 1970	13 specimens, 8 empty
<u>Cloeon dipterum</u>	a	Wissmeyer, 1926	---
<u>C. dipterum</u>	a	Ivanova, 1958	in lab preference study
<u>C. dipterum</u>	a	Brown, 1960	---
<u>Pseudocloeon</u> sp.	a(99) ¹	Coffman, 1967	riffle areas in Aug & Oct
Baetiscidae			
<u>Baetisca bajkovi</u>	a	Lehmkuhl, 1972	---
Caenidae			
<u>Caenis anceps</u>	a(92) ¹	Coffman, 1967	riffle areas in Aug & Oct

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

2

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>References</u>	<u>Remarks</u>
Ephemeroptera (cont.)			
Caenidae			
<u>Caenis horaria</u>	d	Moon, 1938	---
Ephemerellidae			
<u>Drunella</u> sp.	ad(37, 45) ³	Muttkowski & Smith, 1929	25 specimens
<u>Ephemerella aestiva</u>	ad(69, 31) ¹	Coffman, 1967	from riffles in Aug & Oct
<u>E. coloradensis</u>	ad(40, 51) ³	Gilpin & Brusven, 1970	7 specimens
<u>E. doddsi</u>	ad(52, 45) ³	Gilpin & Brusven, 1970	55 specimens, 25 empty
<u>E. doddsi</u>	c	Snow, 1973 unpublished	can be carnivorous on other mayflies
<u>E. dorothea</u>	ad(49, 51) ¹	Coffman, 1967	from riffles in Aug & Oct
<u>E. edmundsi</u>	ad(27, 70) ³	Gilpin & Brusven, 1970	5 specimens, 2 empty
<u>E. flavilinea</u>	ad(41, 56) ³	"	85 specimens, 11 empty
<u>E. grandis</u>	ad(43, 50) ³	"	97 specimens, 16 empty

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Ephemeroptera (cont.)			
EphemereLLidae(cont.)			
<u>E. hecuba</u>	ad(55, 44) ³	Gilpin & Brusven, 1970	41 specimens, 8 empty
<u>E. hystrix</u>	ad(48, 47) ³	"	4 specimens
<u>E. inermes</u> & <u>infrequens</u>	ad(37,60) ³	"	147 specimens, 43 empty
<u>E. ignita</u>	d	Ivanova, 1958	lab preference study
<u>E. margarita</u>	ad(46, 55) ³	Gilpin & Brusven, 1970	20 specimens, 10 empty
<u>E. notata</u>	ad	Jones, 1950	---
<u>E. spinifera</u>	c(80) ³	Gilpin & Brusven, 1970	6 specimens, 5 empty
<u>E. subvaria</u>	a(75) ²	Minckley, 1963	data reported graphically
<u>E. teresa</u>	ad(30, 70) ³	Gilpin & Brusven, 1970	2 specimens
<u>E. tibialis</u>	ad(43, 55) ³	"	96 specimens, 8 empty
<u>Emphemerella</u> sp.	ad(36, 60) ²	Muttkowski & Smith, 1929	28 specimens
<u>Ephemerella</u> sp.	ad(60, 40) ³	Chapman & Demory, 1963	Deer Cr., 7 specimens

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>References</u>	<u>Remarks</u>
Ephemeroptera (cont.)			
Ephemeridae			
<u>Ephemera simulans</u>	c(87) ¹	Coffman, 1967	from riffles in Aug and Oct
<u>E. varia</u>	a(82) ¹	"	"
<u>Hexagenia bilineata</u>	d	Fremling, 1960	---
<u>H. limbata</u>	ad	Hunt, 1953	---
<u>Hexagenia sp.</u>	ad	Morgan, 1911	---
Heptageniidae			
<u>Anepeorus sp.</u>	c	Burks, 1953	on basis of mouthparts
<u>Anepeorus sp.</u>	c	Edmunds, 1957	on basis of mouthparts
<u>Chirotenetes sp.</u>	a	Clemens, 1917	---
<u>Chirotenetes sp.</u>	c	Morgan, 1911	---
<u>Cinygmula sp.</u>	ad(40, 60) ³	Chapman & Demory, 1963	Deer Cr., 229 specimens
<u>Cinygmula sp.</u>	ad(35, 65) ³	Chapman & Demory, 1963	Needle Br., 98 specimens

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

5

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Ephemeroptera (cont.)			
Heptageniidae (cont.)			
<u>Cinygmula</u> sp.	ad(40, 53) ³	Gilpin & Brusven, 1970	128 specimens, 39 empty
<u>Epeorus albertae</u>	ad(46, 54) ³	"	30 specimens, 12 empty
<u>E. grandis</u>	ad(28, 73) ³	"	5 specimens, 3 empty
<u>E. longimanus</u>	ad(38, 61) ³	"	41 specimens, 12 empty
<u>E. pleuralis</u>	d(85) ²	Minshall, J., 1964	25 specimens
<u>Epeorus</u> sp.	ad(60, 40) ³	Chapman & Demory, 1963	Deer Cr., 30 specimens
<u>Epeorus</u> sp.	ad(25, 75) ³	"	Needle Br., 8 specimens
<u>Heptagenia criddlei</u>	ad(55, 45) ³	Gilpin & Brusven 1970	56 specimens, 14 empty
<u>H. lateralis</u>	d	Jones, 1950	---
<u>H. sulphurea</u>	d	Ivanova, 1958	lab preference study
<u>H. sp.</u>	d(76) ²	Muttkowski & Smith, 1929	34 specimens
<u>Heptagenia</u> sp.	ad	Morgan, 1911	---
<u>Pseudiron</u> sp.	c	Edmunds, 1957	on basis of mouthparts
<u>Rhithrogena hageni</u>	ad(45, 55) ³	Gilpin & Brusven, 1970	38 specimens, 11 empty

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Ephemeroptera (cont.)			
Heptageniidae (cont.)			
<u>R. robusta</u>	ad(41, 55) ³	Gilpin & Brusven, 1970	4 specimens, 3 empty
<u>R. semicolorata</u>	d	Jones, 1950	---
<u>Stenonema canadense</u>	a(87) ¹	Coffman, 1967	from riffles in Aug and Oct
<u>S. fuscum</u>	a(80) ¹	"	"
<u>S. tripunctatum</u>	a(79) ¹	"	"
<u>Stenonema</u> sp.	d(80)	Minckley, 1963	data presented graphically
<u>Stenonema</u> undet.	ad(70, 30) ¹	Coffman, 1967	from riffles in Aug & Oct
Isonychidae			
<u>Isonychia albomanicata</u>	c(57) ¹	Coffman, 1967	from riffles in Aug & Oct
<u>Isonychia</u> sp.	c	Burks, 1953	partly predaceous
Leptophlebiidae			
<u>Blasturus</u> sp.	a	Morgan, 1911	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Ephemeroptera (cont.)			
Leptophlebiidae			
<u>Choroterpes</u> sp.	a(79) ¹	Coffman, 1967	from riffles in Aug & Oct
<u>Habrophleboides</u>			
<u>americana</u>	ad(49, 51) ¹	Coffman, 1967	from riffles in Aug & Oct
<u>Leptophlebia marginata</u>	d	Moon, 1938	---
<u>L. vespertina</u>	d	Moon, 1938	---
<u>Paraleptophlebia</u>			
<u>bicornuta</u>	ad(33, 66) ³	Gilpin & Brusven, 1970	30 specimens, 7 empty
<u>P. debilis</u>	ad(36, 61) ³	"	7 specimens, 1 empty
<u>P. heteronea</u> &			
<u>memorialis</u>	ad(31, 65) ³	"	33 specimens, 7 empty
<u>P. columbiae</u>	ad(27, 72) ³	"	6 specimens
<u>P. mollis</u>	d	Hynes & Kaushik, 1968	fed on leaves in lab

a = algal feeding

d = detrital feeding

c = carnivorous

¹ based on food calories

² based on number of food items

³ based on food volume or area

∞

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>References</u>	<u>Remarks</u>
Ephemeroptera (cont.)			
Leptophlebiidae (cont.)			
<u>Paraleptophlebia</u> sp.	d(95) ³	Chapman & Demory, 1963	Deer Cr., 444 specimens
<u>Paraleptophlebia</u> sp.	d(95) ³	Chapman & Demory, 1963	Needle Br., 371 specimens
Oligoneuriidae			
<u>Oligoneuriella rhenana</u>	a	Pinet, 1962	---
Siphonuridae			
<u>Ameletus</u> sp.	d(77) ²	Muttkowski & Smith, 1929	25 specimens
<u>Ameletus</u> sp.	ad(40, 60) ³	Gilpin & Brusven, 1970	57 specimens, 6 empty
<u>Metreturus pecatonia</u>	c	Burks, 1953	on basis of mouthparts
<u>Metreturus</u> sp.	c	Edmunds, 1957	"
<u>Siphonurus lacustris</u>	d	Jones, 1950	---
<u>S. occidentalis</u>	adc	Edmunds, 1960	fed on live mosquito larvae in lab

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Ephemeroptera (cont.)			
Siphonuridae			
<u>S. occidentalis</u>	d(79) ³	Gilpin & Brusven, 1970	5 specimens, 1 empty
<u>Siphonurus</u> sp.	ad	Morgan, 1911	also some mayfly fragments
Tricorythidae			
<u>Tricorythodes minutus</u>	d(83) ¹	Gilpin & Brusven, 1970	5 specimens
Plecoptera			
Filipalpia			
Capniidae			
<u>Allocapnia granulata</u>	ad	Frison, 1929	---
<u>A. vivipara</u>	ad	"	---
<u>A. recta</u>	ad	"	---
<u>A. mystica</u>	ad	"	---
Leuctridae			
<u>Leuctra claasseni</u>	d	"	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>References</u>	<u>Remarks</u>
Plecoptera (cont.)			
Filipalpia (cont.)			
Leuctridae			
<u>L. hippopus</u>	d	Jones, 1950	---
Nemouridae			
<u>Nemoura delosa</u>	d(97) ²	Minshall, 1965	measured monthly
<u>N. vallicularia</u>	d(100) ²	"	"
<u>N. vallicularia</u>	d	Wu, 1923	---
<u>Nemoura</u> sp.	ad	Needham & Claassen, 1925	---
<u>Nemoura</u> sp.	d(100) ³	Chapman & Demory, 1963	23 specimens
Peltoperlidae			
<u>Peltoperla brevis</u>	a(90) ²	Chapman & Demory, 1963	Deer Cr., 4 specimens
<u>P. brevis</u>	d(100) ²	"	Needle Br., 2 specimens
<u>P. maria</u>	d	Wallace, et al, 1970	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Plecoptera (cont.)			
Filipalpia (cont.)			
Pteronarcidae			
<u>Pteronarcys californica</u>	d ²	Muttkowski & Smith, 1929	virtually 100% detritus
<u>P. princeps</u>	d	Grafius, 1974	lab study
<u>P. scotti</u>	d	McDiffett, 1970	lab study
<u>Pteronarcys</u> spp.	ad	Needham & Claassen, 1925	---
Taeniopterygidae			
<u>Brachyptera</u> sp.	a	Hynes, 1941	---
<u>Taeniopteryx fasciata</u>			
(= <u>Strophoteryx</u>)	ad	Frison, 1929	---
<u>T. maura</u>	c(99) ¹	Coffman, 1967	riffle areas in Aug and Oct
<u>T. nivalis</u>	ad	Frison, 1929	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Plecoptera (cont.)			
Setipalpia (cont.)			
Chloroperlidae			
<u>Alloperla</u> sp.	c	Chapman & Demory, 1963	Deer Cr. & Needle Br., 52 specimens
<u>Alloperla</u> sp.	dc	Ellis, 1970	scavengers on dead alevins and salmon eggs
<u>Chloroperla</u> sp.	ad	Frison, 1935	---
<u>Chloroperla</u> sp.	c(99) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Kathroperla perdita</u>	ad(50, 50) ³	Chapman & Demory, 1963	Deer Cr., 6 specimens
<u>K. perdita</u>	d(95) ³	Chapman & Demory, 1963	Needle Br., 20 specimens
Perlidae	c	Frison, 1935	---
<u>Acroneuria californica</u>	c	Chapman & Demory, 1963	Deer Cr. and Needle Br., 35 specimens

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>References</u>	<u>Remarks</u>
Plecoptera (cont.)			
Setipalpia (cont.)			
Perlidae (cont.)			
<u>A. lycorias</u>	c(99) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>A. pacifica</u>	c(77) ²	Muttkowski & Smith, 1929	49 specimens
<u>A. pacifica</u>	c	Chapman & Demory, 1963	Deer Dr., 5 specimens
<u>Phasganophora capitata</u>	c(98) ¹	Coffman, 1967	riffle areas in Aug & Oct
Perlodidae			
<u>Arcynopteryx curvata</u>	ac(70, 21) ³	Thut, 1969a	87.9% chironomids
<u>A. subtruncata</u>	ac(39, 53) ³	Thut, 1969a	animals were 85.6% chironomids
<u>Isogenus decius</u>	c(100) ²	Minshall, 1965	measured monthly
<u>I. nonus</u>	ac(29, 63) ³	Thut, 1969a	animals were 85.6% chironomids
<u>I. subvarius</u>	c	Minckley, 1963	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹ based on food calories

² based on number of food items

³ based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Plecoptera (cont.)			
Setipalpia (cont.)			
Perlodidae (cont.)			
<u>Isoperla bilineata</u>	ad	Frison, 1935	---
<u>I. clio</u>	dc (18, 82) ²	Minshall, 1965	measured monthly
<u>I. confusa</u>	c	Frison, 1935	---
<u>I. decepta</u>	c	"	---
<u>I. duplicata</u>	c	"	---
<u>I. grammatica</u>	c	Jones, 1950	---
<u>I. minuta</u>	ad	Frison, 1935	---
<u>I. mohri</u>	c	"	---
<u>I. richardsoni</u>	c	"	---
<u>Perla verticalis</u>	c ²	Muttkowski & Smith, 1929	virtually 100% carnivores, 5 specimens, 1 empty

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Plecoptera (cont.)			
Setipalpia (cont.)			
Perlodidae (cont.)			
<u>Perlodes mortoni</u>	c	Jones, 1950	---
<u>Rickera sorpta</u>	c(80) ³	Thut, 1969a	animals were 95.4 chironomids
Trichoptera			
Brachycentridae			
<u>Brachycentrus americanus</u>	a(97) ³	Mecom & Cummins, 1964	sometimes feeds on "drift"
<u>B. nigrosoma</u>	ac	Murphy, 1919	older larvae become carnivorous
<u>B. subnubilis</u>	d	Hanna, 1957	also fed on diatoms, etc.
<u>Brachycentrus</u> sp.	ac(73, 18) ²	Muttkowski & Smith, 1929	---
<u>Micrasema</u> sp.	a(90) ³	Chapman & Demory, 1963	Deer Cr., 36 specimens

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Trichoptera (cont.)			
Brachycentridae (cont.)			
<u>Micrasema</u> sp.	a(75) ³	Chapman & Demory, 1963	Needle Br., 10 specimens
Calamatoceratidae			
<u>Ganonema americanum</u>	d	Lloyd, 1921	---
<u>Heteroplectron</u>			
<u>californicum</u>	d	Grafius, 1974	lab studies
Glossosomatidae			
<u>Agapetus fuscipes</u>	a	Douglas, 1958	---
<u>A. fuscipes</u>	a	Anderson, 1972 (unpublished)	---
<u>Glossosoma boltoni</u>	a	Badcock, 1949	mainly diatoms
<u>G. boltoni</u>	a	Jones, 1950	blue-green algae, 10 specimens
<u>G. boltoni</u>	a	Scott, 1958	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>References</u>	<u>Remarks</u>
Trichoptera (cont.)			
Glossosomatidae (cont.)			
<u>G. intermedium</u>	d(88) ²	Minshall, 1965	---
<u>G. nigrrior</u>	a(98) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>G. nigrrior</u>	a	Cummins, <u>et al</u> 1966	---
<u>Glossosoma</u> sp.	a(95) ³	Chapman & Demory, 1963	26 specimens
Goeridae			
<u>Goera calcarata</u>	a(99) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Silo nigricornis</u>	a	Slack, 1936	10 specimens
Helicopsychidae			
<u>Helicopsyche borealis</u>	ac(67, 31) ¹	Coffman, 1967	riffle areas in Aug & Oct
Hydropsychidae			
<u>Cheumatopsyche</u> sp.	ac(50, 42) ¹	Coffman, 1967	"
<u>Diplectrona modesta</u>	d(86) ²	Minshall, 1965	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Trichoptera (cont.)			
Hydropsychidae (cont.)			
<u>Hydropsyche betteni</u>	c(98) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>H. betteni</u>	a	Minckley, 1963	---
<u>H. borealis</u>	ad	Coffman, Cummins & Wuycheck, 1971	no animal fragments
<u>H. bronta</u>	ac(39, 55) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>H. instabilis</u>	a	Jones, 1950	some detritus
<u>H. slossonae</u>	c(80) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Hydropsyche sp.</u>	ac(54, 42) ²	Muttkowski & Smith, 1929	27 specimens
<u>Hydropsyche sp.</u>	adc	Slack, 1936	10 specimens
<u>Hydropsyche sp.</u>	a	Badcock, 1949	some detritus and animals
Hydroptilidae			
<u>Agraylea multiplicata</u>	a(100) ²	Minckley, 1963	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Trichoptera (cont.)			
Hydroptilidae (cont.)			
<u>A. sexmaculata</u>	a	Barnard, 1971	filamentous algae
<u>Orthotrichia tetensii</u>	a	Lepneva, 1964	---
Lepidostomatidae			
<u>Lepidostoma</u> sp.	d(95) ³	Chapman & Demory, 1963	Deer Cr. & Needle Br. 93 specimens
<u>Lepidostoma</u> sp.	d	Brusven, 1969	fed on dead squawfish
<u>Lepidostoma</u> sp.	d	Grafius, 1974	lab study
Leptoceridae			
<u>Arthripsodes transversus</u>	-	Resh, 1972	bacterial grazers
<u>Arthripsodes</u> sp.	c	Lehmkuhl, 1970	freshwater sponges
<u>Arthripsodes</u> spp.	c	Resh, 1972	freshwater sponges
<u>Leptocerus</u> sp.	ac	Slack, 1936	5 specimens

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>References</u>	<u>Remarks</u>
Trichoptera (cont.)			
Leptoceridae (cont.)			
<u>Oecetis</u> spp.	c	Ross, 1944	---
<u>Setodes</u> sp.	ac	Merrill & Wiggins, 1971	---
Limnephilidae			
<u>Anabolia nervosa</u>	adc	Slack, 1936	18 specimens
<u>A. nervosa</u>	d	Jones, 1950	some algae
<u>Apatania muliebris</u>	a	Elliott, 1971	---
<u>Arctoecia consicia</u>	d	Lloyd, 1921	---
<u>Astenophylax argus</u>	d	"	---
<u>Chilostigma difficilis</u>	d	"	---
<u>Drusus annulatus</u>	a	Gower, 1966	---
<u>Dicosmoecus</u> sp.	d	Brusven, 1967	fed on dead squawfish
<u>Ecclisomyia</u> sp.	ad	Grafius, 1974	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>References</u>	<u>Remarks</u>
Trichoptera (cont.)			
Limnephilidae (cont.)			
<u>Ecclisopteryx guttulata</u>	a	Scott, 1958	---
<u>Frenesia difficilus</u>	d	Flint, 1956	---
<u>F. missa</u>	d	Flint, 1956	---
<u>Glyphotaelius</u> sp.	ad	Slack, 1936	10 specimens
<u>Halesochila taylori</u>	d	Winterbourne, 1971	---
<u>H. taylori</u>	d	Grafius, 1974	lab study
<u>Halesus guttifer</u>	d	Lloyd, 1921	---
<u>H. radiatus</u>	d	Jones, 1950	---
<u>Halesus</u> sp.	ad	Slack, 1936	20 specimens
<u>Limnephilus combinatus</u>	ad	Lloyd, 1921	aquatic macrophytes
<u>L. flavicornis</u>	d	Hanna, 1957	also fed on diatoms, etc.
<u>L. indivisus</u>	ad	Lloyd, 1921	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>References</u>	<u>Remarks</u>
Trichoptera (cont.)			
Limnephilidae (cont.)			
<u>L. lunatus</u>	d	Hanna, 1957	also fed on diatoms, etc.
<u>L. rhombicus</u>	d	Slack, 1936	10 specimens
<u>L. submonilifer</u>	ad	Lloyd, 1921	---
<u>Neophylax autumnus</u>	ad(33, 67) ²	Minshall, 1965	varies with season
<u>Neophylax nacatus</u>	a(100) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>N. concinnus & oligius</u>	ad(28-37, 62-73) ¹	Sedell, 1971	40 specimens
<u>Platyphylax designata</u>			
(= <u>Hesperophylax</u>)	a	Lloyd, 1921	---
<u>Potamophylax stellatus</u>	d	Hanna, 1957	also fed on diatoms, etc.
<u>Psychoglypha</u> sp.	d	Brusven, 1969 ^a	fed on dead squawfish
<u>Pycnopsyche antica</u>	d(91) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>P. guttifer</u>	d	Cummins, 1964	also some algae

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Trichoptera (cont.)			
Limnephilidae (cont.)			
<u>P. guttifer</u>	d	Feldmeth, 1970	lab feeding studies
<u>P. lepida</u>	d	Cummins, 1964	also some algae
<u>P. lepida</u>	d	Feldmeth, 1970	lab feeding studies
<u>P. scabripennis</u>	d	Lloyd, 1921	---
<u>Pycnopsyche</u> sp.	d(90) ²	Minckley, 1963	data presented graphically
<u>Stenophylax latipennis</u>	d	Scott, 1958	---
<u>S. stellatus</u>	a	Badcock, 1949	also some detritus and invertebrates
<u>Stenophylax</u> sp.	a	Slack, 1936	10 specimens
<u>Thremma</u> sp.	ad(64, 36) ²	Muttkowski & Smith, 1929	23 specimens
Molannidae			
<u>Molanna augusta</u>	ac	Slack, 1936	12 specimens

a = algal feeding

d = detrital feeding

c - carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>References</u>	<u>Remarks</u>
Trichoptera (cont.)			
Odontoceridae			
<u>Odontocerum albicorne</u>	ac	Slack, 1936	12 specimens
<u>O. albicorne</u>	ac	Jones, 1949	---
<u>O. albicorne</u>	adc	Scott, 1958	---
<u>Psilotreta indecisa</u>	a(97) ¹	Coffman, 1967	riffle areas in Aug & Oct
Philopotamidae			
<u>Chimarra aterrima</u>	ad(65, 31) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Chimarra marginata</u>	a	Badcock, 1949	---
Phryganeidae			
<u>Agrypnia pagetana</u>	a	Haage, 1970	filamentous algae and <u>Fucus</u>
<u>Banksiola crotchi</u>	ac	Winterbourne, 1971	carnivorous in final instar
<u>Neuronia clathrata</u>	a	Gatjen, 1926	---
<u>N. pardalis</u>	d	Lloyd, 1921	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Trichoptera (cont.)			
Phryganeidae (cont.)			
<u>N. phalaenoides</u>	a	Gatjen, 1926	---
<u>N. postica</u>	d	Lloyd, 1921	---
<u>N. reticulata</u>	a	Gatjen, 1926	---
<u>N. rufierus</u>	ad	Kolenkina, 1951	---
<u>N. stygipes</u>	d	Lloyd, 1921	---
<u>Oligostomis reticulata</u>	ad	Kolenkina, 1951	---
<u>Phryganea grandis</u>	-	Smirnov, 1962	aquatic macrophytes
<u>P. grandis</u>	c	Haage, 1970	---
<u>P. interrupta</u>	ad	Lloyd, 1921	---
<u>P. minor</u>	a	Gatjen, 1926	---
<u>P. obsoleta</u>	a	"	---
<u>P. striata</u>	a	"	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Trichoptera (cont.)			
Phryganeidae (cont.)			
<u>P. striata</u>	ad	Kolenkina, 1951	preferred leaf detritus & <u>Cladophora</u>
<u>P. varia</u>	a	Gatjen, 1926	---
<u>P. vestita</u>	ad	Lloyd, 1921	aquatic macrophytes
Psychomyiidae			
<u>Polycentropus</u>	c	Badcock, 1949	---
<u>flavomaculatus</u>			
<u>P. flavomaculatus</u>	c	Jones, 1950	---
<u>Psychomyia flavida</u>	a(94) ¹	Coffman, 1967	riffle areas in Aug & Oct
Rhyacophilidae			
<u>Rhyacophila arnaudi</u>	c(94) ²	Thut, 1969b	---
<u>R. dorsalis</u>	c	Slack, 1936	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Trichoptera (cont.)			
Rhyacophilidae (cont.)			
<u>R. dorsalis</u>	c	Badcock, 1949	---
<u>R. dorsalis</u>	c	Scott, 1958	---
<u>R. fuscua</u>	c	Lloyd, 1921	also some filamentous algae
<u>R. grandis</u>	c(90) ²	Thut, 1969b	2 specimens
<u>R. parantra</u>	d(96) ²	Minshall, 1965	---
<u>R. vaccua</u>	c(93) ²	Thut, 1969b	---
<u>R. vaefes</u>	adc(39, 21, 36) ²	Thut, 1969b	---
<u>R. vagrita</u>	c(87) ²	"	---
<u>R. verrula</u>	-	"	31% algae, 64% vascular plant (i.e. moss)
<u>R. verrula</u>	a	Smith, 1968	<u>Prasiola</u> and watercress

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Trichoptera (cont.)			
Rhyacophilidae (cont.)			
<u>R. vepulsa</u> (narvae)	c(91) ²	Thut, 1969b	---
<u>Rhyacophila</u> sp.	dc(28, 49) ²	Muttkowski & Smith, 1929	32 specimens
<u>Rhyacophila</u> spp.	c	Ross, 1944	---
<u>Rhyacophila</u> sp.	c	Chapman & Demory, 1963	Deer Cr. & Needle Br., 16 specimens
Sericostomatidae			
<u>Sericosoma personatum</u>	d	Slack, 1936	10 specimens
<u>S. personatum</u>	d	Jones, 1950	---
Diptera			
Blepharoceridae			
	a	Wirth & Stone, 1968	---
Chaoboridae			
<u>Chaoborus nyblasi</u>	c	Dodson, 1970	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Diptera (Cont.)			
Chaoboridae (Cont.)			
<u>C. flavicans</u>	c	Dodson, 1970	---
Ceratopogonidae	c	Kajak & Pieczynski, 1966	---
<u>Palpomyia</u> spp.	d(95) ¹	Coffman, 1967	riffle areas in Aug & Oct
Chironomidae	ad(71, 29) ²	Muttkowski & Smith, 1929	6 specimens
Chironomidae (except			
Tanypodinae)			
Orthoclaadiinae	d(90) ³	Chapman & Demory, 1963	Deer Cr., 316 specimens
(=Hydrobaeninae)			
Orthoclaadiinae	d(85) ³	Chapman & Demory, 1963	Needle Br., 255 specimens
Pelopiinae	c	Kajak & Pieczynski, 1966	---
Tanypodinae			
(=Pelopiinae)			
	c	Badcock, 1949	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Diptera (cont.)			
Chironomidae (cont.)			
<u>Brillia flavifrons</u>	ad(42, 58) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Corynoneura</u> sp.	ad(66, 34) ¹	"	"
Genus nr. <u>Corynoneura</u> sp.	ad(68, 32) ¹	"	"
<u>C. taris</u>	a(77) ¹	"	"
<u>Crictopus exilis</u>	a(94) ¹	"	"
<u>C. sp. nr. junus</u>	a(95) ¹	"	"
undet. <u>Crictopus</u>	a(100) ¹	"	"
<u>Cryptochironomus</u> sp. A	d(94) ¹	"	"
<u>Cryptochoronomus</u> sp. B	a(100) ¹	"	"
<u>Cryptochoronomus</u> sp. C	c(100) ¹	"	"
<u>Cryptochoronomus</u> sp. 3	c(100) ¹	"	"
<u>C. vulneratus</u>	ad(57, 43) ¹	"	"

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Diptera (cont.)			
Chironomidae (cont.)			
<u>Einfeldia synchrona</u>	ad	Danks, 1971	no filter-feeding mechanism observed
<u>Endochiromus dispar</u>	a	Walshe, 1951	filter-feeders
<u>E. tendens</u>	a	"	"
<u>E. albipennis</u>	a	"	"
<u>Eukiefferiella</u> sp. 2	ad(57, 43) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Eukiefferiella</u> sp. nr.			
<u>brevinervis</u>	a(99) ¹	"	"
<u>E. sp. nr. longicalcar</u>	a(100) ¹	"	"
<u>E. sp. nr. sordens</u>	a(81) ¹	"	"
genus nr. <u>Eukiefferiella</u>	a(77) ¹	"	"
<u>Glyptotendipes barbipes</u>	ad	Kimerle, 1969	in waste stabilization lagoon

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Diptera (cont.)			
Chironomidae (cont.)			
<u>G. foliicola</u>	d	Walshe, 1951	possibly live plant tissue
<u>G. gripekoveni</u>	d	"	"
<u>G. pallens</u>	d	"	"
<u>G. viridis</u>	d	"	"
<u>Hydrobaenus</u> sp. nr.			
<u>paradoreus</u>	a(88) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Lauterborniella marmorata</u>			
(= <u>Zavreliella</u>)	ad	Walshe, 1951	---
<u>Limnochironomus modestus</u>	a(87) ¹	Coffman, 1967	riffle areas in Aug & Oct
Genus nr. <u>Metriocnemus</u>			
sp.	a(80) ¹	"	"
<u>Micropsectra</u> sp.	d	Walshe, 1951	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Diptera (cont.)			
Chironomidae (cont.)			
<u>Microtendipes pedellus</u>	d	Walshe, 1951	---
<u>M. pedellus</u>	a(87) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Paraclunio alaskensis</u>	a	Morley & Ring, 1972	intertidal
<u>Pelopia punctipennis</u>	c	Kajak & Stanczykowska, 1968	---
<u>Pentaneura auriensis</u>	c(100) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>P. carnea</u> group sp.	c(100) ¹	"	"
<u>P. melanops</u> group sp.	c(100) ¹	"	"
<u>P. psilosella</u>	c(100) ¹	"	"
<u>P. sp. A</u>	c(100) ¹	"	"
<u>Pentapedilum sordens</u>	a	Walshe, 1951	filter-feeder
<u>Phaenopsectra jucundus</u>	a(87) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>P. obediens</u>	ad(43, 57) ¹	"	"

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Diptera (cont.)			
Chironomidae (cont.)			
<u>Polypedilum fallax</u> group sp.	c(93) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>P. scabraenum</u>	a(85) ¹	"	"
<u>P. sp. nr. sordens</u>	a(83) ¹	"	"
<u>Polypedilum</u> sp.	d	Walshe, 1951	---
<u>Procladius</u> sp. nr. <u>abumbratus</u>	c(100) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>P. choreus</u>	c	Kajak & Stanczykowska, 1968	---
<u>Psectrocladius</u> sp. nr. 4	a(80) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Saundersia clavicornis</u>	a	Morley & Ring, 1972	intertidal
<u>S. marinus</u>	a	"	"
<u>S. pacificus</u>	a	"	"

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Diptera (cont.)			
Chironomidae (cont.)			
<u>Sergentia coracina</u>	d	Walshe, 1951	---
<u>Stempellina bausei</u>	d	"	---
<u>S. sp. nr. johannseni</u>	a(86) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Tanytarsus conversus</u>	a(84) ¹	"	"
<u>T. dissimilis</u>	ad	Cavanaugh & Tilden, 1930	"
<u>T. sp. nr. exigua</u>	a(74) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>T. exigua</u>	ad(66, 27) ¹	"	"
<u>T. sp. nr. polita</u>	a(80) ¹	"	"
<u>T. sp. 3</u>	a(87) ¹	"	"
<u>T. sp. 4</u>	ad(47, 53) ¹	"	"
<u>T. sp. 5</u>	a(100) ¹	"	"
<u>T. rivulorum</u>			
(=Rheotanytarsus)	a	Walshe, 1951	filter-feeders

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Diptera (cont.)			
Chironomidae (cont.)			
<u>Tanytarsus</u> sp.	ad	Walshe, 1951	---
<u>Tendipes anthracinus</u>			
(= <u>Chironomus</u>)	d	"	---
<u>Tendipes dissidens</u>			
(= <u>Einfeldia</u>)	d	"	---
<u>T. dorsalis dorsalis</u>	d	"	---
<u>T. riparius</u>	d	"	---
<u>T. plumosus flaveolus</u>	a	"	filter-feeders
<u>T. plumosus plumosus</u>	a	"	"
<u>Trichocladius</u> sp. nr. 3	c(77) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Zavrelia</u> sp. nr.			
<u>pentatoma</u>	ad(61, 27) ¹	"	"

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Diptera (cont.)			
Dixidae			
<u>Dixa</u> sp.	d(95) ³	Chapman & Demory, 1963	Deer Cr., 4 specimens
Empididae			
<u>Hemerodromia</u> sp.	ad(34, 66) ¹	Coffman, 1967	riffle areas in Aug & Oct
Psychodidae			
<u>Psychoda alternata</u>	ad(39, 61) ¹	"	"
Simuliidae			
<u>Prosimulium caudatum</u>	d(87) ³	Speir, 1973 (unpublished)	30 specimens, Feb & Mar
<u>P. dicum</u>	d(84) ³	"	26 specimens, Feb & Mar
<u>Simulium arcticum</u>	-	Fredeen, 1963	reared on bacterial suspension
<u>S. arcticum</u>	ad(39, 58) ³	Speir, 1973 (unpublished)	34 specimens, May

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Diptera (cont.)			
Simuliidae (cont.)			
<u>S. canadense</u>	ad(33, 62) ³	Speir, 1973 (unpublished)	18 specimens, May
<u>S. venustum</u>	-	Fredeen, 1963	reared on bacterial suspension
<u>S. verecundum</u>	-	"	"
<u>S. vittatum</u>	d(88) ³	Speir, 1973 (unpublished)	14 specimens, Oct
<u>S. vittatum</u>	d	Fredeen, 1963	reared on bacterial suspension
<u>Simulium</u> spp.	a (79) ²	Muttkowski & Smith, 1929	---
<u>Simulium</u> sp.	d(95) ³	Chapman & Demory, 1963	33 specimens
<u>Simulium</u> sp.	a(90) ³	Coffman, 1967	riffle areas in Aug & Oct
<u>Twinnia nova</u>	a(81) ³	Speir, 1973 (unpublished)	9 specimens, Mar, this species has no head fans

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Diptera (cont.)			
Tabanidae			
<u>Chrysops</u> spp.	c	Philip, 1931	lab studies
<u>Tabanus</u> sp.	ad(74, 26) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Tabanus</u> sp.	c	Philip, 1931	lab studies
Tipulidae			
<u>Antocha</u> sp.	a(93) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Dicranota</u> sp.	d(78) ¹	"	"
<u>Eriocera fultonensis</u>	c(96) ¹	"	"
<u>Tipula nobilis</u>	a(90) ²	Minckley, 1963	data presented graphically
<u>Tipula</u> sp.	c(76) ¹	Coffman, 1967	riffle areas in Aug & Oct
Megaloptera			
Corydalidae			
<u>Chauliodes</u> sp.	c	Chandler, 1968	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Megaloptera (cont.)			
Corydalidae (cont.)			
<u>Nigronia serricornis</u>	dc(85, 15) ³	Minshall, 1965	---
Sialidae			
<u>Sialis californica</u>	c	Azam, 1968	---
<u>S. joppa</u>	c(100) ³	Minshall, 1965	---
<u>S. rotunda</u>	c	Azam, 1968	---
<u>Sialis</u> spp.	c	Ross, 1937	---
Coleoptera			
Dryopidae			
	-	Leech & Chandler, 1968	larvae are probably root feeders
<u>Helicus</u> sp. (adults)	ad(26, 74) ¹	Coffman, 1967	riffle areas in Aug & Oct
Dytiscidae			
<u>Acilius</u> sp.	c	Balduf, 1935	adults & larvae all carnivorous

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Coleoptera			
Dytiscidae			
<u>Colymbetes</u> sp.	c	Balduf, 1935	adults & larvae all carnivorous
<u>Coptotomus</u> sp.	c	"	"
<u>Cybister</u> spp.	c	"	"
<u>Dytiscus</u> spp.	c	"	"
<u>Limnaea</u> sp.	c	"	"
<u>Hydroporus</u> sp.	c	"	"
Elmidae	ad	Badcock, 1949	---
<u>Dubiraphia</u> sp. (adults)	ad(26, 74) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Optioservus</u> <u>quadrimaculatus</u> (adults)	ad(65, 35) ³	Chapman & Demory, 1963	Deer Cr., 6 specimens

a = algal feeding

d = detrital feeding

c = carnivorous

¹based on food calories

²based on number of food items

³based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Coleoptera (cont.)			
Elmidae (cont.)			
<u>O. quadrimaculatus</u>			
(adults)	d(100) ³	Chapman & Demory, 1963	Needle Br., 1 specimen
<u>Optioservus</u> sp. (larvae)	a(90) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Optioservus</u> sp. (adults)	a(76) ¹	"	"
<u>Stenelmis beameri</u>			
(larvae)	ac(71, 22) ¹	"	"
<u>S. beameri</u> (adults)	ad(59, 41) ¹	"	"
Hydrophilidae			
all larvae except			
<u>Berosus</u> spp.	c	Balduf, 1935	---
<u>Berosus</u> spp.	a	"	adults and larvae
<u>Enochrus</u> sp.	a	"	adults

a = algal feeding

d = detrital feeding

c = carnivorous

¹ based on food calories

² based on number of food items

³ based on food volume or area

<u>Taxa</u>	<u>Food Categories and % Composition</u>	<u>Reference</u>	<u>Remarks</u>
Coleoptera (cont.)			
Hydrophilidae (cont.)			
<u>Laccobius</u> sp.	a	Balduf, 1935	adults
<u>Hydrophilus obtusatus</u>	a	"	"
<u>Tropisternis</u> sp.	a	"	"
<u>Hydrous triangularis</u> (= <u>Hydrophilus</u>)	-	Wilson, 1923	adults may be predaceous
<u>Hydrophilus piceus</u>	c	Miller, 1963	adults & larvae
<u>Tropisternis</u> sp.	ad	Spangler, 1960	adults
Psephenidae			
<u>Ectoporia</u> sp.	a(98) ¹	Coffman, 1967	riffle areas in Aug & Oct
<u>Psephenus herricki</u>	a(98) ¹	Coffman, 1967	"
<u>P. herricki</u>	a	Murvosh, 1971	---

a = algal feeding

d = detrital feeding

c = carnivorous

¹ based on food calories

² based on number of food items

³ based on food volume or area

References

- Anderson, N. H. 1972. Unpublished data. Entomology Department, Oregon State Univ., Corvallis, Ore.
- Azam, K. M. 1968. Life history and production studies of Sialis californica Banks and Sialis rotunda Banks (Megaloptera: Sialidae). Ph. D. Thesis. Oregon State Univ., Corvallis, Ore. 111 pp.
- Badcock, R. M. 1949. Studies on stream life in tributaries of the Welch Dee. *J. Anim. Ecol.* 18:193-208.
- Barnard, P. C. 1971. The larva of Agraylea sexmaculata Curtis (Trichoptera: Hydroptilidae). *Ent. Gazette.* 22:253-257.
- Brown, D. S. 1960. The ingestion and digestion of algae by Cloeon dipterum L. (Ephemeroptera). *Hydrobiol.* 16:81-96.
- Brown, D. S. 1961. The food of the larvae of Cloeon dipterum L. and Baetis rhodani (Pictet) (Ephemeroptera). *J. Anim. Ecol.* 30:55-75.
- Brusven, M. A. and A. C. Scoggan. 1969. Sarcophagous habits of Trichoptera larvae on dead fish. *Ent. News.* 80:103-105.
- Burks, B. D. 1953. The mayflies, or Ephemeroptera, of Illinois. *Bull. Ill. St. Nat. Hist. Surv.* 26:1-216.
- Cavanaugh, W. J. and J. E. Tilden. 1930. Algal food, feeding, and case building habits of the larvae of the midge fly, Tanytarsus dissimilis. *Ecology* 11:281-287.
- Chandler, H. P. 1968. Megaloptera. pp. 229-233. In Usinger, R. L. (ed.). *Aquatic Insects of California*. University of California Press. Berkeley and Los Angeles. 508 pp.

- Chapman, D. W. and R. Demory. 1963. Seasonal changes in the food ingested by aquatic insect larvae and nymphs in two Oregon streams. *Ecology* 44:140-146.
- Clemens, W. A. 1917. An ecological study of the mayfly Chironetes. Univ. Toronto Stud. Biol. 17.
- Coffman, W. P. 1967. Community structure and trophic relations in a small woodland stream, Linesville Creek, Crawford County, Pennsylvania. Ph. D. thesis. University of Pittsburgh. 573 pp.
- Coffman, W. P., K. W. Cummins, and J. C. Wuycheck. 1971. Energy flow in a woodland stream ecosystem: I. Tissue support trophic structure of the autumnal community. *Arch. Hydrobiol.* 68:232-276.
- Cummins, K. W. 1964. Factors limiting the microdistribution of larvae of the caddisflies Pycnopsyche lepida (Hagen) and Pycnopsyche guttifer (Walker) in a Michigan stream. *Ecol. Monogr.* 34:271-295.
- Cummins, K. W., W. P. Coffman, and P. A. Roff. 1966. V. Running waters. Trophic relationships in a small woodland stream. *Verh. Internat. Verein. Limnol.* 16:627-638.
- Danks, H. V. 1971. Life history and biology of Einfeldia synchrona (Diptera: Chironomidae). *Can. Ent.* 103:1597-1606.
- Dodson, S. I. 1970. Complementary feeding niches sustained by size-selective predation. *Limnol. Oceanogr.* 15:131-137.
- Edmunds, G. F. Jr. 1957. The predaceous mayfly nymphs of North America. *Proc. Utah Acad. Sci. Arts and Letters* 34:23-24.
- Edmunds, G. F. Jr. 1960. The food habits of the nymph of the mayfly Siphonurus occidentalis. *Proc. Utah Acad. Sci. Arts and Letters.* 37:73-74.

- Elliott, J. M. 1971. The life history and biology of Apatania muliebris Mclachlan (Trichoptera). Ent. Gazette. 22:245-251.
- Ellis, R. J. 1970. Alloperla stonefly nymphs: Predators or scavengers on salmon eggs and alevins? Trans. Am. Fish. Soc. 99:677-683.
- Feldmeth, C. R. 1970. The respiratory energetics of two species of stream caddisfly larvae in relation to water flow. Comp. Biochem. Physiol. 32:193-202.
- Flint, O. S. Jr. 1956. The life and biology of the genus Frenesia (Trichoptera: Limnephilidae). Bull. Brooklyn Ent. Soc. 51:93-108.
- Fredeen, F. H. J. 1963. Bacteria as a source of food for black-fly larvae (Diptera: Simuliidae) in laboratory cultures and in natural streams. Can. J. Zool. 42:527-538.
- Fremling, C. R. 1960. Biology of a large mayfly, Hexagenia bilineata (Say), of the upper Mississippi River. Res. Bull. Iowa Ag. Exp. Stn. 482:842-852.
- Frison, T. H. 1929. Fall and winter stoneflies, or Plecoptera, of Illinois. Bull. Ill. St. Nat. Hist. Surv. 18:345-409.
- Frison, T. H. 1935. The stoneflies, or Plecoptera, of Illinois. Bull. Ill. Nat. Hist. Surv. 20:281-471.
- Gatjen, L. 1926. Nahrungsuntersuchung bei Phryganidenlarven (Phrygania und Neuronia). Arch. Hydrobiol. 16:649-667.
- Gilpin, B. R. and M. A. Brusven. 1970. Food habits and ecology of mayflies of the St. Maries River in Idaho. Melanderia 4:19-40.
- Gower, A. M. 1966. The life cycle of Drusus annulatus Steph. (Trichoptera: Limnephilidae) in watercress beds. Ent. mon. Mag. 101:113-141.
- Grafius, E. J. 1973. The conversion of allochthonous material by stream detritivores. M.S. thesis. Entomology Dept., Oregon State Univ., Corvallis, Ore.

- Haage, P. 1970. On the feeding habits of two Baltic species of caddis larvae (Trichoptera). *Entomol. Scand.* 1: 282-290.
- Hanna, H. M. 1957. A study of the growth and feeding habits of the larvae of four species of caddisflies. *Proc. R. ent. Soc. Lond. (A)* 32:139-146.
- Hunt, B. P. 1953. The life history and economic importance of a burrowing mayfly, Hexagenia limbata, in Southern Michigan lakes. *Bull. Inst. Fish. Res.* 4:151 pp.
- Hynes, H. B. N. 1941. The taxonomy and ecology of the nymphs of British Plecoptera, with notes on the adults and eggs. *Trans. R. ent Soc. Lond.* 91:459-557.
- Hynes, H. B. N. 1970. The ecology of running waters. Univ. Toronto Press. 555 pp.
- Hynes, H. B. N. and N. K. Kaushik. 1968. Experimental study of the role of autumnshed leaves in aquatic environments. *J. Ecol.* 56:229-243.
- Ivanova, S. S. 1958. Nutrition of some mayfly larvae. *Proc. Mikoyan Moscow Tech. Inst. Fishing Industry.* 9:102-120. (Russian) In Brown, D. S. 1960. The ingestion and digestion of algae by Cloeon dipterum L. (Ephemeroptera). *Hydrobiol.* 16:81-96.
- Jones, J. R. E. 1949. An ecological study of the River Rheidal, North Cardiganshire, Wales. *J. Anim. Ecol.* 18:67-88.
- Jones, J. R. E. 1950. A further ecological study of the River Rheidal: The food of the common insects of the main-stream. *J. Anim. Ecol.* 19:159-174.
- Kajak, Z. and E. Pieczynski. 1966. Wplyw drapiezcow bezkregowych na liczebnośc organizmow bentosowych (glownie Chironomidae). *Ekologia Polska. seria B.* 12:165-179 (English summary).

- Kajak, Z. and A. Stanczykowska. 1968. Influence of mutual relations of organisms, especially Chironomidae, in natural benthic communities, on their abundance. *Ann. Zool. Fenn.* 5:49-56.
- Kolenkina, L. V. 1951. Nutrition of the larvae of some caddis flies (Trichoptera). *Trudy Vsesoyuznogo Gidrobiologicheskogo Obshchestva.* 3:44-57. (Russian) In Lepneva, S. G. 1964. Fauna of the U.S.S.R.: Trichoptera. Vol. II(1). Translated by the Israel Program for Scientific Translations. 1970.
- Kimerle, R. A. 1969. Production biology of Glyptotendipes barbipes (Staeger) (Diptera:Chironomidae) in a waste stabilization lagoon. Ph. D. thesis. Oregon State Univ., Corvallis. 125 pp.
- Leathers, A. L. 1922. Ecological study of the aquatic midges and some related insects with special reference to feeding habits. *Bull. U. S. Bur. Fish.* 38:1-61.
- Leech, H. B. and H. P. Chandler. 1968. Aquatic Coleoptera. pp. 293-371. In Usinger, R. L. (ed.). *Aquatic Insects of California.* University of California Press. Berkeley and Los Angeles. 508 pp.
- Lehmkuhl, D. M. 1970. A North American trichopteran larva which feeds on freshwater sponges (Trichoptera:Leptoceridae;Porifera:Spongillidae). *Amer. Midl. Nat.* 84:278-280.
- Lehmkuhl, D. M. 1972. Baetisca (Ephemeroptera:Baetiscidae) from the western interior of Canada with notes on the life cycle. *Can. J. Zool.* 50:1017-1019.
- Lepneva, S. G. 1964. Fauna of the U. S. S. R.: Trichoptera. Vol II(1). Translated by the Israel Program for Scientific Translations. 1970.

- Lloyd, J. T. 1921. The biology of North American caddis fly larvae.
Bull. Lloyd Libr. Ent. Ser. 1:124 pp.
- McDiffett, W. F. 1970. The transformation of energy by a stream
detritivore, Pteronarcys scotti (Plecoptera). Ecology 5:975-988.
- Mecom, J. O. and K. W. Cummins. 1964. A preliminary study of the trophic
relationships of the larvae of Brachycentrus americanus (Banks)
(Trichoptera:Brachycentridae). Trans. Am. Microscop. Soc. 83:233-243.
- Merrill, D. and G. B. Wiggins. 1971. The larvae and pupa of the caddisfly
genus Setodes in North America (Trichoptera:Leptoceridae). Life Sci.
Occ. Pap., R. Ont. Mus. 19:1-12.
- Minckley, W. L. 1963. The ecology of a spring stream, Doe Run, Meade
County, Kentucky. Wildl. Monogr. 11:1-124.
- Minshall, G. W. 1965. Community dynamics and economics of a woodland
springbrook: Morgan's Creek, Meade County, Kentucky. Ph. D. thesis,
Department of Biology, Univ. of Louisville. 261 pp.
- Minshall, G. W. 1967. Role of allochthonous detritus in the trophic
structure of a woodland springbrook community. Ecology 48:139-144.
- Minshall, J. N. 1964. An ecological life history of Epeorus pleuralis
(Banks) in Morgan's Creek, Meade County, Kentucky. M. S. thesis.
Univ. of Louisville. 79 pp.
- Moon, H. P. 1938. The growth of Caenis horaria, Leptophlebia marginata
and L. vespertina. Proc. zool. Soc. Lond. (A) 108:507-512.
- Morgan, A. H. 1911. Mayflies of Fall Creek. Ann. Ent. Soc. Amer. 4:93-119.
- Morley, R. L. and R. A. Ring. 1972. The intertidal Chironomidae (Diptera)
of British Columbia II. Life history and population dynamics. Can.
Ent. 104:1099-1121.

- Murphy, H. 1919. Observations on the egg laying of the caddisfly Brachycentrus nigrosoma Banks, and the habits of the young larvae. J. New York Ent. Soc. 27:154-158.
- Muttkowski, R. A. and G. M. Smith. 1929. The food of trout stream insects in Yellowstone National Park. Roosevelt Wildlife Ann. 2:241-263.
- Needham, J. H. and P. W. Claassen. 1925. A monograph of the Plecoptera or stoneflies of America north of Mexico. Thomas Say Foundation. Ent. Soc. Amer. 2:1-397.
- Philip, C. B. 1931. The Tabanidae (horseflies) of Minnesota. With special reference to their biologies and taxonomy. Univ. Minn. Agric. Exp. Sta. Tech. Bull. 80:128 pp.
- Pinet, J. M. 1962. Observations sur la biologie et l'ecologie d'Oligoneuriella rhenana Imhoff (Ephemeroptera). Annl. Stn. Cent. Hydrobiol. 9:301-331. In Hynes, H. B. N. 1970. Ecology of running waters. Univ. Toronto Press. 555 pp.
- Ross, H. N. 1944. The caddis flies, or Trichoptera, of Illinois. Bull. Ill. Nat. Hist. Surv. 23:326 pp.
- Ross, H. H. 1937. Neartic alder flies of the genus Sialis (Megaloptera, Sialidae). Ill. Nat. Hist. Surv. 21:57-78.
- Scott, D. 1958. Ecological studies on the Trichoptera of the River Dean, Cheshire. Arch. Hydrobiol. 54:340-392.
- Sedell, J. R. 1971. Trophic ecology and natural history of Neophylax concinnus and N. oligius (Trichoptera: Limnephilidae). Ph. D. thesis. Univ. Pittsburgh. 154 pp.
- Slack, H. D. 1936. The food caddisfly larvae (Trichoptera). J. Anim. Ecol. 5:105-115.

- Smirnov, N. N. 1962. On nutrition of caddis worms Phryganea grandis L. Hydrobiol. 19:252-261.
- Snow, S. P. 1973. Unpublished data. Entomology Department, Oregon State Univ., Corvallis, Ore.
- Speir, J. A. 1973. Unpublished data. Entomology Department, Oregon State Univ., Corvallis, Ore.
- Strenger, A. 1953. Zur Kopfmorphologie der Ephemeridenlarven. Erster Teil. Ost. Zool. Z. 4:191-228. In Hynes, H. B. N. 1970. Ecology of running waters. Univ. Toronto Press. 555 pp.
- Thut, R. N. 1969a. Feeding habits of stonefly nymphs of the suborder Setipalpia. Unpublished Paper presented at Midwest Benthological Society Annual Meeting.
- Thut, R. N. 1969b. Feeding habits of larvae of seven Rhyacophila (Trichoptera). Ann. Ent. Soc. America. 62:894-898.
- Wallace, J. B., W. R. Woodall, and F. F. Sherberger. 1970. Breakdown of leaves by feeding of Peltoperla maria nymphs (Plecoptera:Peltoperlidae). Ann. Ent. Soc. of America. 63:562-567.
- Walshe, B. M. 1951. The feeding habits of certain chironomid larvae (subfamily Tendipedinae). Proc. zool. Soc. Lond. 121:63-79.
- Winterbourne, M. J. 1971. The life histories and trophic relationships of the Trichoptera of Marion Lake, British Columbia. Can. J. Zool. 49:623-635.
- Wirth, W. W. and A. Stone. 1968. Aquatid Diptera pp. 372-482. In Usinger, R. L. (ed.). Aquatic Insects of California, University of California Press. Berkeley and Los Angeles. 508 pp.
- Wissmeyer, A. 1926. Nahrungsuntersuchungen bei Ephemeridenlarven. Arch. Hydrobiol. 16:668-698.
- Wu, C. F. 1923. Morphology, anatomy and ethology of Nemoura. Lloyd Bull. Libr. 23:1-81