

Carbon budget estimation of a subarctic catchment using a dynamic ecosystem model at high spatial resolution

The Faculty of Oregon State University has made this article openly available.
Please share how this access benefits you. Your story matters.

Citation	Tang, J., Miller, P. A., Persson, A., Olefeldt, D., Pilesjö, P., Heliasz, M., ... & Christensen, T. R. (2015). Carbon budget estimation of a subarctic catchment using a dynamic ecosystem model at high spatial resolution. <i>Biogeosciences</i> , 12, 933-980. doi:10.5194/bgd-12-933-2015
DOI	10.5194/bgd-12-933-2015
Publisher	Copernicus Publications
Version	Version of Record
Terms of Use	http://cdss.library.oregonstate.edu/sa-termsfuse

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Carbon budget estimation of a subarctic catchment using a dynamic ecosystem model at high spatial resolution

J. Tang^{1,2}, P. A. Miller¹, A. Persson¹, D. Olefeldt³, P. Pilesjö¹, M. Heliasz¹,
M. Jackowicz-Korczynski¹, Z. Yang⁴, B. Smith¹, T. V. Callaghan^{5,6,7}, and
T. R. Christensen^{1,8}

¹Department of Physical Geography and Ecosystem Science, Lund University, Sölvegatan 12, 223 62, Lund, Sweden

²Department of Resource and Environmental Science, Wuhan University, Road Luoyu 129, Wuhan, China

³Department of Renewable Resources, University of Alberta, Edmonton, AB T6G 2H1, Canada

⁴Department of Forest Ecosystems and Society, Oregon State University, Corvallis, 973 31, Oregon, USA

⁵Royal Swedish Academy of Sciences, Lilla Frescativägen 4A, 114 18, Stockholm, Sweden

⁶Department of Animal and Plant Sciences, University of Sheffield, S10 2TN Sheffield, UK

⁷Department of Botany, National Research Tomsk State University, 36, Lenin Ave., Tomsk, 634050, Russia

⁸Arctic Research Centre, Aarhus University, C.F. Møllers Allé 8, 8000 Aarhus C, Denmark

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Received: 10 October 2014 – Accepted: 20 November 2014 – Published: 16 January 2015

Correspondence to: J. Tang (lu.gistangjing@gmail.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

12, 933–980, 2015

**C budget estimation
of a subarctic
catchment**

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Large amount of organic carbon is stored in high latitude soils. A substantial proportion of this carbon stock is vulnerable and may decompose rapidly due to temperature increases that are already greater than the global average. It is therefore crucial to quantify and understand carbon exchange between the atmosphere and subarctic/arctic ecosystems. In this paper, we combine an arctic-enabled version of the process-based dynamic ecosystem model, LPJ-GUESS (version LPJG-WHyMe-TFM) with comprehensive observations of terrestrial and aquatic carbon fluxes to simulate long-term carbon exchange in a subarctic catchment comprising both mineral and peatland soils. The model is applied at 50m resolution and is shown to be able to capture the seasonality and magnitudes of observed fluxes at this fine scale. The modelled magnitudes of CO₂ uptake generally follow the descending sequence: birch forest, non-permafrost *Eriophorum*, *Sphagnum* and then tundra heath during the observation periods. The catchment-level carbon fluxes from aquatic systems are dominated by CO₂ emissions from streams. Integrated across the whole catchment, we estimate that the area is a carbon sink at present, and will become an even stronger carbon sink by 2080, which is mainly a result of a projected densification of birch forest and its encroachment into tundra heath. However, the magnitudes of the modelled sinks are very dependent on future atmospheric CO₂ concentrations. Furthermore, comparisons of global warming potentials between two simulations with and without CO₂ increase since 1960 reveal that the increased methane emission from the peatland could double the warming effects of the whole catchment by 2080 in the absence of CO₂ fertilization of the vegetation. This is the first process-based model study of the temporal evolution of a catchment-level carbon budget at high spatial resolution, integrating comprehensive and diverse fluxes including both terrestrial and aquatic carbon. Though this study also highlights some limitations in modelling subarctic ecosystem responses to climate change including aquatic system flux dynamics, nutrient limitation, herbivory and other disturbances and peatland

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



C budget estimation of a subarctic catchment

J. Tang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Roulet et al., 2007; Christensen et al., 2012; McGuire et al., 2012). Year-around observations are needed because there is clear evidence that C fluxes in the cold seasons are very important (Larsen et al., 2007b; Mastepanov et al., 2008).

2. Observations of interactions between terrestrial and aquatic systems are lacking (Lundin et al., 2013; Olefeldt et al., 2013). Quantifications of terrestrial lateral loss of C are needed not only because they represent a significant fraction of net ecosystem exchange but also because they are intrinsically linked to downstream aquatic C cycling (Lundin et al., 2013). Integrating terrestrial and aquatic C cycling is of high importance for our understanding of the C balance at the catchment scale, particularly at high latitudes. Northern peatlands are large sources of dissolved organic carbon (DOC), while receiving lakes generally are net sources of both CO₂ and CH₄ (Tranvik et al., 2009).

Many environmental characteristics of the Stordalen catchment, located in the subarctic discontinuous permafrost region of northern Sweden, have been measured since the 1950s (Bäckstrand, 2008) and many studies covering a variety of disciplines are still on-going (Callaghan et al., 2010, 2013). Observations related to the dynamics of almost all the relevant C components have been made in different landcover types (Christensen et al., 2012; Callaghan et al., 2013). However, the various observations over different landcover types have not yet been integrated into a comprehensive, year-round catchment-level C budget. A significant aspect of this area is that it contains permafrost that is rapidly thawing (Åkerman and Johansson, 2008). This makes more C hydrologically available (Olefeldt and Roulet, 2012) and the large stock of C in these tundra heath soils becomes available to microbial processes (Sjögersten and Wookey, 2002; Fox et al., 2008; Hartley et al., 2012). A rapid changing environment together with comprehensive observations have established the unique importance of this area as a model system for furthering our process-based understanding of the role of climate changes in northern regions. Furthermore, this understanding, gained in an accessible

and highly instrumented area, can be applied to vast areas where large C stocks exist but long term measurements are lacking.

The Stordalen catchment contains several distinct landcover types, including tundra heath, birch forest with heath understory, peatlands and lakes/rivers. Hereafter, the peatland is divided into three groups named “palsa”, “*Sphagnum* site” and “*Eriophorum* site”, based on surface hydrology, permafrost condition as well as characteristic plant communities. An earlier compilation of the C balance of the larger Torneträsk catchment, which encompasses the Stordalen catchment, indicated that there was a significant sink capacity in the birch forest as well as across the peatland (Christensen et al., 2007). This assessment, however, lacked year-around measurement of CO₂ and CH₄ emissions and did not include direct measurements of aquatic C fluxes (Christensen et al., 2007). Subsequent observations in the Stordalen catchment have focused on filling the missing components. Consequently, recently updated year-around CO₂ and CH₄ measurements in the peatland identified the wetter non-permafrost *Eriophorum* sites to be strong C sinks ($-46 \text{ gC m}^{-2} \text{ yr}^{-1}$) with high CH₄ emissions ($18\text{--}22 \text{ gC m}^{-2} \text{ yr}^{-1}$) (Jackowicz-Korczyński et al., 2010; Christensen et al., 2012), while measurements conducted on the drier palsa (where permafrost is present) showed a relatively weaker uptake ($-39.44 \text{ gC m}^{-2} \text{ yr}^{-1}$) (Olefeldt et al., 2012). Total waterborne C exports (DOC plus particulate organic C (POC) and dissolved inorganic C (DIC)) from the terrestrial ecosystems (both peatland and forest) were also monitored (Olefeldt et al., 2013) and found to represent a significant component of the net ecosystem C balance, ranging from 2.77 to 7.31 $\text{gC m}^{-2} \text{ yr}^{-1}$. In contrast, four years of continuous Eddy-Covariance (EC)-tower based CO₂ measurements in the birch forest revealed very variable C sink/source functionality, which in two out of four years has been found to be a C source to the atmosphere (Heliasz, 2012). Tundra heath, another important landcover type in this region, has lower C uptake (Christensen et al., 2007; Fox et al., 2008) and both birch forest and tundra heath were found to have high spatial heterogeneity (Fox et al., 2008; Heliasz, 2012). Altogether, different landcover types show diverse contributions to this subarctic ecosystem’s C balance.

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



With pronounced future warming expected in this region, the structure and function of the different vegetation types are expected to vary dramatically as has been observed during warming in the recent past (Callaghan et al., 2013). In addition, changes to soil conditions due to warming and permafrost thaw will likely stimulate C fluxes to the atmosphere and affect the long-term accumulated C (Wolf et al., 2008a; McGuire et al., 2012), but this likely C release needs to be weighed against the possibility of increased uptake by increased primary productivity resulting from longer growing seasons and/or potential CO₂ fertilization.

The field measurements described above provide an insight into the ongoing processes and current ecosystem status, but until now, no modeling exercises have been implemented in this region in combination with the comprehensive measured data. Moreover, high spatial resolution predictions of future potential dynamics of both vegetation and soil processes and their responses to the projected climate are lacking in this region. In this study, therefore, we aim to assess the Stordalen catchment C budget in a retrospective as well as in a prognostic way by implementing a process-based dynamic ecosystem model (Smith et al., 2001; Miller and Smith, 2012) integrated with a distributed hydrology model (Tang et al., 2014a, b) at high spatial resolution (50 m by 50 m, (Yang et al., 2012)). We quantify the overall C budget of the study catchment by synthesizing diverse C fluxes and specifically address the following questions: (1) will this subarctic catchment become a C source or a larger C sink in the near future? (2) How differently will the catchment's vegetation micro-types respond to the climate drivers? And (3) what are the major limitations in the model's prognostic ability?

To answer these questions, we implemented an arctic-enabled version of the dynamic ecosystem model, LPJ-GUESS WHyMe (Wania et al., 2009a, b, 2010). This model has been widely and successfully implemented for estimating and predicting ecosystem function in high-latitude regions (McGuire et al., 2012; Miller and Smith, 2012; Zhang et al., 2013). LPJ-GUESS WHyMe includes comprehensive process descriptions to capture the interactions between atmosphere–vegetation–soil domains

and it explicitly describes permafrost and peatland processes, which are important components of our study catchment. Importantly, previous studies (Tang et al., 2014a, b) dedicated to integrate a distributed hydrology scheme to LPJ-GUESS WHyMe have demonstrated the necessity of considering lateral water movements to accurately capture water and C cycling in this region. The model with distributed hydrology is called LPJG-WHyMe-TFM where TFM stands for “Triangle Form-based Multiple flow algorithm” (Pilesjö and Hasan, 2014). The study presented here is the first modeling exercise to combine all the available year-around measured data in the Stordalen catchment and at high spatial resolution.

2 Model descriptions

The process-based dynamic ecosystem model, LPJG-WHyMe-TFM, was chosen as the platform for studying the subarctic catchment C balance. The processes in the model include vegetation growth, establishment and mortality, disturbance, competition between plant individuals for light and soil water (Smith et al., 2001, 2014) and soil biogeochemical processes (Sitch et al., 2003). These processes are operated in a number of independent and replicate patches. Vegetation in the model is defined and grouped by plant function types (PFTs), which are based on plant phenological and physiological features combined with bioclimatic limits (Hickler et al., 2004; Wramneby et al., 2008). Bioclimatic conditions determine which PFTs can potentially grow in study regions, and vertical stand structure together with soil water availability further influence PFTs establishment based on PFTs’ shade tolerance and drought tolerance characteristics. A list of the simulated PFTs in this catchment can be found in Table S1 in the Supplement. The parameterization and choice of non-peatland PFTs in this study cover all the main vegetation types found in the study region (boreal forest, shrubs, open-ground grasses, peatland mosses and flood-tolerant C3 graminoids) and are based on previous studies for the arctic region using LPJ-GUESS (Wolf et al., 2008a; Hickler et al., 2012; Miller and Smith, 2012; Zhang et al., 2013). For peatland

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



grid cells, two new peatland PFTs, flood-tolerant graminoids and *Sphagnum* moss, are introduced by Wania et al. (2009b).

The model is driven by climate data and includes both non-peatland and peatland hydrological processes (Fig. 1). The vertical water movement between atmosphere, vegetation and soil is based on Gerten et al. (2004) while the lateral water movement between grid cells was implemented by Tang et al. (2014b) based on topographical variations. More recently, an advanced multiple flow algorithm TFM (Pilesjö and Hasan, 2014) was chosen to distribute water among grid cells, due to its better treatment of flow continuity and flow estimation over flat surfaces (Tang et al., 2014a). Within the catchment boundary, surface and subsurface runoff can move from one upslope cell to multiple downslope cells, which greatly improves hydrological flux estimations (Tang et al., 2014a) and results in a better estimate of C fluxes in peatland region. The soil temperature estimation is driven by surface air temperature, and the Crank–Nicolson heat diffusion algorithm (Crank and Nicolson, 1996) is implemented to calculate the soil temperature profile on a daily time-step (Wania et al., 2009a). The C cycling descriptions in LPJG-WHyMe-TFM for peatland cells are based on Wania’s developments (Wania et al., 2009a, b, 2010), while the non-peatland grid cell C cycling is kept the same as in LPJ-GUESS (Smith et al., 2001; Sitch et al., 2003). The full hydrological processes and peatland C cycling descriptions in LPJG-WHyMe-TFM can be found in Sect. S1 in the Supplement. A brief summary of relevant C cycling processes in the model will be presented below.

A modified Farquhar photosynthesis scheme (Haxeltine and Prentice, 1996; Haxeltine et al., 1996; Sitch et al., 2003) is used to estimate gross primary production (GPP), which is related to air temperature (T), atmospheric CO_2 concentration, absorbed photosynthetically active radiation (PAR) and stomatal conductance. Part of the GPP is respired to the atmosphere by maintenance and growth respiration (R_a), and the remaining part is net primary production (NPP) for each PFT. The reproduction costs are subtracted from NPP, and thereafter the remaining NPP is allocated to different living tissues in accordance with a set of PFT-specific allometric relationships

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Smith et al., 2001). Leaves, fine root biomass and root exudates are transferred to the litter pool with a given turnover rate, and above-ground plant materials can also provide inputs to the litter pool due to stochastic natural disturbance events and mortality (Smith et al., 2001; Thonicke et al., 2001). The majority (70%) of the litter is respired as CO₂ directly to atmosphere, with a fixed fraction entering into fast- and slow-turnover soil organic pools (fSOM and sSOM) (Sitch et al., 2003). The overall decomposition rate in the model is strongly influenced by soil temperature and moisture (Sitch et al., 2003). Additionally, the model also estimates emissions of biogenic volatile organic compounds (BVOC) per PFT (Arneeth et al., 2007; Schurgers et al., 2009). However, the modelled BVOC values are not compared in the current study, because they only represent a very small fraction of the modelled NEE and, additionally, there is insufficient growing season measurement data in the study domain to evaluate model performance. The major C cycling pathways can be found in the solid line-box area of Fig. 1. For peatland cells, an extra potential C pool for methanogens (C_{CH₄}) has been added (see dashed line-box area in Fig. 1) and mainly includes root exudates and easily-decomposed materials (Wania et al., 2010). The majority of C_{CH₄} is located in the acrotelm layer (Sect. S1) and the oxidation and production of CH₄ together determine the net emission of CH₄. In the model, the oxidation of CH₄ through methanotrophic bacteria is turned into CO₂, whereas the un-oxidized CH₄ can be released to the atmosphere by plant-transport, diffusion and ebullition (see the lines with a solid arrow in Fig. 1). The oxidation level is mainly determined by the location of water table position (WTP) in the model (Wania et al., 2010).

The biomass production in the current version of LPJ-GUESS WHyMe has no representation of nitrogen (N) limitation and neither N fluxes nor C–N interactions are included (Sitch et al., 2007). The latest version of LPJ-GUESS does include N cycling and N limitation on plant production (Smith et al., 2014), but this capability has not yet been integrated with the customized arctic version of the model adopted in the present paper. Moreover, processes determining concentrations of DOC and DIC in soil water have not yet been explicitly described in the model (Fig. 1). To cover the majority of

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



dissolved C losses and gains in our assessment of the catchment C budget, DOC is estimated by combining modelled runoff with observed DOC concentrations. Measured DIC export data is used directly, based on observations by Olefeldt et al. (2013).

3 Study site and materials

3.1 Stordalen catchment

The Stordalen catchment, located in the discontinuous permafrost region, is situated around 9.5 km east of the Abisko Naturvetenskapliga Station (ANS, Abisko Research Station). The catchment, covering around 16 km², has a steep topography on the southern part and turns into the lower flat peatland region to the north (Fig. 2b). The mean air temperature obtained from ANS records was -0.7°C for the period 1913–2002 (Christensen et al., 2004) and 0.49°C for the period 2002–2011 (Callaghan et al., 2013). Climate details of the two warming periods in 20th century are given in Callaghan et al. (2010). The Stordalen catchment is composed of birch forest, tundra heath, peatland, lakes and rivers. Permafrost is only found in the palsa and recent permafrost loss and conversion of palsa into non-permafrost *Eriophorum* sites have been related to the observed warming trends in the region (Åkerman and Johansson, 2008). The *Sphagnum* sites have a variable active layer depth and are dominated by *Sphagnum* spp. (PFT: pmoss), while the *Eriophorum* sites are dominated by flood-tolerant *Eriophorum* spp. (PFT: WetGRS). Birch forest (PFT: IBS, e.g. *Betula pubescens* ssp. *czerepanovii*), which forms the tree line in this region, is mainly distributed around and to the south of the peatlands and extends upwards to mountain slope areas (Fig. 2). Above the birch tree line, there are extensive areas of sub arctic ericaceous dwarf-shrub heath dominated by the evergreen species *Vaccinium vitis-idaea* and the deciduous species *V. uliginosum* and *V. myrtillus*. Willow shrubs (*Salix* spp.) and dwarf birch shrubs (*Betula nana*) occur at the fringes of the forest and in wetter areas above tree line. The heath shrub species also occur as a ground layer

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in the forest (see Callaghan et al. (2013) for details of vegetation distribution and changes). Around 5% of the catchment is covered by lakes or small rivers (Lundin et al., 2013). A detailed description of catchment hydrological conditions can be found in Olefeldt et al. (2013). The map in Fig. 2a is based on an object-based vegetation analysis in this area (Lundin et al., 2014). The classification is based on an aerial imagery dataset collected in August 2008, with a spatial resolution of 0.08 m by 0.08 m.

3.2 Model inputs and modelling protocol

3.2.1 Model inputs

Monthly climate data at 50 m resolution, including temperature, precipitation, and cloudiness, together with annual CO₂ concentration were used to drive the model from 1913 to 2080 for the Stordalen catchment. The detailed descriptions of data sources for the period of 1913–2010 can be found in Tang et al. (2014b). The high spatial resolution of monthly temperature for the period 1913–2000 was developed by Yang et al. (2011) covering the whole catchment, while precipitation and cloudiness were extracted and interpolated from the Climatic Research Unit 1.2 dataset (Mitchell et al., 2004). For the period 2001–2012, we obtained 12 years of observed temperature and precipitation data from ANS and interpolated these data to the whole region (Olefeldt et al., 2013). The annual CO₂ concentrations from 1913 to 2010 were obtained based on McGuire et al. (2001) and TRENDS (<http://cdiac.esd.ornl.gov/trends/co2/contents.htm>). The model was spun up in order to achieve vegetation and C pools in equilibrium with the climate at the beginning of the study period by using the first 30 years of climate forcing data to repeatedly drive the model for 300 years. To run the model into the future, the monthly anomalies of simulation outputs from the Rossby Centre Atmosphere Ocean (RCAO) regional climate model (Koenigk et al., 2011) for the grid cell nearest Stordalen were estimated and applied to the historical datasets at 50 m, assuming the same anomaly for all cells in the study region. RCAO climate data was downscaled for

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



an arctic domain using boundary forcing from a general circulation model forced by emissions from the SRES A1B scenario for the period 2013–2080 (Zhang et al., 2013).

A soil map provided by the Geological Survey of Sweden was used to identify the peatland fraction of each cell. Notably, the detailed rock area shown in Fig. 2a is not represented by the soil map used. Imagery Digital Elevation Model (DEM) from the National Land Survey of Sweden was used and the TFM algorithm (written in a Matlab script, R2012) was used to calculate topographic indices (Pilesjö and Hasan, 2014) used in the model LPJG-WHyMe-TFM.

3.2.2 Additional model parameterization and sensitivity testing

An artificially-adjusted snow density (100 kg m^{-3} , following the measurements ranges provided in Judson and Doesken, 2000) was implemented for birch forests to mimic deeper trapped snow in the forest. The observed thick top organic soil in the birch forest (Heliasz, 2012) is also represented in the model by increasing the organic fraction of the top 0.1 m soil layer (Olefeldt and Roulet, 2014) in the model. Additionally, the model simulation uses a single replicate patch in each 50 m grid cell. Finally, an additional simulation was performed in which we kept the CO_2 concentration constant after 1960 and allow the remaining inputs to vary as normal to diagnose the extent to which the CO_2 concentration influences on ecosystem dynamics and C fluxes.

3.3 Model evaluation data

The observations used to evaluate the model include: (1) EC-tower measured CO_2 NEE covering the palsa and *Sphagnum* sites during 2008 and 2009 (Olefeldt et al., 2012), (2) EC-tower measured CO_2 NEE at the *Eriophorum* site from 2006–2008 (Christensen et al., 2012), (3) two years (2006–2007) of EC-tower measured CH_4 fluxes at the *Eriophorum* site (Jackowicz-Korczyński et al., 2010), (4) analyzed DOC concentrations at the palsa/*Sphagnum* site and *Eriophorum* site from 2008 (Olefeldt and Roulet, 2012), (5) measured DIC export and DOC concentration at six sampling

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



points from 2007 to 2009 (Olefeldt et al., 2013), (6) EC-tower measured birch forest CO₂ NEE for four continuous years, 2007–2010 (Heliasz, 2012), (7) catchment-level annual CO₂ and CH₄ emission from lakes and streams from 2008 to 2011 (Lundin et al., 2013).

5 3.4 Calculation of the catchment carbon budget

The catchment level C budget was calculated using estimated C fluxes (CF_{*i*}) (after first accounting for observed DOC or DIC fluxes, where available) weighted by landcover fractions (*f_i* in Eq. 1). The symbol *i* in Eq. (1) represents different landcover types, in our case, birch forest, peatland (*Sphagnum* and *Eriophorum* sites), tundra heath and lake/river. To identify the temporal variations of C fluxes, the identification different landcover (except peatland region) is based on the dominant PFTs for each grid cell during the reference period of 2001–2012. The aerial photo-based classification of different peatland types is used when estimating peatland fluxes as a whole (Olefeldt et al., 2012).

$$15 \quad C_{\text{all}} = \sum_{i=1}^{i=5} CF_i \cdot f_i \quad (1)$$

4 Results

The modelled C fluxes for birch forest, peatland and tundra heath are first compared with the measured data for the period of observation. The seasonality and magnitudes of the fluxes are evaluated and then used to estimate the catchment-level C budget. The modelled long-term C dynamics during the period 1913–2080 are then presented for different landcover types and the catchment-level C budget considering all the available C components is also assessed.

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.1 Evaluation of the carbon fluxes in the historical period

4.1.1 Peatland

Both the plant communities and hydrological conditions in the peatland differ among the *palsa*, *Sphagnum* and *Eriophorum* sites. Notably, the measured NEE (see grey bars in Fig. 3a and b) covers both dry hummock (*palsa*) and semi-wet (*Sphagnum*) vegetation (Olefeldt et al., 2012), whereas our model cannot represent the dry conditions of the *palsa*. Therefore, the observed fluxes over both *palsa* and *Sphagnum* site were compared with the modelled fluxes at the *Sphagnum* site. The C fluxes magnitudes (including both wintertime emission and summertime uptake) are larger at the *Eriophorum* site, when compared with the *palsa/Sphagnum* site for both measured and modelled data. The modelled NEE at both sites generally captures the seasonality and magnitude of measured NEE, from being a strong sink (negative NEE) of CO₂ during the summer (mainly June–August) to being a wintertime CO₂ source. However, the model is unable to fully capture the C source/sink functionality in September at both sites. Furthermore, the modelled winter respiration at the *Eriophorum* site is very close to the observations, though the model overestimates the wintertime emissions at the *palsa/Sphagnum* site. The mean annual cumulative NEE reveals that the model estimations of C fluxes for both parts of the peatland are within the observed ranges, though with around 20 % of underestimation in the three-year averaged annual uptake (−39.76 and −50.18 gCm^{−2}yr^{−1} for the modelled and observed fluxes at the *palsa/Sphagnum* site; −71.54 and −90.34 gCm^{−2}yr^{−1} for the modelled and observed fluxes at the *Eriophorum* site). For the *Eriophorum* site, the three-year mean growing season uptake of C is underestimated by the model (Fig. 3d), which indicates that the modelled photosynthetic rates may be too low, or summer respiration rates may be too high, or both.

Two full years (2006 and 2007) of EC-tower measured CH₄ emissions were used to evaluate modelled estimations, and three different pathways of modelled CH₄ are also presented (Fig. 3e and f). The seasonal variability is well-described by the model

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the model's predictions. Furthermore, the observed CO₂ fluxes in September indicate a C source but the modelled fluxes are close to zero.

The red-shaded areas in Fig. 4a reveal high spatial variations during the growing season and the accumulated variations over the year (the average monthly SDs of the four years, see Fig. 4b) demonstrate a remarkable spatial variability of the model-estimated annual NEE. Interestingly, the observed mean annual NEE in both 2007–2008 and 2009–2010 fall within the wide spatial variations indicated by the model.

4.1.3 Tundra heath

Around 29.8% of the catchment with alpine terrain was covered by heaths and dwarf shrubs during the reference period 2001–2012. The model predicts that low-growing evergreen shrubs (PFT: LSE, e.g. *Vaccinium vitis-idaea*) currently dominate in this area, with tall summer-green shrub (PFT: HSS, e.g. *Salix* spp.) dominant in the future predictions (2051–2080) (Fig. 5). Since there is no available year-around observations of the C balance in this part of the Stordalen catchment, a synthesis of published data from similar environments is used to evaluate the model estimations (Table 1).

Four periods with averaged C uptake values from our model are presented (see the last four columns in Table 1) and a clear increase of summer uptake can be found with increased temperature and CO₂ concentration. The modelled, whole-year uptake also follows the increasing trend, except for the period of 2051–2080. The model-estimated summer C uptake is much stronger than the observations made at the high arctic heath of NE Greenland, which is reasonable when considering the longer growing season in our catchment. The study conducted in an alpine area in Abisko by Fox et al. (2008) shows a wide range of estimated NEE over 40 days during the summer time of 2004. The wide range of NEE values are because three levels of vegetation coverage were studied and the lowest uptake is from sparse vegetation dominated by bare rock and gravels, which is similar to the situation in our tundra heath sites (Fig. 2a). The modelled summer NEE between the years of 2001–2012 integrates the whole summer season (longer than 40 days), and falls within the observed ranges

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



presented by Fox et al. (2008), with values slightly higher than the lowest observed values.

4.1.4 Aquatic systems

Lateral waterborne C fluxes

5 The modelled DOC exports based on runoff estimations are compared with the measured DOC fluxes. The model generally underestimated annual runoff (the measured and modelled mean annual runoff for six points are 279.65 and 207.75 mm, respectively), but the modelled accuracy varies from point to point as well as for different years (Tang et al., 2014a). The average DOC export of the birch forest
10 over three years across sampling sites from 2007 to 2009 are 3.65, 2.77, and 2.33 $\text{gCm}^{-2}\text{yr}^{-1}$ from the observations and 3.09, 2.03, and 1.69 $\text{gCm}^{-2}\text{yr}^{-1}$ from the model estimations, respectively. The downward trend over the three years is captured and is related to lower precipitation during the latter years. The observed DOC export rates at the *palsa/Sphagnum* and *Eriophorum* sites in 2008 are directly
15 used to represent different types of peatland export level and the values are 3.35 and 7.55 $\text{gCm}^{-2}\text{yr}^{-1}$ for the *palsa/Sphagnum* and *Eriophorum* sites, respectively (Olefeldt and Roulet, 2012). DIC export is currently beyond the scope of our model but nonetheless contributes to the whole C budget. We used an averaged DIC export value of 1.22 $\text{gCm}^{-2}\text{yr}^{-1}$ based on the published data in Olefeldt et al. (2013) and DIC
20 export is included in the estimation of the catchment C budget below.

Carbon fluxes from lakes and streams

Investigations of aquatic system C emission in the Stordalen catchment were conducted by Lundin et al. (2013) during the years 2008–2011. Around 5% of the total catchment area (0.75 km^2) is classified as aquatic systems, with lakes accounting
25 for 96% of the aquatic area. Both lakes and streams contribute to the emissions of

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



CH₄ and CO₂, but the streams dominate CO₂ emission while the lakes dominate CH₄ emissions. Averaging across the catchment area, the measured annual CO₂ and CH₄ emissions from streams are 10.1 and 0.06 gCm⁻²yr⁻¹, respectively, while the lake emitted 0.5 gCm⁻²yr⁻¹ as CO₂ and 0.1 gCm⁻²yr⁻¹ as CH₄ (Lundin et al., submitted).

5 Since river and lake processes are at present beyond the scope of the model, the contribution of aquatic systems to the catchment C fluxes is purely based on the observed data.

4.2 Modelled whole-catchment carbon balance

10 One of the benefits of using a dynamic vegetation model is that it allows us to investigate the vegetation and C budget responses to climate change. In this section, annual variations of both climate variables (temperature and precipitation) and different ecosystem C fluxes are presented for the whole simulated period. In addition, the normalized catchment C budget is also estimated.

15 Our simulation results suggest that the temperature increase (around 2 °C, see Fig. 6-1a–c) together with a CO₂ increase to 639 ppm by 2080 could greatly increase the productivity of the birch forest (Figs. 6-2a–c and 5) as well as peatland CH₄ emissions (see Fig. 6-6a–c). The DOC export from birch sites also increases slightly over the study period (Fig. 6-3a–c). Besides leading to an increase in ecosystem biomass in grid cells currently occupied by birch forest (Fig. 5c and d), warming will also result in the current tundra heath close to the birch treeline being replaced by an upward expansion of the birch forest (Fig. 5a and b). This is indicated by the changes to birch treeline's uppermost elevation in Fig. 6-8a–c. The dramatic increase of C uptake in the tundra heath region (Fig. 6-5a–c) is largely a result of this vegetation succession. Meanwhile, the successive degradation of permafrost and slightly higher annual precipitation may result in more anaerobic conditions for the modelled peatland. Combined with warmer soil conditions, these result in both higher decomposition rates of soil organic matter and greatly increased CH₄ production. Moreover, a stronger response of respiration than NPP to the temperature increase reduces the net C

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



uptake in the peatland (see Fig. 6-4a–c). The catchment, as a whole, shows an increased C uptake (see the trend line in Fig. 6-7a) over the 1913–2080 period. The averaged C budgets for the two selected periods (1961–1990 and 2051–2080) are -1.47 and $-11.23 \text{ g C m}^{-2} \text{ yr}^{-1}$, respectively, with the increase being dominated by the large increase seen in the uptake rates at the birch forest and tundra heath sites.

During the 2051–2080 period, both boreal needle-leaf shade-tolerant spruce (PFT: BNE, e.g. *Picea abies*) and boreal needle-leaf (but less shade-tolerant) pine (PFT: BINE, e.g. *Pinus sylvestris*) start to appear in the birch-dominated regions (Fig. 5). In the current tundra regions, the coverage of HSS greatly increases at the expense of LSE. For the northern parts of the catchment, birch forest densification is observed with future warming, while the greatest relative changes in biomass occur near the treeline. The increased temperature together with increased CO_2 concentration by 2080 are very likely to increase CO_2 uptake in both tundra heath and forested areas, though the nutrient limitations are not included in this version of the model (but see Smith et al., 2014). A summary of the modelled catchment C components and C fluxes from different sources is given in Table 2. Additionally, the modelled estimations during the warm period of 2000–2005 are also presented in Table 2 and the positive value of catchment C-budget is seen to be an exception compared to the main reference period.

To illustrate the impact of CO_2 increases alone on the modelled C budget, the differences between simulations (ΔC fluxes) with and without a CO_2 increase since 1960 are shown in Fig. 7. The original outputs of those two simulations are plotted separated in Fig. S1 in the Supplement. Interestingly, the simulation with constant CO_2 forcing after 1960 greatly reduces the birch and tundra uptake (positive values in Fig. 7), whereas the peatland NEE and CH_4 are not strongly influenced by the CO_2 increase. The catchment C budget dynamics are consistent with the changes seen in the birch and tundra heath regions.

To account for the fact that CH_4 is a more potent greenhouse gas than CO_2 , an estimation of the global warming potential (GWP) of the two simulations can be made. Assuming the relative climate impact of CH_4 is 28 times greater than CO_2 over a 100

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



year period (IPCC, 2013), the calculated GWP for the simulation with atmospheric CO₂ increase is three times larger in the period 1961–1990 (27.3 g CO₂-eq) than the period 2051–2080 (8.8 g CO₂-eq). However, in the simulation without CO₂ increase, the GWP in 1961–1990 (40.7 g CO₂-eq) is approximately half of the GWP value for the 2051–2080 period (93.3 g CO₂-eq). This shows that the change in global warming potential found in the model simulations is strongly influenced by the CO₂ concentration used to force the model, as the CO₂ trajectory can alter the balance between the GWP changes resulting from C uptake in the birch forest and tundra, and the peatland CH₄ emissions.

5 Discussion

To our knowledge, our model simulation is the first attempt to create a C budget of a subarctic catchment based on a dynamic global vegetation model applied at the local scale and to predict its evolution in response to changing climate. The C budget estimations in this paper include major flux components (CO₂, CH₄ and hydrological C fluxes) based on a process-based approach. The magnitude and seasonality of modelled C fluxes compare well with the measurements (Figs. 3 and 4 and Tables 1 and 2), which gives us confidence in the ability of our model to represent the main processes influencing the C balance in this region. Our hope is that, by using a process-based modelling approach at high spatial resolution, our methodology will be more robust in estimating the C budget in a changing future climate than other budget estimation methods (Christensen et al., 2007; Worrall et al., 2007). In response to a climate warming scenario, our model shows a general increase in the C sink in both birch forest and tundra heath ecosystems, along with greater CH₄ emissions in the catchment (Fig. 6). Integrated over the catchment, our modelled C budget indicates that the region will be a greater sink of C by 2080, though these estimates are sensitive to the atmospheric CO₂ concentration. Nevertheless, the current model set-up and simulations still contain limitations in both the historical estimations and predictions.

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Below we discuss the model's current performance as well as a few existing limitations, and further propose some potential model developments. Most of the studies referred to below have been conducted specifically in the area near the Stordalen catchment or in other regions with similar environments.

5.1 Modelling carbon fluxes for the historical period

5.1.1 Peatland CO₂ and CH₄ emission

The model estimations in the peatland demonstrated skill in representing the relative differences of CO₂ fluxes between the *palsa/Sphagnum* site and *Eriophorum* site. The current model version cannot represent the very dry conditions of the *palsa* due to the acrotelm-catotelm soil structure (Wania et al., 2009a); therefore, the overestimated emissions rates of CO₂ (Fig. 3a and b) are expected when using the modelled fluxes from the *Sphagnum* site to compare with the measured data covering both the *palsa* and *Sphagnum* sites (Larsen et al., 2007a; Bäckstrand et al., 2010). The cold-season respiration at the *Sphagnum* site can account for at least 22 % of the annual CO₂ emissions (Larsen et al., 2007a), which is a further reason for the model overestimations. Moreover, at the *Eriophorum* site, the model slightly underestimated summertime uptake, which was found to be related to modelled summer respiration being higher than dark chamber-measured respiration, though with high spatial variations (Tang et al., 2014a).

The accurate representations of annual CH₄ emissions at the *Eriophorum* sites reflect the model's improved estimations of both hydrological conditions and dynamic C inputs to the C_{CH₄} pools (Fig. 1). With the inclusion of the distributed hydrology at 50 m resolution (Tang et al., 2014a, b), the lower-lying peatland can receive water from the upslope regions and the fact that water can rise to a depth of 10 cm above the surface both creates anaerobic conditions in the model and favours the establishment of flood-tolerant graminoid PFTs such as *Carex* spp., which can transport CH₄ to the atmosphere. Under such anaerobic conditions, decomposition rates are restricted and

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



part of the decomposed C becomes CH₄ (Wania et al., 2010). Nevertheless, it remains the case that excluding the palsa type could result in a general overestimation of CH₄ emission over the whole peatland.

5.1.2 Wintertime carbon fluxes, thickness of organic layer and disturbance in the birch forest

The high wintertime CO₂ fluxes to the atmosphere observed in the birch forest and the underestimations by our model (Fig. 4) highlight the importance of the representation of winter season C fluxes for the birch forest, particularly as winter temperatures (*T*) are increasing more than summer *T* (Callaghan et al., 2010). In the model, the respiration rate is linked to soil *T* and moisture and the underestimated respiration could be attributed to the lower *T* estimations in the wintertime (covering all months except June–September) when compared with the observed soil temperature at 5 and 10 cm depth in 2009 and 2010 (the modelled wintertime soil *T* is around 1.9°C lower than that observed at both depths). Also, the investigations of snow depth insulation influence on soil *T* and respiratory activity by Grogan and Jonasson (2003) have shown that soil *T* contributes to the greatest variations in respiratory activity at our birch sites. The birch forest is located on the relatively lower parts of the catchment (Fig. 2) and traps the wind-shifted lighter snow from the upland tundra heath, creating a much thicker snow pack and therefore significantly increasing the soil *T* (Groendahl et al., 2007; Larsen et al., 2007b; Luus et al., 2013). Snow depth measurements in the Abisko birch and tundra heath sites from 24 March to 7 April 2009 (<http://www.nabohome.org/cgi-bin/explore.pl?seq=131>) revealed the snowpack in the birch forest was 26.62 cm deeper on average than the snowpack in the tundra heath. The snow density for the birch forest was artificially decreased to 100 kg m⁻³ in the model in order to increase the snow depth in the absence of a wind-redistribution mechanism in our model, but it is still hard to capture the high soil *T* as well as high emission rates in winter.

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**C budget estimation
of a subarctic
catchment**

J. Tang et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The thickness of organic soil layer is another crucial component controlling birch site respiration in our catchment (Sjögersten and Wookey, 2002; Heliasz, 2012). The thickness of organic soil is most likely connected to the past transformation of the sites from heath to forest and is expected to decrease if birch biomass continues to increase (Hartley et al., 2012, 2013). Furthermore, Hartley et al. (2012) showed evidence of the decomposition of older soil organic matter during the middle of the growing season and concluded that with more productive forests in the future, the soil stocks of C will become more labile. From a model perspective, faster turnover rates of soil C pools (Sitch et al., 2007) could be used in climate warming model experiments to reflect the accelerated C mobilization in the soil. This could, to some degree, offset the stronger C uptake in the birch forest site in the simulation results shown here. In addition, the current parameterization of the organic fraction in the birch forest may not fully represent the organic horizons in reality soil.

The birch forest is cyclically influenced by the outbreak of moths (*E. autumnata*) and an outbreak in 2004 greatly affected the birch forest, resulting in a much lower C uptake than the average (Heliasz et al., 2011). Even though it is assumed that the fluxes during 2007–2010 had returned to the pre-defoliation levels (M. Heliasz, personal communication, 2013), it is still difficult to completely exclude the influence of insect disturbance in this forest. For the current model simulation, stochastic mortality and patch-destroying disturbance events have been included to account for the impacts of these random processes on ecosystem C cycling. However, to give a more accurate and reliable representation of the C fluxes in the birch forest, an explicit representation of insect impacts, outbreaks and their periodicity should be included (Wolf et al., 2008b) as well as other disturbances (Callaghan et al., 2013; Bjerke et al., 2014). For example, a warm event in winter 2007 caused a 26 % reduction in biomass over an area of over 1424 km² in the summer 2008 (Bokhorst et al., 2009).

5.1.3 Benefits of high spatial resolution and limitations from monthly temporal resolution

Subarctic ecosystems are characterized by small-scale variations in vegetation composition, hydrological conditions, nutrient characteristics and C fluxes (Lukeno and Billings, 1985; McGuire et al., 2002; Callaghan et al., 2013). The spatial resolution of 50 m in this model application allowed us to capture the diverse vegetation micro-types and their altitudinal gradient in the catchment, as well as their differential responses to climate changes (Figs. 5 and 6). This is unlikely to be well represented by a simple-averaging approach across the landscape, e.g. RCMs or GCMs. Furthermore, it is worth noting that the carbon cycling in the peatland, especially CH₄ fluxes, is sensitive to the WTP estimations (Tang et al., 2014a). Without spatially-distinguished climate and topographical data, it becomes impossible to implement our distributed hydrological scheme and thereby capture the peatland WTP dynamics. To our knowledge, the use of 50 m climate data as forcing of LPJG-WHyMe-TFM (Yang et al., 2011) is among the most detailed and comprehensive modelling exercises related to C cycling. Although this study focuses on C cycling, the innovative projects changes of vegetation micro-types at high spatial resolution are relevant to local stakeholders such as conservation managers and reindeer herders.

Nevertheless, the high spatial resolution data is currently produced with coarser monthly temporal resolution, which could restrict the model's ability to accurately estimate C fluxes at the start and end of the growing season (Figs. 3 and 4). Due to the dramatic variations of day length at high latitudes, a few days of misrepresenting the starting date of the growing season could significantly alter the estimated plant C uptake (Heliasz, 2012). The daily variations are difficult to capture from the interpolated quasi-daily values used in the model. Indeed, such variations strongly highlight the need for climate forcing at a higher temporal resolution for this region, due to the long daylight duration and short growing seasons.

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5.2 Projection uncertainties

A temperature increase of 2°C (Fig. 6) and elevated CO₂ concentrations could greatly increase vegetation growth and thus the C sink of the whole catchment. However, the densification and expansion of birch forest as well as the increased presence of boreal spruce and pine PFTs in our projected period (Fig. 5) could be strongly influenced by reindeer grazing and herbivore outbreaks (Hedenås et al., 2011; Callaghan et al., 2013), even though those detected changes are consistent with other models simulations (Wolf et al., 2008a; Miller and Smith, 2012) and general historical trends (Barnekow, 1999). Furthermore, climate warming may favor the spread of insect herbivores, so an assessment of ecosystem responses to future climate change cannot ignore these disturbances (Wolf et al., 2008b) and also other factors such as winter warming events (Bjerke et al., 2014)

Temperature increase results in a larger extent of permafrost degradation in the future. Meanwhile, the increased amount of available water from precipitation and lateral inflow may increase the degree of anoxia and further favors the flood-tolerant WetGRS PFT growth as well as CH₄ production. However, the exact extent of wetting or drying of peatland in the future is still highly uncertain, and the model prediction depends strongly on the climate scenario, permafrost thawing and the resulting balance between increased water availability and increased evapotranspiration. If the peatland drying is large enough, the reduced degree of anoxia could reduce CH₄ emissions in the future (Wrona et al., 2006; Riley et al., 2011). In our case, the peatland, located at the lower part of the catchment, receives water from the southern mountain region, and is more likely to become wetter in the near future in response to the increases in water availability (Wrona et al., 2006). Additionally, the determination of grid cell peat fractions in our simulation is dependent on the current (historical) soil map; therefore, the projections of peatland expansion to non-peatland cells cannot be reflected in the current model predictions. This could bring some additional uncertainties into the catchment C-budget estimations (Malmer et al., 2005; Marushchak et al., 2013).

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



To cover all the major C components in this catchment, the current estimations of the C budget used available, observed aquatic emissions and DOC concentration and DIC export values, all of which were assumed constant for the whole simulation period, 1913–2080. However, the direction and magnitude of changes to aquatic C fluxes are hard to quantify without modelling additional processes in these systems. Previous studies have found that the substantial thawing of permafrost as well as increased precipitation in recent decades have significantly increased total organic carbon (TOC) concentrations in lakes (Kokfelt et al., 2009; Karlsson et al., 2010). The increased loadings of nutrients and sediments in lakes are very likely to increase productivity of aquatic vegetation, but the effects may be offset by the increased inputs from terrestrial organic C and increases in respiration (Wrona et al., 2006; Karlsson et al., 2010). Emissions of CH₄ from aquatic systems are more likely to increase due to the longer ice-free season (Callaghan et al., 2010), increased methanogenesis in sediments and also increased CH₄ transport by vascular plants (Wrona et al., 2006). Furthermore, a recent study by Wik et al. (2014) found that CH₄ ebullition from lakes is strongly related to heat fluxes into the lakes. Therefore, future changes to energy fluxes together with lateral transports of dissolved C from terrestrial ecosystems to the aquatic ecosystem are especially important for predicting C emissions from aquatic systems.

As we have discussed, the dynamics of birch forest and to a lesser extent tundra heath C assimilation largely determines the catchment's C budget (Figs. 5 and 6), whereas the dramatic increase of CH₄ can slightly offset the net climate impact of the projected C uptake. Furthermore, both modelled C budget and GWP values are very sensitive to the atmospheric CO₂ levels. However, to date, there is no clear evidence showing significant long-term CO₂ fertilization effects in the arctic region (Oechel et al., 1994; Gwynn-Jones et al., 1997; Olsrud et al., 2010). Two possible reasons for this lack of CO₂ fertilization response might be that CO₂ levels close to the moss surface in birch forest can reach as high as 1140 ppm already and are normally within the range of 400–450 ppm (Sonesson et al., 1992) while tall

**C budget estimation
of a subarctic
catchment**J. Tang et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

5 vegetation such as shrubs and birch trees as well as peatland species have not been manipulated by CO₂ fertilization. Over the long-term, vegetation growth is likely a result of complex interactions between nutrient supplies (McNown and Sullivan, 2013), UV-B exposure (Schipperges and Gehrke, 1996; Johnson et al., 2002), temperature and growing season length (Heath et al., 2005) and forest longevity (Bugmann and Bigler, 2011). Therefore, more field experiments are urgently needed in order to quantify and understand the CO₂ fertilization effects on the various vegetation micro-types of the subarctic environment, and particularly tall vegetation types.

10 Overall, the current model application has been valuable in pointing to this gap in process understanding and meanwhile shows the importance of including vegetation dynamics in studies of C balance. Furthermore, a current inability to include the potential impacts of peatland expansion, potential increases of emissions from aquatic systems as well as the potential nutrient limitations on plants (but see Smith et al., 2014) and disturbances (Bjerke et al., 2014), make it likely that our projections of the catchment C-budget and CO₂-GWP will vary from those that may be observed in the future. However, our high spatial resolution, process-based modelling in the subarctic catchment provides an insight into the complexity of responses to climate change of a subarctic ecosystem while simultaneously revealing some key uncertainties that ought to be dealt with in future model development. These developments would be
20 aided by certain new observations and environmental manipulations, particularly of CO₂ with FACE experiments of shrubs and trees, in order to improve our understanding and quantification of complex subarctic processes.

**The Supplement related to this article is available online at
doi:10.5194/bgd-12-933-2015-supplement.**

25 *Acknowledgements.* The authors would like to thanks Rita Wania for providing the original LPJ-WHyMe code. We want to thank Thomas Holst, Guy Schurgers and Anneli Poska for interesting and informative discussions. We thank Ralf Döscher and Torben Koenig (SMHI)

for the providing us with regional climate projection data and anomalies. PAM acknowledges financial support from the ADSIMNOR (Advanced Simulation of Arctic Climate and Impact on Northern Regions) project funded by FORMAS (Swedish Research Council), and the Lund University Centre for the study of Climate and Carbon Cycle (LUCCI) funded by VR. The study is a contribution to the strategic research area Modelling the Regional and Global Earth System (MERGE), the Nordic Centre of Excellence DEFROST and the EU PAGE21 project. TVC and TRC wish to thank FORMAS for funding for the project “Climate change, impacts and adaptation in the sub-Arctic: a case study from the northern Swedish mountains” (214-2008-188) and the EU Framework 7 Infrastructure Project “INTERACT” (<http://www.eu-interact.org/>).

References

- Åkerman, H. J. and Johansson, M.: Thawing permafrost and thicker active layers in sub-arctic Sweden, *Permafrost Periglac.*, 19, 279–292, doi:10.1002/ppp.626, 2008.
- AMAP: Arctic Climate Issues 2011: Changes in Arctic Snow, Water, Ice and Permafrost, Oslo, 2012.
- Arneeth, A., Niinemets, Ü., Pressley, S., Bäck, J., Hari, P., Karl, T., Noe, S., Prentice, I. C., Serça, D., Hickler, T., Wolf, A., and Smith, B.: Process-based estimates of terrestrial ecosystem isoprene emissions: incorporating the effects of a direct CO₂-isoprene interaction, *Atmos. Chem. Phys.*, 7, 31–53, doi:10.5194/acp-7-31-2007, 2007.
- Bäckstrand, K.: Carbon Gas Biogeochemistry of a Northern Peatland – in a Dynamic Permafrost Landscape, PhD thesis, Faculty of Science, Department of Geology and Geochemistry, Stockholm University, 2008.
- Bäckstrand, K., Crill, P. M., Jackowicz-Korczyński, M., Mastepanov, M., Christensen, T. R., and Bastviken, D.: Annual carbon gas budget for a subarctic peatland, Northern Sweden, *Biogeosciences*, 7, 95–108, doi:10.5194/bg-7-95-2010, 2010.
- Barnekow, L.: Holocene tree-line dynamics and inferred climatic changes in the Abisko area, northern Sweden, based on macrofossil and pollen records, *The Holocene*, 9, 253–265, doi:10.1191/095968399676322637, 1999.
- Bjerke, J. W., Stein Rune, K., Kjell Arild, H., Eirik, M., Jane, U. J., Sarah, L., Dagrún, V.-S., and Hans, T.: Record-low primary productivity and high plant damage in the Nordic Arctic

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Region in 2012 caused by multiple weather events and pest outbreaks, *Environ. Res. Lett.*, 9, 084006, doi:10.1088/1748-9326/9/8/084006, 2014.

Bokhorst, S. F., Bjerke, J. W., Tømmervik, H., Callaghan, T. V., and Phoenix, G. K.: Winter warming events damage sub-Arctic vegetation: consistent evidence from an experimental manipulation and a natural event, *J. Ecol.*, 97, 1408–1415, doi:10.1111/j.1365-2745.2009.01554.x, 2009.

Bugmann, H. and Bigler, C.: Will the CO₂ fertilization effect in forests be offset by reduced tree longevity?, *Oecologia*, 165, 533–544, doi:10.1007/s00442-010-1837-4, 2011.

Callaghan, T. V., Bergholm, F., Christensen, T. R., Jonasson, C., Kokfelt, U., and Johansson, M.: A new climate era in the sub-Arctic: accelerating climate changes and multiple impacts, *Geophys. Res. Lett.*, 37, L14705, doi:10.1029/2009GL042064, 2010.

Callaghan, T. V., Jonasson, C., Thierfelder, T., Yang, Z., Hedenås, H., Johansson, M., Molau, U., Van Bogaert, R., Michelsen, A., Olofsson, J., Gwynn-Jones, D., Bokhorst, S., Phoenix, G., Bjerke, J. W., Tømmervik, H., Christensen, T. R., Hanna, E., Koller, E. K., and Sloan, V. L.: Ecosystem change and stability over multiple decades in the Swedish subarctic: complex processes and multiple drivers, *Philos. T. R. Soc. B*, 368, 20120488, doi:10.1098/rstb.2012.0488, 2013.

Christensen, T. R., Johansson, T., Åkerman, H. J., Mastepanov, M., Malmer, N., Friborg, T., Crill, P., and Svensson, B. H.: Thawing sub-arctic permafrost; effects on vegetation and methane emissions, *Geophys. Res. Lett.*, 31, L04501, doi:10.1029/2003GL018680, 2004.

Christensen, T. R., Johansson, T., Olsrud, M., Strom, L., Lindroth, A., Mastepanov, M., Malmer, N., Friborg, T., Crill, P., and Callaghan, T. V.: A catchment-scale carbon and greenhouse gas budget of a subarctic landscape, *Philos. T. R. Soc. A*, 365, 1643–1656, 2007.

Christensen, T. R., Jackowicz-Korczyński, M., Aurela, M., Crill, P., Heliasz, M., Mastepanov, M., and Friborg, T.: Monitoring the multi-year carbon balance of a subarctic palsa mire with micrometeorological techniques, *Ambio*, 41, 207–217, doi:10.1007/s13280-012-0302-5, 2012.

Crank, J. and Nicolson, P.: A practical method for numerical evaluation of solutions of partial differential equations of the heat-conduction type, *Adv. Comput. Math.*, 6, 207–226, doi:10.1007/BF02127704, 1996.

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Fox, A. M., Huntley, B., Lloyd, C. R., Williams, M., and Baxter, R.: Net ecosystem exchange over heterogeneous Arctic tundra: scaling between chamber and eddy covariance measurements, *Global Biogeochem. Cy.*, 22, GB2027, doi:10.1029/2007GB003027, 2008.
- Gerten, D., Schaphoff, S., Haberlandt, U., Lucht, W., and Sitch, S.: Terrestrial vegetation and water balance-hydrological evaluation of a dynamic global vegetation model, *J. Hydrol.*, 286, 249–270, doi:10.1016/j.jhydrol.2003.09.029, 2004.
- Groendahl, L., Friborg, T., and Soegaard, H.: Temperature and snow-melt controls on interannual variability in carbon exchange in the high Arctic, *Theor. Appl. Climatol.*, 88, 111–125, doi:10.1007/s00704-005-0228-y, 2007.
- Grogan, P. and Jonasson, S.: Controls on annual nitrogen cycling in the understory of a Subarctic birch forest, *Ecology*, 84, 202–218, 2003.
- Grogan, P. and Jonasson, S.: Ecosystem CO₂ production during winter in a Swedish subarctic region: the relative importance of climate and vegetation type, *Glob. Change Biol.*, 12, 1479–1495, doi:10.1111/j.1365-2486.2006.01184.x, 2006.
- Gwynn-Jones, D., Lee, J. A., and Callaghan, T. V.: Effects of enhanced UV-B radiation and elevated carbon dioxide concentrations on a sub-arctic forest heath ecosystem, *Plant Ecol.*, 128, 243–249, doi:10.1023/A:1009771125992, 1997.
- Hartley, I. P., Garnett, M. H., Sommerkorn, M., Hopkins, D. W., Fletcher, B. J., Sloan, V. L., Phoenix, G. K., and Wookey, P. A.: A potential loss of carbon associated with greater plant growth in the European Arctic, *Nature Climate Change*, 2, 875–879, doi:10.1038/nclimate1575, 2012.
- Hartley, I. P., Garnett, M. H., Sommerkorn, M., Hopkins, D. W., and Wookey, P. A.: The age of CO₂ released from soils in contrasting ecosystems during the arctic winter, *Soil Biol. Biochem.*, 63, 1–4, doi:10.1016/j.soilbio.2013.03.011, 2013.
- Haxeltine, A. and Prentice, I. C.: BIOME3: An equilibrium terrestrial biosphere model based on ecophysiological constraints, resource availability, and competition among plant functional types, *Global Biogeochem. Cy.*, 10, 693–709, doi:10.1029/96GB02344, 1996.
- Haxeltine, A., Prentice, I. C., and Creswell, I. D.: A coupled carbon and water flux model to predict vegetation structure, *J. Veg. Sci.*, 7, 651–666, 1996.
- Heath, J., Ayres, E., Possell, M., Bardgett, R. D., Black, H. I. J., Grant, H., Ineson, P., and Kerstiens, G.: Rising atmospheric CO₂ reduces sequestration of root-derived soil carbon, 309, 1711–1713, doi:10.1126/science.1110700, 2005.

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hedenås, H., Olsson, H., Jonasson, C., Bergstedt, J., Dahlberg, U., and Callaghan, T.: Changes in tree growth, biomass and vegetation over a 13-year period in the Swedish sub-Arctic, *Ambio*, 40, 672–682, doi:10.1007/s13280-011-0173-1, 2011.

Heliasz, M.: Spatial and Temporal Dynamics of Subarctic Birch Forest Carbon Exchange, PhD thesis, Department of Physical Geography and Ecosystems Science, Lund University, Sweden, 132 pp., 2012.

Heliasz, M., Johansson, T., Lindroth, A., Mölder, M., Mastepanov, M., Friborg, T., Callaghan, T. V., and Christensen, T. R.: Quantification of C uptake in subarctic birch forest after setback by an extreme insect outbreak, *Geophys. Res. Lett.*, 38, L01704, doi:10.1029/2010GL044733, 2011.

Hickler, T., Smith, B., Sykes, M. T., Davis, M. B., Sugita, S., and Walker, K.: Using a generalized vegetation model to simulate vegetation dynamics in northeastern USA, *Ecology*, 85, 519–530, doi:10.1890/02-0344, 2004.

Hickler, T., Vohland, K., Feehan, J., Miller, P. A., Smith, B., Costa, L., Giesecke, T., Fronzek, S., Carter, T. R., Cramer, W., Kühn, I., and Sykes, M. T.: Projecting the future distribution of European potential natural vegetation zones with a generalized, tree species-based dynamic vegetation model, *Global Ecol. Biogeogr.*, 21, 50–63, doi:10.1111/j.1466-8238.2010.00613.x, 2012.

IPCC: Climate Change 2013: The Physical Science Basis, NY, USA, 2013.

Jackowicz-Korczyński, M., Christensen, T. R., Bäckstrand, K., Crill, P., Friborg, T., Mastepanov, M., and Ström, L.: Annual cycle of methane emission from a subarctic peatland, *J. Geophys. Res.-Biogeo.*, 115, G02009, doi:10.1029/2008JG000913, 2010.

Johnson, D., Campbell, C. D., Lee, J. A., Callaghan, T. V., and Gwynn-Jones, D.: Arctic microorganisms respond more to elevated UV-B radiation than CO₂, *Nature*, 416, 82–83, doi:10.1038/416082a, 2002.

Jonasson, S. and Michelsen, A.: Nutrient cycling in Subarctic and Arctic ecosystems, with special reference to the Abisko and Torneträsk Region, *Ecol. Bull.*, 45, 45–52, 1996.

Judson, A. and Doesken, N.: Density of freshly fallen snow in the Central Rocky Mountains, *B. Am. Meteorol. Soc.*, 81, 1577–1587, doi:10.1175/1520-0477(2000)081<1577:DOFFSI>2.3.CO;2, 2000.

Karlsson, J., Christensen, T. R., Crill, P., Förster, J., Hammarlund, D., Jackowicz-Korczynski, M., Kokfelt, U., Roehm, C., and Rosén, P.: Quantifying the relative importance of lake emissions

C budget estimation of a subarctic catchment

J. Tang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



in the carbon budget of a subarctic catchment, *J. Geophys. Res.-Biogeo.*, 115, G03006, doi:10.1029/2010JG001305, 2010.

Keuper, F., van Bodegom, P. M., Dorrepaal, E., Weedon, J. T., van Hal, J., van Logtestijn, R. S. P., and Aerts, R.: A frozen feast: thawing permafrost increases plant-available nitrogen in subarctic peatlands, *Glob. Change Biol.*, 18, 1998–2007, doi:10.1111/j.1365-2486.2012.02663.x, 2012.

Koenigk, T., Döscher, R., and Nikulin, G.: Arctic future scenario experiments with a coupled regional climate model, *Tellus A*, 63, 69–86, doi:10.1111/j.1600-0870.2010.00474.x, 2011.

Kokfelt, U., Rosen, P., Schoning, K., Christensen, T. R., FÖRster, J., Karlsson, J., Reuss, N., Rundgren, M., Callaghan, T. V., Jonasson, C., and Hammarlund, D.: Ecosystem responses to increased precipitation and permafrost decay in subarctic Sweden inferred from peat and lake sediments, *Glob. Change Biol.*, 15, 1652–1663, doi:10.1111/j.1365-2486.2009.01880.x, 2009.

Koven, C. D., Ringeval, B., Friedlingstein, P., Ciais, P., Cadule, P., Khvorostyanov, D., Krinner, G., and Tarnocai, C.: Permafrost carbon–climate feedbacks accelerate global warming, *P. Natl. Acad. Sci. USA*, 108, 14769–14774, doi:10.1073/pnas.1103910108, 2011.

Larsen, K. S., Grogan, P., Jonasson, S., and Michelsen, A.: Respiration and microbial dynamics in two subarctic ecosystems during winter and spring thaw: effects of increased snow depth, *Arct. Antarct. Alp. Res.*, 39, 268–276, doi:10.1657/1523-0430(2007)39[268:RAMDIT]2.0.CO;2, 2007a.

Larsen, K. S., Ibrom, A., Jonasson, S., Michelsen, A., and Beier, C.: Significance of cold-season respiration and photosynthesis in a subarctic heath ecosystem in Northern Sweden, *Glob. Change Biol.*, 13, 1498–1508, doi:10.1111/j.1365-2486.2007.01370.x, 2007b.

Lukeno, J. O. and Billings, W. D.: The influence of microtopographic heterogeneity on carbon dioxide efflux from a subarctic bog, *Ecography*, 8, 306–312, doi:10.1111/j.1600-0587.1985.tb01183.x, 1985.

Lundin, E. J., Giesler, R., Persson, A., Thompson, M. S., and Karlsson, J.: Integrating carbon emissions from lakes and streams in a subarctic catchment, *J. Geophys. Res.-Biogeo.*, 118, 1200–1207, doi:10.1002/jgrg.20092, 2013.

Luus, K. A., Lin, J. C., Kelly, R. E. J., and Duguay, C. R.: Subnivean Arctic and sub-Arctic net ecosystem exchange (NEE): towards representing snow season processes in models of NEE using cryospheric remote sensing, *Prog. Phys. Geog.*, 37, 484–515, doi:10.1177/0309133313491130, 2013.

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Malmer, N., Johansson, T., Olsrud, M., and Christensen, T. R.: Vegetation, climatic changes and net carbon sequestration in a North-Scandinavian subarctic mire over 30 years, *Glob. Change Biol.*, 11, 1895–1909, doi:10.1111/j.1365-2486.2005.01042.x, 2005.

Marushchak, M. E., Kiepe, I., Biasi, C., Elsakov, V., Friborg, T., Johansson, T., Soegaard, H., Virtanen, T., and Martikainen, P. J.: Carbon dioxide balance of subarctic tundra from plot to regional scales, *Biogeosciences*, 10, 437–452, doi:10.5194/bg-10-437-2013, 2013.

Mastepanov, M., Sigsgaard, C., Dlugokencky, E. J., Houweling, S., Strom, L., Tamstorf, M. P., and Christensen, T. R.: Large tundra methane burst during onset of freezing, *Nature*, 456, 628–630, doi:10.1038/nature07464, 2008.

McGuire, A. D., Sitch, S., Clein, J. S., Dargaville, R., Esser, G., Foley, J., Heimann, M., Joos, F., Kaplan, J., Kicklighter, D. W., Meier, R. A., Melillo, J. M., Moore, B., Prentice, I. C., Ramankutty, N., Reichenau, T., Schloss, A., Tian, H., Williams, L. J., and Wittenberg, U.: Carbon balance of the terrestrial biosphere in the Twentieth Century: analyses of CO₂, climate and land use effects with four process-based ecosystem models, *Global Biogeochem. Cy.*, 15, 183–206, doi:10.1029/2000GB001298, 2001.

McGuire, A. D., Wirth, C., Apps, M., Beringer, J., Clein, J., Epstein, H., Kicklighter, D. W., Bhatti, J., Chapin, F. S., de Groot, B., Efremov, D., Eugster, W., Fukuda, M., Gower, T., Hinzman, L., Huntley, B., Jia, G. J., Kasischke, E., Melillo, J., Romanovsky, V., Shvidenko, A., Vaganov, E., and Walker, D.: Environmental variation, vegetation distribution, carbon dynamics and water/energy exchange at high latitudes, *J. Veg. Sci.*, 13, 301–314, doi:10.1111/j.1654-1103.2002.tb02055.x, 2002.

McGuire, A. D., Christensen, T. R., Hayes, D., Heroult, A., Euskirchen, E., Kimball, J. S., Koven, C., Lafleur, P., Miller, P. A., Oechel, W., Peylin, P., Williams, M., and Yi, Y.: An assessment of the carbon balance of Arctic tundra: comparisons among observations, process models, and atmospheric inversions, *Biogeosciences*, 9, 3185–3204, doi:10.5194/bg-9-3185-2012, 2012.

McNown, R. W. and Sullivan, P. F.: Low photosynthesis of treeline white spruce is associated with limited soil nitrogen availability in the Western Brooks Range, Alaska, *Funct. Ecol.*, 27, 672–683, doi:10.1111/1365-2435.12082, 2013.

Miller, P. A. and Smith, B.: Modelling tundra vegetation response to recent Arctic warming, *Ambio*, 41, 281–291, doi:10.1007/s13280-012-0306-1, 2012.

C budget estimation of a subarctic catchment

J. Tang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Mitchell, T. D., Carter, T. R., Jones, P. D., Hulme, M., and New, M.: A Comprehensive Set of High-Resolution Grids of Monthly Climate for Europe and the Globe: the Observed Record (1901–2000) and 16 Scenarios (2001–2100), Tyndall Centre, UEA, Norwich, UK, 2004.

Oechel, W. C., Cowles, S., Grulke, N., Hastings, S. J., Lawrence, B., Prudhomme, T., Riechers, G., Strain, B., Tissue, D., and Vourlitis, G.: Transient nature of CO₂ fertilization in Arctic tundra, *Nature*, 371, 500–503, 1994.

Olefeldt, D. and Roulet, N. T.: Effects of permafrost and hydrology on the composition and transport of dissolved organic carbon in a subarctic peatland complex, *J. Geophys. Res.-Biogeo.*, 117, G01005, doi:10.1029/2011jg001819, 2012.

Olefeldt, D. and Roulet, N. T.: Permafrost conditions in peatlands regulate magnitude, timing, and chemical composition of catchment dissolved organic carbon export, *Glob. Change Biol.*, 20, 3122–3136, doi:10.1111/gcb.12607, 2014.

Olefeldt, D., Roulet, N. T., Bergeron, O., Crill, P., Bäckstrand, K., and Christensen, T. R.: Net carbon accumulation of a high-latitude permafrost tundra similar to permafrost-free peatlands, *Geophys. Res. Lett.*, 39, L03501, doi:10.1029/2011GL050355, 2012.

Olefeldt, D., Roulet, N., Giesler, R., and Persson, A.: Total waterborne carbon export and DOC composition from ten nested subarctic peatland catchments – importance of peatland cover, groundwater influence, and inter-annual variability of precipitation patterns, *Hydrol. Process.*, 27, 2280–2294, doi:10.1002/hyp.9358, 2013.

Olsrud, M., Carlsson, B. Å., Svensson, B. M., Michelsen, A., and Melillo, J. M.: Responses of fungal root colonization, plant cover and leaf nutrients to long-term exposure to elevated atmospheric CO₂ and warming in a subarctic birch forest understory, *Glob. Change Biol.*, 16, 1820–1829, doi:10.1111/j.1365-2486.2009.02079.x, 2010.

Pilesjö, P. and Hasan, A.: A triangular form-based multiple flow algorithm to estimate overland flow distribution and accumulation on a Digital Elevation Model, *Transactions in GIS*, 18, 108–124, doi:10.1111/tgis.12015, 2014.

Riley, W. J., Subin, Z. M., Lawrence, D. M., Swenson, S. C., Torn, M. S., Meng, L., Mahowald, N. M., and Hess, P.: Barriers to predicting changes in global terrestrial methane fluxes: analyses using CLM4Me, a methane biogeochemistry model integrated in CESM, *Biogeosciences*, 8, 1925–1953, doi:10.5194/bg-8-1925-2011, 2011.

Roulet, N. T., Lafleur, P. M., Richard, P. J. H., Moore, T. R., Humphreys, E. R., and Bubier, J.: Contemporary carbon balance and late Holocene carbon accumulation in a northern peatland, *Glob. Change Biol.*, 13, 397–411, doi:10.1111/j.1365-2486.2006.01292.x, 2007.

- Schipperges, B. and Gehrke, C.: Photosynthetic characteristics of subarctic mosses and lichens, *Ecol. Bull.*, 45, 121–126, 1996.
- Schurgers, G., Arneth, A., Holzinger, R., and Goldstein, A. H.: Process-based modelling of biogenic monoterpene emissions combining production and release from storage, *Atmos. Chem. Phys.*, 9, 3409–3423, doi:10.5194/acp-9-3409-2009, 2009.
- 5 Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J. O., Levis, S., Lucht, W., Sykes, M. T., Thonicke, K., and Venevsky, S.: Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model, *Glob. Change Biol.*, 9, 161–185, doi:10.1046/j.1365-2486.2003.00569.x, 2003.
- 10 Sitch, S., McGuire, A. D., Kimball, J., Gedney, N., Gamon, J., Engstrom, R., Wolf, A., Zhuang, Q., Clein, J., and McDonald, K. C.: Assessing the carbon balance of circumpolar Arctic tundra using remote sensing and process modeling, *Ecol. Appl.*, 17, 213–234, doi:10.1890/1051-0761(2007)017[0213:ATCBOC]2.0.CO;2, 2007.
- 15 Sjögersten, S. and Wookey, P. A.: Climatic and resource quality controls on soil respiration across a forest–tundra ecotone in Swedish Lapland, *Soil Biol. Biochem.*, 34, 1633–1646, doi:10.1016/S0038-0717(02)00147-5, 2002.
- Smith, B., Prentice, I. C., and Sykes, M. T.: Representation of vegetation dynamics in the modelling of terrestrial ecosystems: comparing two contrasting approaches within European climate space, *Global Ecol. Biogeogr.*, 10, 621–637, doi:10.1046/j.1466-822X.2001.t01-1-00256.x, 2001.
- 20 Smith, B., Wårlind, D., Arneth, A., Hickler, T., Leadley, P., Siltberg, J., and Zaehle, S.: Implications of incorporating N cycling and N limitations on primary production in an individual-based dynamic vegetation model, *Biogeosciences*, 11, 2027–2054, doi:10.5194/bg-11-2027-2014, 2014.
- 25 Sonesson, M., Gehrke, C., and Tjus, M.: CO₂ environment, microclimate and photosynthetic characteristics of the moss *Hylocomium splendens* in a subarctic habitat, *Oecologia*, 92, 23–29, doi:10.1007/BF00317258, 1992.
- Tagesson, T., Mastepanov, M., Tamstorf, M. P., Eklundh, L., Schubert, P., Ekberg, A., Sigsgaard, C., Christensen, T. R., and Ström, L.: Satellites reveal an increase in gross primary production in a greenlandic high arctic fen 1992–2008, *Biogeosciences Discuss.*, 7, 1101–1129, doi:10.5194/bg-7-1101-2010, 2010.
- 30

**C budget estimation
of a subarctic
catchment**J. Tang et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

C budget estimation of a subarctic catchment

J. Tang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Tang, J., Miller, P. A., Crill, P. M., Olin, S., and Pilesjö, P.: Investigating the influence of two different flow routing algorithms on soil–water–vegetation interactions using the dynamic ecosystem model LPJ-GUESS, *Ecohydrology*, doi:10.1002/eco.1526, online first, 2014a.

Tang, J., Pilesjö, P., Miller, P. A., Persson, A., Yang, Z., Hanna, E., and Callaghan, T. V.: Incorporating topographic indices into dynamic ecosystem modelling using LPJ-GUESS, *Ecohydrology*, 7, 1147–1162, doi:10.1002/eco.1446, 2014b.

Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., and Zimov, S.: Soil organic carbon pools in the northern circumpolar permafrost region, *Global Biogeochem. Cy.*, 23, GB2023, doi:10.1029/2008GB003327, 2009.

Thonicke, K., Venevsky, S., Sitch, S., and Cramer, W.: The role of fire disturbance for global vegetation dynamics: coupling fire into a Dynamic Global Vegetation Model, *Global Ecol. Biogeogr.*, 10, 661–677, doi:10.1046/j.1466-822X.2001.00175.x, 2001.

Tranvik, L. J., Downing, J. A., Cotner, J. B., Loiselle, S. A., Striegl, R. G., Ballatore, T. J., Dillon, P., Finlay, K., Fortino, K., Knoll, L. B., Kortelainen, P. L., Kutser, T., Larsen, S., Laurion, I., Leech, D. M., Leigh McCallister, S., McKnight, D. M., Melack, J. M., Overholt, E., Porter, J. A., Prairie, Y., Renwick, W. H., Roland, F., Sherman, B. S., Schindler, D. W., Sobek, S., Tremblay, A., Vanni, M. J., Verschoor, A. M., Von Wachenfeldt, E., and Weyhenmeyer, G. A.: Lakes and reservoirs as regulators of carbon cycling and climate, *Limnol. Oceanogr.*, 54, 2298–2314, 2009.

Van Bogaert, R., Jonasson, C., De Dapper, M., and Callaghan, T. V.: Competitive interaction between aspen and birch moderated by invertebrate and vertebrate herbivores and climate warming, *Plant Ecology & Diversity*, 2, 221–232, doi:10.1080/17550870903487456, 2009.

Wania, R., Ross, I., and Prentice, I. C.: Integrating peatlands and permafrost into a dynamic global vegetation model; 1, Evaluation and sensitivity of physical land surface processes, *Global Biogeochem. Cy.*, 23, GB3014, doi:10.1029/2008gb003412, 2009a.

Wania, R., Ross, I., and Prentice, I. C.: Integrating peatlands and permafrost into a dynamic global vegetation model; 2, Evaluation and sensitivity of vegetation and carbon cycle processes, *Global Biogeochem. Cy.*, 23, GB3015, doi:10.1029/2008gb003413, 2009b.

Wania, R., Ross, I., and Prentice, I. C.: Implementation and evaluation of a new methane model within a dynamic global vegetation model: LPJ-WHYMe v1.3.1, *Geosci. Model Dev.*, 3, 565–584, doi:10.5194/gmd-3-565-2010, 2010.

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Wik, M., Thornton, B. F., Bastviken, D., MacIntyre, S., Varner, R. K., and Crill, P. M.: Energy input is primary controller of methane bubbling in subarctic lakes, *Geophys. Res. Lett.*, 41, 555–560, doi:10.1002/2013GL058510, 2014.

Wolf, A., Callaghan, T., and Larson, K.: Future changes in vegetation and ecosystem function of the Barents Region, *Climatic Change*, 87, 51–73, doi:10.1007/s10584-007-9342-4, 2008a.

Wolf, A., Kozlov, M., and Callaghan, T.: Impact of non-outbreak insect damage on vegetation in northern Europe will be greater than expected during a changing climate, *Climatic Change*, 87, 91–106, doi:10.1007/s10584-007-9340-6, 2008b.

Worrall, F., Burt, T., Adamson, J., Reed, M., Warburton, J., Armstrong, A., and Evans, M.: Predicting the future carbon budget of an upland peat catchment, *Climatic Change*, 85, 139–158, doi:10.1007/s10584-007-9300-1, 2007.

Wramneby, A., Smith, B., Zaehle, S., and Sykes, M. T.: Parameter uncertainties in the modelling of vegetation dynamics – effects on tree community structure and ecosystem functioning in European forest biomes, *Ecol. Model.*, 216, 277–290, doi:10.1016/j.ecolmodel.2008.04.013, 2008.

Wrona, F. J., Prowse, T. D., Reist, J. D., Hobbie, J. E., Lévesque, L. M. J., and Vincent, W. F.: Climate change effects on aquatic biota, ecosystem structure and function, *Ambio*, 35, 359–369, doi:10.1579/0044-7447(2006)35[359:CCEOAB]2.0.CO;2, 2006.

Yang, Z., Hanna, E., and Callaghan, T. V.: Modelling surface–air-temperature variation over complex terrain around Abisko, Swedish Lapland: uncertainty of measurements and models at different scales, *Geogr. Ann. A*, 93, 89–112, doi:10.1111/j.1468-0459.2011.00005.x, 2011.

Yang, Z., Sykes, M., Hanna, E., and Callaghan, T.: Linking fine-scale sub-Arctic vegetation distribution in complex topography with surface–air-temperature modelled at 50-m resolution, *Ambio*, 41, 292–302, doi:10.1007/s13280-012-0307-0, 2012.

Zhang, W., Miller, P. A., Smith, B., Wania, R., Koenigk, T., and Döscher, R.: Tundra shrubification and tree-line advance amplify arctic climate warming: results from an individual-based dynamic vegetation model, *Environ. Res. Lett.*, 8, 034023, doi:10.1088/1748-9326/8/3/034023, 2013.

C budget estimation
of a subarctic
catchment

J. Tang et al.

Table 2. Summary of catchment C-budget (in the unit of $\text{gC m}^{-2} \text{yr}^{-1}$) for different time periods. Negative values indicate ecosystem carbon uptake, while positive values indicate that mean ecosystem carbon is lost through respiration. The mean temperature (T , $^{\circ}\text{C}$) of each period is listed.

Periods	Birch CO_2^a	<i>Eriop.</i> CO_2^a	Palsa/Sphag. CO_2^a	Peatland CH_4	Tundra heath	Streams CO_2^c	Lakes CO_2^c	Streams CH_4^c	Lakes CH_4^c	Birch DIC	Peatland DIC	Birch DOC	<i>Eriop.</i> DOC	Palsa/Sphag. DOC	C budget
1961–1990 ($T = -1.26$)	-17.52	-36.83	-18.04	14.08	-4.31							2.52			-1.47
2000–2005 ($T = 0.13$)	-3.48	-32.85	-10.49	17.65	-1.63							3.25			7.96
2051–2080 ($T = 0.9$)	-24.56	-28.84	-8.43	22.94	-26.88							2.86			-11.23
Measured (Years)	0.88 ^g (2007– 2010)	-90.34 ^f (2006– 2008)	-50.18 ^d (2008– 2009)	20.23 ^b (2006– 2007)	-3 ^{d,h}	10.1 ± 4.4 ⁱ (2009– 2011)	0.5 ± 0.2 ^j (2008– 2011)	0.06 ⁱ (2009– 2011)	0.1 ± 0.1 ⁱ (2010)	1.22 ^j (2007– 2009)	1.22 ^j (2007– 2009)	3.17 ^j (2008– 2009)	7.55 ^k (2008)	3.35 ^k (2008)	

^a fluxes taken away DOC and DIC export;

^b observed CH_4 fluxes for the *Eriophorum* site;

^c values are normalized to the catchment area;

^d is a reference-estimated value;

^e Heliasz (2012);

^f Jackowicz-Korczyński et al. (2010);

^g Olefeldt et al. (2012);

^h Christensen et al. (2007);

ⁱ Lundin et al. (2014);

^j Olefeldt et al. (2013);

^k Olefeldt and Roulet (2012);

Eriop.: *Eriophorum* and *Sphag.*: *Sphagnum*.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



C budget estimation of a subarctic catchment

J. Tang et al.




Figure 1. Schematic of carbon components and cycling in LPJG-WHyMe-TFM. The solid line-box includes carbon cycling for non-peatland cell, whereas the items inside the dashed line-box represent processes particular to the peatland cells. Only DOC&DIC in red text is not explicitly presented in the current model. T : air temperature, P : precipitation, R : radiation, CO_2 : atmospheric CO_2 concentration, $Decom_C_{CH_4}$: decomposed materials allocated to C_{CH_4} , $Decom$: decomposition.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



C budget estimation of a subarctic catchment

J. Tang et al.




Figure 2. Object-based vegetation classification map of the Stordalen catchment **(a)** and schematic of landcover types and features of the approximate transect A–B in **(b)**.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



C budget estimation of a subarctic catchment

J. Tang et al.




Figure 3. Monthly (left) and mean annual cumulative (right) CO₂ NEE and CH₄ fluxes for the peatland. Positive values indicate ecosystem release to the atmosphere and negative values denote ecosystem uptake. Msd stands for the measured.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



C budget estimation of a subarctic catchment

J. Tang et al.




Figure 4. Monthly (a) and cumulative (b) NEE between the years 2007 and 2010 for the birch forest. Modelled NEE with spatial variability (red shadow) for each year is shown.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**C budget estimation
of a subarctic
catchment**

J. Tang et al.




Figure 5. Catchment dominated PFTs and their biomass for the periods of 1961–1990 and 2051–2080.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)


[Printer-friendly Version](#)

[Interactive Discussion](#)



C budget estimation of a subarctic catchment

J. Tang et al.



[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Figure 6. Time-series (column **a**) of annual mean temperature, precipitation sum, NEE ($\text{g C m}^{-2} \text{ yr}^{-1}$), averaged DOC export ($\text{g C m}^{-2} \text{ yr}^{-1}$) from the birch forest, birch tree line elevation (m) and catchment C-budget ($\text{g C m}^{-2} \text{ yr}^{-1}$) values between 1913 and 2080. The NEE data for the birch forest and the peatland have had the corresponding DOC and DIC values subtracted. Here the averaged C fluxes from the Stordalen peatland (the northeast side of the catchment) are used to represent for the averaged C fluxes for the whole catchment peatlands since only the Stordalen peatland DOC and DIC observations are currently available. The aerial photo-based classification of the *Eriophorum* and *Palsa/Sphagnum* peatland fractions within the Stordalen peatland is used to scale up the C fluxes. The trend of each dataset is shown with a red dashed line. The second and third columns (**b** and **c**) of the figure focus on the periods 1961–1990 and 2051–2080 (these two periods are also indicated in the first column with shaded area). The numbers in bold type in columns (**b**) and (**c**) show the annual average of each quantity for the respective time period. The fractions of peatland, birch forest, tundra and lakes/rivers are 5.7, 57.69, 29.76 and 5 %, respectively. There are around 1.6 % of the catchment area are estimated with the dominance of C3G and HSE which are not included in the above classification. The last row shows the birch tree line maximum elevation changes over the period.

BGD

12, 933–980, 2015

C budget estimation of a subarctic catchment

J. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



C budget estimation of a subarctic catchment

J. Tang et al.




Figure 7. Carbon flux differences (ΔC fluxes) between simulations with and without a CO_2 increase since 1960. Positive values imply a higher uptake (NEE) or a lower emission of CH_4 compared to the simulation without a CO_2 increase.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)