
Oregon Agricultural College

Extension Service

PAUL V. MARIS

Director

Cooperative Extension Work in Agriculture and Home Economics
Oregon Agricultural College and United States Department of Agriculture, Cooperating
Printed and distributed in furtherance of the Acts of Congress of
May 8 and June 30, 1914

Suggestive Points on Feeding for Egg Production

By

H. E. COSBY

Extension Specialist in Poultry Husbandry

If satisfactory results are to be secured from feeding for egg production the hens must be properly housed, have sufficient exercise, be healthful and vigorous, and possess an inborn tendency to lay. The feeding problem then becomes one of selecting the right feeds, proportioning them correctly and exercising judgment and common sense in carrying out the feeding operations.

Food is consumed by fowls to build up body tissues and to furnish energy. About seventy-five percent of the feed of a laying hen is required for body maintenance. In feeding for egg production it is imperative that the hen be given enough feed to supply her body needs and have a surplus of proper food elements to use in the production of eggs. It is expensive to feed sparingly, thus giving the hen no surplus above body maintenance.

No phase of poultry keeping requires better judgment or closer observation than that of feeding poultry for profit. Better results are secured when the fowls are fed according to their appetite than when they are fed according to rules. There is no best poultry feed, or ration, other than the one which supplies most economically the necessary food elements. When food value is considered, some poultry feed at \$3.00 per hundred pounds may be more expensive than other feeds at \$5.00 per hundred pounds.

Hens cannot do well on a whole grain ration. No grain ration supplies the various elements of food necessary in producing eggs. The continued use of a straight grain ration not only results in poor egg production but also results in digestive disorders. A ration should consist of grain and ground feeds. Poultrymen usually refer to the whole or cracked grains as "scratch food" and to the combination of ground feeds as "mash."

Hens being fed for egg production should be fed liberally on foods having the food elements found in the egg. An analysis of the egg shows that it is composed of ash 12.2 percent; water 65.7 percent; protein 11.4 percent; and fat 8.9 percent. The hen must be supplied with the necessary raw materials or she cannot manufacture the finished product. The hen cannot counterfeit her product.

NUTRITIVE ELEMENTS OF FOOD COMPOSITION

Ash produces bone and shell.

Protein produces muscle, tendons, blood, feathers, and whites of eggs.

Carbohydrates produce fat, heat, energy, and yolks of eggs.

Fats produce same as carbohydrates but are two and a quarter times as efficient.

Crude Fibre is only slightly digestible and has practically no nutritive value. It adds bulk to the ration and without it the concentrated feeds would form in a compact mass, thus preventing the digestive juices from functioning efficiently. A ration having more than five or six percent of crude fibre is not relished by poultry.

BALANCED RATION AND NUTRITIVE RATIO

A balanced ration is a ration which supplies in a palatable form the correct proportion of nutritive elements found in the egg.

Nutritive ratio is the ratio between the digestive protein and the digestive carbohydrates plus two and a quarter times the fat.

A ration which contains one pound of protein to from four to five pounds of carbohydrates plus the fat meets the requirements of laying hens. (Fowls in production require that a part of the protein in the ration come from animal sources, in addition to the vegetable protein found in the scratch and mash food.) The above ration would be termed a narrow ration and would be expressed as N. R. 1:4 or 1:5.

The nutritive ratio does not always indicate the true value of a ration and is no guarantee of high egg production. Nutritive ratio as herein figured is based upon the total amount of food elements and not upon the amounts actually digestible. Experiments have proved the superiority of the narrow ration.

AVERAGE PERCENTAGE COMPOSITION AND NUTRITIVE RATIO OF FOODS

	Fibre	Ash	Protein	Carbo- hydrates	Fat	Nutritive ratio
Corn	1.9	1.5	10.4	70.3	5.0	1:7.9
Wheat	1.8	1.8	11.9	71.9	2.1	1:6.3
Oats	9.5	3.0	11.8	59.7	5.0	1:6.1
Barley	2.7	2.4	12.4	69.8	1.8	1:6.
Kafir	1.4	1.5	9.9	74.9	3.0	1:8.3
Milo maize	2.4	2.8	8.7	66.2	2.2	1:8.2
Buckwheat	8.7	2.0	10.0	64.5	2.2	1:7.
Bran	9.0	5.8	15.4	53.9	4.0	1:4.1
Middlings	4.6	3.3	15.6	60.4	4.0	1:4.7
Mill-run	7.6	5.2	12.9	45.1	4.0	1:4.2
Gluten feed	6.4	1.3	23.2	54.7	6.3	1:2.9
Oil meal	9.5	5.8	33.2	38.4	3.0	1:1.4
Cocoonut meal	11.2	4.9	18.8	42.0	8.1	1:3.2
Meat scraps*	8.0	58.0	32.9	1:1.4
Fish scraps*	34.0	6.5	1:0.4

*Composition has a wide variation.

Knowing approximately the required nutritive ratio of a balanced ration for laying fowls, a study of the methods used in determining the nutritive ratio of any ration may be made by applying them to the following ration, which may or may not be desirable.

	Fibre	Protein	Carbo- hydrates	Fat
Scratch food				
50 pounds wheat9	5.95	35.95	1.05
25 pounds corn47	2.1	17.6	1.25
25 pounds oats	2.37	2.95	14.9	1.25
Mash food				
25 pounds bran	2.25	3.85	13.5	1.0
25 pounds middlings	1.5	3.9	15.1	1.0
15 pounds ground oats	1.43	1.77	10.96	.75
15 pounds corn meal23	1.56	10.55	.75
5 pounds oil meal47	1.66	1.92	.15
15 pounds meat scrap	0.	8.7	0.	1.65
	9.62	32.44	120.48	8.85

To determine the nutritive ratio of the above ration, multiply the fats by $2\frac{1}{4}$ ($8.85 \times 2\frac{1}{4}$ equals 19.91). Add the product to the carbohydrates (120.48 plus 19.91 equals 140.39). The nutritive ratio of this ration would be the ratio between 32.44 and 140.39. Dividing each number by 32.44 the nutritive ratio is found to be 1:4.4.

The 200-pound combination of feeds has 9.62 pounds of crude fibre or 4.8 percent. It is composed of grains each one of which the fowls like. It has one pound of protein to each four or five pounds of carbohydrates plus the fat, and should therefore be a reasonably good ration when properly fed.

RATIONS RECOMMENDED BY EXPERIMENT STATIONS IN DIFFERENT SECTIONS OF THE COUNTRY

Oregon Agricultural College Ration—

Scratch food
25 pounds wheat
15 pounds oats

Mash
5 pounds mill feed (bran and shorts)
5 pounds ground oats
5 pounds ground corn
5 pounds linseed meal (or)
7 pounds cocoanut meal
7 pounds meat or fish scraps

University of Missouri Ration—

Scratch food
2 parts cracked corn
1 part wheat

Mash
1 part bran
1 part middlings
1 part corn meal
1 part meat scrap

West Virginia Experiment Station Ration—

Scratch food
60 pounds cracked corn
60 pounds wheat
40 pounds oats
20 pounds barley
10 pounds buckwheat

Mash
200 pounds wheat bran
100 pounds corn meal
100 pounds ground oats
100 pounds gluten feed
100 pounds middlings
100 pounds meat scrap

METHOD OF FEEDING

The grain is usually fed in the litter morning and evening, about one third as much being fed in the morning as in the evening. The dry mash hopper should be open at all times. Fresh water, grit, and oyster shell should also be accessible at all times.

The amount of mash consumed is regulated by the amount of grain fed. Most poultrymen prefer to have the hens in the laying flock consume approximately equal parts by weight of mash and scratch feeds. This proportion will vary according to the weather and work being done by the hens. A flock of hens laying heavily will consume more mash than the same flock not laying. In cold weather the hens should ordinarily consume more grain than mash. The hens should have, in any case, not a limited number of pounds of grain, but all they will eat before going to roost.

If a moist, crumbly mash is fed, it is more advisable to feed it on a time basis rather than a quantity basis. After ten or fifteen minutes time, the moist mash that has not been consumed should be removed.

It is generally believed that for the average poultryman moist mash has more value as an emergency feed than as a regular part of the daily feeding system. It may be used advantageously in bringing a flock of pullets into an even lay in the fall. It is valuable as an emergency feed when a marked falling off in egg production is noted at any season of the year. After the moist mash has served its purpose it may be gradually eliminated from the feeding system.

Milk may be substituted for the meat or fish scrap in the mash. It is advisable, however, to have a small percentage of meat scrap in the mash because it makes the mash more palatable. Where milk is to be substituted for the full amount of animal protein, no water should be given the fowls, otherwise the substitution will not supply the needed animal protein.

A liberal supply of green feed should be given the fowls daily. The efficiency of any feed depends largely upon the amount of succulent feed consumed.

In addition to the five classes of food elements herein discussed, investigators have found present certain elements necessary to nutrition known as *vitamines*. The exact nature of *vitamines* is not accurately known and the exact amounts necessary in an economical ration are still in the stages of experimentation.

If fowls are worth keeping they are worth taking care of. It is more advisable to sell the flock than to neglect it.