

AN ABSTRACT OF THE THESIS OF

NIELS AAGE SKOV for the DOCTOR OF PHILOSOPHY  
(Name) (Degree)

in OCEANOGRAPHY presented on October 24, 1967  
(Major) (Date)

Title: THE ICE COVER OF THE GREENLAND SEA: AN EVALUA-  
TION OF OCEANOGRAPHIC AND METEOROLOGICAL  
CAUSES FOR YEAR-TO-YEAR VARIATIONS

Abstract approved: Redacted for Privacy  
Robert L. Smith

A criterion is defined to compare seasonal ice coverage in the Greenland Sea for the years 1900-57, and the areal coverage is graphed using the 1898-1913 average as a standard. The factors wielding possible influence on short-term variations of the ice cover are examined individually and their relative importance established.

The influence of ocean currents is evaluated by analysis of hydrosections across the East Greenland Current at 74°-76° N and across the North Atlantic Current in the Faeroe-Shetland channel. Data from the latter area are used for numerical analysis of heat imported to the Greenland Sea by the North Atlantic Current in the 1927-52 period. Details about the Irminger Current's behavior are derived from station data from Denmark Strait and from surface temperatures at Selvogsbanki south of Iceland. Year-to-year variations are found to exist in the flow volumes of all three currents,

and correlations with seasonal ice coverage in the Greenland Sea are shown.

Above-average precipitation in conjunction with below-average storm activity is found to have negligible influence on the ice regime, and no significant correlation is found. The possible effect of evaporation is computed to be far below the threshold of detectability. Air temperatures in the Norwegian-Greenland Sea region display a trend of increase throughout the period studied, in harmony with a concurrent trend of decreasing ice cover; but no causal relationship is in evidence.

The effects of strengths and directions of predominant winds are examined, and good correlations are shown between ice cover fluctuations and easterly wind components at Norwegian coastal stations. At the points of major currents' entrances to and exits from the Greenland Sea the wind effects are complex and cannot be fully evaluated on the basis of existing data.

The fluctuations of ocean currents entering and leaving the Greenland Sea and of water movements within the Greenland Sea remain as the apparent determinant of year-to-year variations of the ice cover.

The Ice Cover of the Greenland Sea: An Evaluation  
of Oceanographic and Meteorological Causes  
for Year-to-Year Variations

by

Niels Aage Skov

A THESIS

submitted to

Oregon State University

in partial fulfillment of  
the requirements for the  
degree of

Doctor of Philosophy

June 1968

As soon as one has passed over the deepest part of the ocean, he will encounter such a great superfluity of ice on the sea that I know of no equal of it anywhere else in the whole world. . . . These ice floes have peculiar habits. Sometimes they lie as quiet as can be . . . at other times they travel with a speed so swift and violent that a ship with a fair wind behind it is no more speedy; and when once in motion, they travel as often against the wind as with it.

Konungs Skuggsjá

c. 1250

APPROVED:

Redacted for Privacy

---

Assistant Professor of Oceanography  
in charge of major

Redacted for Privacy

---

Acting Chairman of the Department of Oceanography

Redacted for Privacy

---

Dean of Graduate School

Date thesis is presented October 24, 1967

Typed by Gwendolyn Hansen for Niels Aage Skov

## ACKNOWLEDGEMENTS

The writing of this thesis would not have been possible without the valuable advice of Dr. R. L. Smith and initial inspiration and encouragement by Dr. P. K. Weyl.

I owe a special debt of gratitude to Dr. June G. Pattullo for many fruitful discussions of oceanographic problems in the course of four years of work at Oregon State University.

The manuscript was read critically by Dr. J. Brookes Spencer, whose constructive comments engendered considerable improvement.

## TABLE OF CONTENTS

Chapter	Page
I. INTRODUCTION	1
II. ARCTIC ICE: DESCRIPTION AND DEFINITIONS	4
Classification of Arctic Sea Ice	4
The East Greenland Pack	6
An Ice Index	9
III. PREVIOUS WORK	14
IV. SOURCES AND PROCESSING OF DATA	19
V. CURRENT EFFECTS	27
The East Greenland Current	27
The North Atlantic Current	36
The Irminger Current	48
VI. CLIMATIC EFFECTS	56
Relevant Weather Stations	56
Weyl's Precipitation Hypothesis	58
Evaporation	61
Temperature Variations	62
Wind-Current Relationships	65
VII. DISCUSSION	75
BIBLIOGRAPHY	80

## LIST OF TABLES

Table	Page
1. Stations used for hydrosections across the East Greenland Current.	20
2. Stations in Denmark Strait from which data were used.	22
3. Stations in the Faeroe-Shetland channel from which data have been used.	23
4. Meteorological stations from which weather data have been used.	24
5. Heat content in cals $\times 10^{10}$ of the section, 182 km wide, 100 deep and 1 cm thick traversing the peak current intensity in Faeroe-Shetland channel	45
6. Summation of monthly anomalies in years with ice severity classifiable as rising, peak, declining or low.	47
7. Tabulation of years in which ice cover (x) deviated from its mean by a value larger than s (Figure 3).	55
8. Change in mean atmospheric pressure at selected stations from the period 1900-19 to 1920-39.	68
9. Correlation coefficients between Greenland Sea ice fluctuations and Wind Effect Indices (WEI) at selected stations in and near the Greenland Sea.	72
10. Anomalies of Wind Effect Indices in years the ice cover x deviated from its mean by an amount greater than s.	74



## LIST OF FIGURES

Figure	Page
1. Distribution of Arctic ice types.	5
2. Mean April ice limits for the period 1898-1913.	8
3. Extent of ice cover relative to 1898-1913 average (0-line).	11
4. Surface currents in the Greenland Sea.	28
5. Location of station lines B-05, <u>Belgica</u> expedition 1905; D-06/08, <u>Danmark</u> expedition 1906-08; G-30, <u>Godthaab</u> expedition 1930; and EG-91/92, <u>E. Greenland</u> expedition 1891-92.	32
6. Positions of isotherms in July 1905 and August 1906-08.	33
7. Positions of isotherms in August 1891 and July 1930.	35
8. Monthly temperature averages plus/minus one standard deviation for the area 60° N, 3°-5° W and 61° N, 5°-6° W.	39
9. Temperature ranges in Marsden Squares #217 and #218.	40
10. Average temperature, maximum range of temperature, and standard deviation by one-degree unit areas covering Iceland-Faeroe-Shetland section, as shown in Figure 8.	42
11. Average annual surface salinity of the area 58°-63° N, 10°-15° W.	48
12. Surface isotherms (blue) delineating boundary between Atlantic Water of Irminger Current and Polar Water of East Greenland Current in Denmark Strait, based on <u>Anton Dohrn</u> data from June 1955.	49

Figure	Page
13. July ice limits and mean positions of 34‰ isohalines in the years 1929, 1931 and 1932.	51
14. Seasonal variation of temperature at 100 m depth from annual mean in the eight degree unit areas shown.	52
15. February-May surface temperature anomaly at Selvogsbanki and ice cover curve (from Figure 3).	53
16. Locations of weather stations from which data were used in this study.	57
17. Precipitation anomalies of Angmagssalik, Grimsey, Jan Mayen, Spitzbergen, Bear Island and Tromsø summed by year and juxtaposed with ice cover graph (from Figure 3).	59
18. Air temperature anomalies of Spitzbergen and Grimsey-Spitzbergen-Røst-Thorshavn combined.	64
19. 5-year moving averages of ice cover curve (Figure 3) and annual combined temperature anomaly of stations Grimsey, Spitzbergen, Røst and Thorshavn.	66
20. Mean pressure distribution in the European Arctic (after Rodewald).	67
21. Difference in annual mean pressure between Thorshavn and Stykkisholm.	69

## PREFACE

Of all the diverse regions of the world ocean the Atlantic Arctic is to me the most tempting to investigate. Nowhere do the forces of nature reveal their interplay more vividly: intricate currents traverse the region in ever changing patterns, making the steady state an elusive dream; air-sea energy exchanges force from here the cool flows which travel for centuries through the bowels of the seas; and the albedo of a million square kilometers of glittering ice reflects the sun's heat, profoundly affecting weather patterns in ways that are as yet dimly understood.

An attempt to determine causes for past ice fluctuations must perforce rest on data gathered by earlier researchers. The work has been frequently frustrating, for although a quantity of material exists, the bulk is a conglomerate of data, fragmented in time and space. Rarely, if ever, does a series of reliable hydrographic measurements extend over a significant period of time. If at times I have been precipitate in drawing conclusions on fluid premises, I offer my regrets but hope that those who question my conjectures will themselves delve into the problems and fare better.

# THE ICE COVER OF THE GREENLAND SEA: AN EVALUATION OF OCEANOGRAPHIC AND METEOROLOGICAL CAUSES FOR YEAR-TO-YEAR VARIATIONS

## I. INTRODUCTION

Investigators of Arctic ice fluctuations have almost to a man concerned themselves with effects of rather than causes for variations in the severity of ice seasons. Consequently, the profound influence exerted by the ice cover upon European weather has been well documented by Knudsen (1905), Henry (1929), Petterson (1929), Montgomery (1940) and others, but almost no effort has been made to explore the factors that might bear on the phenomenon, including those in the realm of oceanography.

In a recent manuscript on the ocean's role in climatic change (Weyl, 1966) it was suggested that low storm activity in the period between ice melting and freezing in the Atlantic Arctic could significantly influence subsequent ice formation. The author argued that normal excess of precipitation over evaporation would tend to form a low-salinity surface water lens. With stable quiet weather preventing admixture of deeper, high-salinity water, sea ice in the following winter would freeze more rapidly and extend farther south than would be the case after a stormy summer. The possibility of the existence of any such simple direct relationship seemed utterly intriguing.

As it turns out, no such causal relationship is in evidence, but

the path of inquiry led to the broader question of possible relationships between ice fluctuations and measurable variations in oceanographic and meteorological parameters. The factors which could influence the ice regime can be enumerated as follows:

1. The magnitudes, distributions into Greenland Sea, salinities and surface temperatures of -
  - a) the East Greenland Current,
  - b) the North Atlantic Current
  - c) the Irminger Current
2. Precipitation, as it dilutes the surface water and raises the freezing point;
3. Evaporation, as it cools the surface and raises the surface salinity;
4. Air temperatures, as they affect the surface water temperatures;
5. Winds, as they affect the ocean currents and the ice transports.

Since variations do exist in both ice conditions and the oceanographic and meteorological parameters, it is my thesis that close, causal relationships exist, some of which can be roughly verified on the basis of existing data, and all of which can be quantified and perhaps predicted, given enough observational material. In the study supporting this thesis it is endeavored to encompass all of the factors

influencing the ice conditions, rather than concentrating on one or two.

## II. ARCTIC ICE: DESCRIPTION AND DEFINITIONS

### Classification of Arctic Sea Ice

Arctic sea ice presents a complex natural phenomenon which is still poorly understood, in part due to meagerness of observations. The three main ice types and their areas of formation can briefly be stated. In the northern hemisphere most of the sea ice forms on the surface in the north polar basin, where a thickness of about 2 meters is average (Transehe, 1928). This main body of floating ice is called the north polar cap ice, and it covers some 70% of the Arctic Ocean, or 2,000,000 square miles. Its margin closely parallels the 1000-meter isobath, Figure 1.

Another important ice type originates in the coastal regions in high latitudes, where in shallow bays and inlets and on wide, shallow shelves sea water in the cold season often freezes clear to the bottom and fast to shore. An estimated 150,000 square miles of this so-called fast ice forms every winter (Smith, 1931).

Surrounding the cap ice and extending fingers into the Greenland Sea and Baffin Bay we find the pack ice. The word "pack" does not necessarily denote that the ice is tightly packed together, for it may have large or small areas of open water within its bounds. Both the north polar cap ice and the pack ice are in continuous movement,




Figure 1. Distribution of Arctic ice types.



depending on the particular location and on the circulations of hydrosphere and atmosphere. Beside the varying movement of cap and pack several features of a general, slow drift have been identified, the dominant of which is the Transpolar Drift Stream. This wide swath of moving ice is the region of the Fram and Sedov drifts, and tracking of ice islands in recent years (Dunbar, 1962) has shown the motion to be clockwise at a rate of one mile per day (H. O. #705) north of the North American Continent and one to two miles per day along the Asian Continent, giving a net transport from the direction of Bering Strait toward the Greenland Sea, the main outlet for northern ice into warmer waters.

The pack in the Greenland Sea is fed not only from the stream traveling down from the Arctic Ocean, but also by a lesser stream passing seasonally from the Barents and Kara Seas south of Spitzbergen, and by fast ice dislodged during the summer. Local freezing in the Greenland Sea adds an amount of relatively thin ice rarely exceeding one meter in thickness (Schell, 1940). Apparently no one has tried to estimate the relative magnitude of this latter source, but Smith (1931) has characterized the Greenland Sea and the Baffin Bay as "the two most prolific regions of (pack) ice production."

#### The East Greenland Pack

In the Greenland Sea, pack ice terminates to the south in a zone

called the fringe of the pack, which shrinks in late summer and autumn and widens in winter and spring. The seasonal movement of the fringe limit denotes by its amplitude the severity of an ice year: when the limit is displaced far southeastward, ice conditions are termed heavy; when the limit stays well to the north-west, conditions are said to be light.

As winter settles, heavy pans and floes begin their influx into the Greenland Sea, reinforced by quantities of locally formed ice. The accumulation spreads along the coast of East Greenland, signalling at successively lower latitudes the beginning of the ice season. Throughout the winter and spring, large masses of ice continue to push southward and to spread away from the coast. Denmark Strait is normally more or less choked in spring, while in a bad year the ice blankets the northern coast of Iceland at that season.

The configuration of the mean ice limit points up the influences of oceanographic and meteorological factors upon the ice regime. North of Iceland the fringe in spring displays a characteristic curve toward the east, as shown in Figure 2, probably under the control of the East Iceland Current. An equally impressive feature is a V-shaped retrenchment immediately west of Spitzbergen, an unmistakable effect of the lingering warmth in the eddies of the northward thrusting branch of the North Atlantic Current. A less distinct indent just north of Jan Mayen has been in evidence in most seasons,




Figure 2. Ice limits for the month of April  
averaged over the years 1898-1913

----- maximum limit  
————— mean limit  
- · - · - minimum limit.

presumably caused by a smaller counter-clockwise eddy off the West Spitzbergen Current.<sup>1</sup>

### An Ice Index

Establishment of a practical criterion for severity of a given ice season, or definition of an "ice index," has been attempted in the past by Meinardus (1906) and others.<sup>2</sup> The method devised by Meinardus takes into account the duration of ice occurrence on the Iceland coast as well as the severity of the observed condition. The effect of weighting severe months is to enhance a graphic display and facilitate identification of the cyclic nature of the variations. Despite this advantage, Meinardus's method will not be emulated here, as the assessment of severity is a subjective matter open to substantial error.

The criterion used in the following is one of areal extent of sea surface coverage between the meridian through Cape Farewell and

---

<sup>1</sup> A favorite seal hunting ground in times past, it was known to sealers as "Nord-Bukta" (North Bay).

<sup>2</sup> Schott (1904) considered the position of the ice limit for each month of the season being evaluated. Smith (1931) used a scale of 1-10 to indicate degree of coverage in the area. Hansen's method (1943) is mentioned in chapter III.

15° E.<sup>3</sup> The Danish Meteorological Institute has compiled a detailed record of ice conditions in the Arctic seas since 1896, and its annually published charts and reports have provided the basic data.<sup>4</sup>

Figure 2 shows the average and extreme ice limits in the month of April for the period 1898-1913. The April average for this period serves in the present study as a "standard" against which all subsequent April charts have been compared. Deviations from this standard have been measured for each of the years 1900-39 and 1946-57, counted positive when the ice cover exceeded the area enclosed by the standard limit, negative when in a given year the ice covered a lesser area of the sea surface.

The same procedure has been followed for the months of May through August, and the average of all five months provides the final index for a given year. The ice fluctuations are graphically shown in Figure 3, where the 0-line represents the 1898-1913 average, and the ice cover's deviation is shown by year in ten-thousands of square kilometers. It is apparent that a trend exists toward less severe ice winters. Assuming the regression to be linear, the estimated

---

<sup>3</sup>There is apparently no correlation between ice conditions in the Greenland Sea and those in the Barents Sea (Schell, 1940). Hence separate treatment of Greenland Sea is justified.

<sup>4</sup>The charts are identical in scale and layout from 1899 to 1957, hence providing a good comparison between years of this period. No data exist from the war years 1940-45.




Figure 3. Extent of ice cover relative to 1898-1913 average (0-line).

regression equation is

$$\bar{y}_x = a + b(x - \bar{x})$$

where  $a = \frac{\Sigma y}{n} = -5.72$  and  $b = \frac{SP}{SS} = -0.066$ . The resulting regression line is indicated in the graph, together with the lines representing  $\pm s$ , where  $s = \frac{(\Sigma(y - y_x)^2)^{1/2}}{n - 2}$ .

Meinardus (1906) found April and May to be the peak months of the ice season. Ice was sighted most frequently and prevailed longer in those months than in any other. The average of April and May has been shown separately in Figure 3.

The source material is subject to errors arising from differences in observers' experience, size (height) of reporting vessel, faulty positioning of reporting vessel<sup>5</sup> and particularly from occasional paucity of observations<sup>6</sup> necessitating extensive extrapolations in drawing the charts.

In the present study, however, extreme precision in the relative magnitude of an ice season is of less significance than a clear indication of a change from light to severe ice conditions, or vice

---

<sup>5</sup> Prior to the 1920s the observing vessels lacked wireless, a circumstance severely impairing longitude determination.

<sup>6</sup> Decline of seal hunting early in the century was accompanied by a corresponding reduction in ice reports from some crucial areas north of Jan Mayen.

versa, in consecutive years. Several such sequences are readily identifiable from the graph and are used in the following.



### III. PREVIOUS WORK

Although the explorations of the 18th and 19th centuries were motivated largely by a quest for geographical discovery, the task of gathering scientific knowledge early became an important adjunct to the aims of the explorers. Scoresby thus gave account in 1820 of ice drift on the East Greenland Current, and Beechey followed with a discussion of that current in 1843. Drifting ice off Iceland was mentioned by Irminger in 1861, and at about the same time we find the first reference, by Lyell,<sup>7</sup> to the ice drifts' influence and periodicity. Subsequent work on the ice fluctuations can be classified into one of two categories labelled cause and effect, respectively. The latter is extensive, embracing work of numerous investigators,<sup>8</sup> whereas the former is quite limited, though of primary concern here.

The original and classic work on ice fluctuations per se is a

---

<sup>7</sup>"It is a well-known fact that every four or five years a large number of icebergs, floating down from Greenland, double Cape Langaness, and are stranded on the west coast of Iceland. The inhabitants are then aware that their crops of hay will fail, in consequence of fogs which are generated almost incessantly; and the dearth of food is not confined to the land, for the temperature of the water is so changed that the fish entirely desert the coast." (Lyell, 1866)

<sup>8</sup>With no pretense of a complete bibliography, some of the major works were published by Hann (1904), Brennecke (1904), Knudsen (1905), Helland Hansen (1920), Wiese (1924), Brooks and Quennell (1928), Nansen (1928), Petterson (1929), and Schell (1955).

paper by Thoroddsen from 1884. After researching original Icelandic sources, Thoroddsen described years of severe ice conditions off the Icelandic coasts in some detail, covering the period from 1233 to 1883.

In 1906, Meinardus published his previously mentioned study of ice conditions in the 19th century, in which he showed peak occurrences to be repetitive with a period of 4-1/2 years. He believed sunspots to be a cause and endeavored to show a relationship with an 11.1-year sunspot cycle.

Of more recent date is an interesting study by Hansen and Sveistrup (1943) of Arctic ice fluctuations in Julianehaab Bay during the years 1901-37. Julianehaab is located some 90 miles northwest of Cape Farewell, and in different years Arctic ice in varying quantities round the Cape and accumulate along the Greenland coast in the Labrador Sea. The two workers used this ice cover west of the Cape as their index and looked into its correlations with surface water temperatures, air temperatures and winds. The surface temperature data derived from sundry merchant ships' reports from the area between the Hebrides and the Faeroes. Air temperatures were taken from Cape North in Norway, Svalbard, Jan Mayen and Bear Island, and the winds from Grimsey and Angmagssalik. Hansen and Sveistrup found correlations of some significance and concluded that the fluctuations have no simple cause but are due to

the interaction of a number of factors.

A detailed treatment of the East Greenland ice was published by Koch in 1945, concluding a lifetime of field work in the region. Koch discussed at length the manifestations and possible reasons for the long-term fluctuations, which caused the demise of the Norse culture in Greenland in the 15th or early 16th century. The short-term behavior of the ice during the period 1898-1938 was also described with great thoroughness and superbly documented, but no short term causes were sought. Meinardus is thus the last scientist to concern himself with the causes for the ice fluctuations. In the more than six decades since his work was published, research on the subject shows a void.

Much effort has been expended, however, on collection of basic data which in the following are brought to bear on the Greenland Sea's ice phenomena. It is therefore appropriate to discuss some of their sources. In the oceanographic field, the Fram voyage in 1893-96 marked the starting point of modern systematic investigations in the Arctic, and since Nansen's successful return numerous expeditions, both single-ship missions and more elaborate land-sea

efforts, have brought back hydrographic data.<sup>9</sup>

Regular collection and issuance of weather data began with the publication of the Danish annual Meteorologisk Aarbog in 1873, which lists observations from several Greenland stations. In Norway, Jahrbuch des norwegischen meteorologischen Instituts commenced in 1874 to provide data from points in Norway, adding Spitzbergen in 1915, Jan Mayen in 1922 and Bear Island in 1928. Icelandic weather data have been published in Veðráttan since 1920, when this task was transferred from the Danish to the Icelandic meteorological service. At the same time, the reporting was expanded to comprise many additional stations in Iceland proper and on offshore islands. From the Shetlands, weather data have been listed in the British Weekly Weather Report published by the Meteorological Office in London since 1892.

Centralized ice reporting was organized by the Danish Meteorological Institute in 1896, and the task was formally requested

---

<sup>9</sup> A complete enumeration of these efforts is difficult, as many Danish investigations of the Greenland regions touch upon hydrography incidentally and peripherally, but the following post-Fram ventures must be included: Nathorst expedition (1898 and 1899); Gjøa (1901); Michael Sars (1901-04); Belgica (1905); Danmark Ekspeditionen (1906-08); Alabama (1909-12); Conrad Holmboe (1923); Godthaab (1924, 1930); Øst (1929); Meteor (1928, 1929, 1933, 1935); Tre Aars Ekspeditionen (Danish, 1931-34); Polarbjørn (1931-32); Polaris (1932); Heimland I (1933); North Pole (1937-39); Atha (1954); Scotia (1954); Dana (1955-59); Ob (1956); Johan Hjort (1958, 1961-62); Conrad Holt (1960); Edisto (1961, 1964, 1965); Westwind (1963).

from the institute by the VIIth International Geographical Congress in Berlin in 1899. The reports were issued uniformly until 1956. Since 1959, air reconnaissance has furnished the data by photography and radar, and the report format was changed. Supplementary reports on sea ice in Denmark Strait have been printed since 1953 in Jökull, sponsored by the National Research Council in Reykjavik. The Icelandic reports are compiled from sightings by coastal vessels and from aerial photographs.

#### IV. SOURCES AND PROCESSING OF DATA

Evaluation of the sundry factors influencing the ice cover has been made after gathering all such data as might contribute to the complete picture. The collection has been unevenly distributed, as more data can be found on some items than on others. Moreover, after preliminary investigation revealed precipitation and evaporation to be relatively insignificant, further efforts were directed primarily toward enriching the material pertaining to appraisal of the more dominant causes.

East Greenland Current data have been reconstructed from hydrographic material preserved in old expedition reports, notably from the East Greenland, Belgica, Danmark, Godthaab, Tre Aars and Meteor expeditions. These records in some cases contain original lists of stations and hydrocasts, and Table 1 contains the stations from which data were used. As the original records were not compiled to serve the purpose of this study, only a modest part has been useable, but it sufficed to construct four hydrosections across the current at 74°-76° N.

Tracing of water movements around Iceland and in the Faeroe-Shetland channel was made possible largely through the cooperation of the staff at the Oceanographic Data Center in Washington, D. C., who supplied magnetic tape records of all hydrographic data in their

Table 1. Stations used for hydrosections across the East Greenland Current.

Expedition or Ship	Year	Station	N. Latitude	W. Longitude	Depth m
E. Greenland	1891	IX	75°57'	4°15'	>1800
"	"	X	75°37'	6°40'	>1800
"	"	XI	75°06'	10°29'	>1800
"	"	XII	74°45'	11°42'	>1800
"	"	XIII	74°17'	15°20'	228
"	"	XIV	74°07'	17°30'	185
Belgica	1905	25	76°38'	3°40'	3240
"	"	26	76°31'	4°59'	3262
"	"	27	75°53'	8°33'	1290
"	"	28	75°49'	9°11'	1296
"	"	29a	75°34'	10°28'	1255
"	"	29b	75°39'	11°16'	347
"	"	30	75°40'	12°03'	366
"	"	31a	75°44'	13°00'	354
"	"	32	75°56'	14°00'	327
"	"	33	76°31'	14°42'	195
"	"	35	76°37'	15°00'	205
Danmark	1906	3	74°45'	3°23'	2080
"	"	4	75°27'	10°00'	2440
"	"	5	75°43'	10°15'	840
"	"	6	75°54'	10°00'	823
"	"	7	76°00'	14°34'	310
"	"	8	76°05'	16°04'	170
"	"	9	76°10'	16°44'	83
"	1908	78	76°06'	15°46'	180
"	"	79	75°57'	14°44'	300
Godthaab	1930	1	74°20'	14°56'	351
"	"	2	74°18'	15°17'	302
"	"	3	74°26'	16°22'	249
"	"	4	74°26'	17°01'	177
"	"	5	74°21'	19°26'	146

files from Marsden Squares 217, 218, 220, 254 and 288. This yielded 61,542 temperature and salinity data from the upper 100 m at approximately 840 stations. These stations have not been individually identified. All of the station data were instead categorized by one-degree unit areas, each Marsden Square being subdivided into 100 such unit areas. Programming and data processing were accomplished in Oregon State University's computer center, which provided computation by unit areas, month and year of means, ranges and standard deviations. These parameters in turn permitted interpretation of the time dependence of flow intensities and location of high flow intensities.

Seasonal variations in Denmark Strait were computed from the Data Center's tapes, and additional information was obtained from the records of the Dana and Øst stations listed in Table 2. Surface isotherms, isohalines and ice limits were drawn from the latter data group, and the positions at different times of the front separating Polar Water and Atlantic Water were determined.

A body of sea surface temperatures at Selvogsbanki on Iceland's south coast, 504 monthly values from the years 1895-1936, was obtained from a study by Thomsen (1937), who in turn had compiled them from ships' logs in the Copenhagen files of the International Association for the Exploration of the Sea. The data have been used here to correlate means of the months February-May with the ice



Table 2. Stations in Denmark Strait from which data were used.

Expedition or Ship	Station	Year	Expedition or Ship	Station	Year
Dana	4216	1931	Dana	4655	1933
"	4218	"	"	4656	"
"	4219	"	"	4657	"
"	4221	"	"	4658	"
"	4223	"	"	4659	"
"	4224	"	"	4660	"
"	4226	"	"	4662	"
"	4227	"	"	4665	"
"	4228	"	"	4666	"
"	4229	"	Øst	12	1929
"	4230	"	"	14	"
Dana	4429	1932	"	15	"
"	4430	"	"	17	"
"	4424	"	"	18	"
"	4432	"	"	19	"
"	4433	"	"	20	"
"	4434	"	"	22	"
"	4435	"	"	23	"
"	4436	"	"	24	"
"	4437	"	"	25	"
"	4438	"	"	27	"
"	4441	"	"	28	"
"	4442	"	"	30	"
"	4444	"	"	31	"
"	4445	"	"	34	"
"	4447	"	"	35	"
"	4448	"	"	36	"
"	4449	"	"	37	"
"	4450	"			

cover. This has been done by subtracting each annual February-May mean from the total February-May mean of the period 1900-36. The correlation coefficient between the ensuing anomaly and the ice cover was then computed.

In the Faeroe-Shetland channel data were obtained from the stations listed in Table 3. The station values were used to compute manually by planimetry a time series of heat transport through those parts of the sections Nolsø-Flugga and Faeroe Bank-Butt of Lewis which straddle the meandering current. Anomalies by month and year were determined in 17 years of the 1927-52 period, and annual volumes of flow were inferred from the results. A correlation between flow and ice cover was then established.

Table 3. Stations in the Faeroe-Shetland channel from which data have been used. The stations are only identified by location and have been variously occupied by Dana, Explorer and Scotia.

N. Lat.	W. Long.	Depth m	N. Lat.	W. Long.	Depth m
61°14'	2°40'	1142	59°31'	7°05'	1033
61°21'	3°10'	1442	59°44'	7°15'	992
61°28'	3°42'	1254	59°56'	7°27'	613
61°35'	4°15'	670	60°12'	7°40'	622
61°42'	4°51'	238	60°24'	7°50'	805
61°49'	5°21'	175	60°35'	8°00'	878
58°50'	6°26'	119	60°45'	8°08'	695
59°00'	6°40'	148	60°54'	8°16'	176
59°17'	6°53'	567	61°08'	8°29'	97

The meteorological data derive from the stations listed in Table 4. Pressure distributions are determined from 5568 data summarized from 169,220 observations. They are used to find the trend of annual pressure averages at the stations where barometric pressure has been consistently recorded. The means of the first two decades of this century is compared with the means of the second two decades, and the differences are used to infer a displacement of the Icelandic low pressure cell.

Table 4. Meteorological stations from which weather data have been used.

Station	N. Latitude	Longitude	Nationality
Aberdeen	57°10'	2°96'W	British
Angmagssalik	65°37'	37°16'W	Danish
Bear Island	74°28'	19°37'E	Norwegian
Gjesvaer	71°06'	25°22'E	Norwegian
Grimsey	66°33'	18°01'W	Icelandic
Haparanda	65°50'	24°09'E	Swedish
Hopen	76°30'	25°04'E	Norwegian
Jan Mayen	70°59'	8°18'W	Norwegian
Kinn	61°33'	4°48'E	Norwegian
Lerwick	60°09'	1°11'W	British
Røst	67°30'	12°04'E	Norwegian
Spitzbergen	78°02'	14°14'E	Norwegian
Stykkisholm	65°05'	22°46'W	Icelandic
Thorshavn	62°02'	6°44'W	Danish
Tromsø	69°39'	18°58'E	Norwegian
Vestmanneyjar	63°24'	20°17'W	Icelandic

Precipitation graphs are worked up from 6168 data, which are monthly values based on 187,832 observations of Danish, Icelandic and Norwegian origin. From the former data, annual anomalies are

calculated for a series of stations in and around the Greenland Sea. The sum of these anomalies is deemed to be a value representative of the ice region, and its correlation with the ice cover is examined, as is the effect of average wind force in the period 1914-37.

Air temperature work is based on 10,028 data, also monthly values derived from 220,920 observations. As in the case of precipitation, variation of annual temperature in the ice region is inferred from summation of anomalies calculated for selected stations. Spitzbergen's annual and January-March anomalies are calculated and graphed separately. Correlation coefficients between ice cover and temperature are determined by all three criteria.

The effect of wind on the formation and retention of the ice cover is sought by defining an index quantifying the influence in certain months each season when persistence of a particular component at a given station may affect ice formation. The basic material consists of 105,204 monthly data on wind direction covering 9,627,260 observations, and from 5928 monthly data on wind strength covering 1,808,040 observations. The index is established at each station by multiplying monthly wind force by the number of times the pertinent component was observed at that station. The correlations between the indices and the ice cover fluctuations are then calculated.

The search for supporting information has entailed review of a number of papers on the Arctic, which are listed in the bibliography.

This has been done not as a process of judging contents on their intrinsic merits but rather for the sake of relating their contents and conclusions to the task at hand, a purpose for which none of them was intended.

In total, the work has been a process of search for information and data on events long past, using them deductively and inferentially to gain knowledge of the causes for the ice variations. The material used has practically all been in the nature of basic data. The data have not been used to this effect before, nor does there seem any other way of gaining the knowledge.

## V. CURRENT EFFECTS

### The East Greenland Current

The surface waters of the Greenland Sea are formed by the confluence of water from the East Greenland Current through the Jan Mayen Current and the East Iceland Current with water from the North Atlantic Current, transported through the Irminger Current, the Norwegian Current and, northernmost, the West Spitzbergen Current, as shown in Figure 4. Most of the surface mixing takes place in two gyres, one located south of Jan Mayen and the other northeast of that island.

This current picture has been known in general outline since the first internationally coordinated survey was launched in 1902-04,<sup>10</sup> but recent observational programs, particularly during the IGY, have refined and somewhat modified several of its details. The picture thus emerging reveals the current structure at any given time to be both complex and ephemeral in the area under discussion.

A dearth of information makes Western oceanographers speak with caution about the East Greenland Current, but Treshnikov states

---

<sup>10</sup> This first attempt at a synoptic survey was planned at the International Oceanographic Congresses in Stockholm in 1899 and in Christiania in 1901.




Figure 4. Surface currents in the Greenland Sea.

— Atlantic Water  
 - - - Polar Water

the annual outflow between Greenland and Spitzbergen to be  $162,000 \text{ km}^3$ ,<sup>11</sup> or an average 5.1 Sv.<sup>12</sup> In 1962, Mosby summarized information from sundry sources in a general description of this current, but since then a profound and illuminating study has been published by Aagaard (1966).

It appears that in the northern part of the East Greenland Current Polar Water is found from the surface to a depth somewhat greater than 150 m. The temperature is between  $0^\circ \text{C}$  and the freezing point, and the salinity varies linearly from about 30‰ at the surface to slightly more than 34‰ at the bottom of the layer. Underlying the PW is the Atlantic Intermediate Water, advected through the current paths mentioned above and extending to a mean depth of 800 m. The temperature of this water mass remains above  $0^\circ \text{C}$  with a maximum evident between 200 m and 400 m, while the salinity increases downward until it reaches a value between 34.88‰ and 35.00‰. Below 400 m the salinity remains virtually constant at this value. Below 800 m Deep Water is found, but for present purposes its characteristics are of no consequence.

---

<sup>11</sup> Although Treshnikov speaks unequivocally, he does not disclose the exact nature or quantity of observations, making only the statement: "During the last eleven years Soviet high-latitude air expeditions and drifting stations have thoroughly explored the Arctic Basin, at a wide network of deep-water oceanographic stations" (Treshnikov, 1959).

<sup>12</sup> The Sverdrup unit is used here,  $1 \text{ Sv} = 10^6 \text{ m}^3/\text{sec}$ .



Calculation of current speeds and volume transports has been made by Aagaard (1966) who found typical speeds of 10-15 cm/sec. He further calculated the PW transport at 7.7 Sv, dominated by a barotropic mode, the AIW transport at 21.3 Sv and a deep water transport of 2.5 Sv. These figures add up to a rather startling 31.5 Sv total, which is an order of magnitude greater than previously believed,<sup>13</sup> but they are well-supported.

Hydrographic data from the passage at Nordøstrundingen are too scarce to give any indication as to fluctuations in the PW outflow. On the basis of heat budget considerations Aagaard concluded that significant seasonal variations are likely.<sup>14</sup>

Lee (1962) has estimated West Spitzbergen Current transports at 74° N, using 11 years of hydrographic measurements, and found strong seasonal fluctuations in the upper 400 m with a maximum transport in February and a minimum in April. Timofeyev (1962) has reported on 13 years of data across the West Spitzbergen Current

---

<sup>13</sup> It should be noted that Aagaard's results exceed as stated such Western investigators as Vowinckel and Orvig (1962), who have estimated a total transport of 3.4-3.7 Sv. The PW far exceeds Mosby's estimate (1962) of about 2 Sv, but is fairly close to the above mentioned Russian figure of 5.1 Sv (Treshnikov, 1959).

<sup>14</sup> Chaplygin (1960) has stated that the East Greenland Current is of highest intensity in winter due to prevailing strong northerly winds, but he offers no evidence. Laktinov, Shamontev and Yanes (1960) are of the same view but also fail to provide substantiation.

at 78° N, finding a maximum in December-January and a minimum in May-June, the former being about twice that of the latter. These fluctuations may well contain a clue to East Greenland Current fluctuations, for several workers have claimed finding a coupling between the two streams; but unfortunately there is disagreement on whether or not the currents are in phase.<sup>15</sup>

The available hydrodata from the area between Nordøstrundingen and Spitzbergen do not allow an evaluation to be made of year-to-year variations, but a fairly sizeable body of material exists from the records of past expeditions in the area from 74° to 76° N, and further south. The station lines are shown in Figure 5, and the current in these latitudes crosses obliquely the Arlis II station line which was oriented 210° T; it probably closely parallels the 100 m isobath, giving a current direction of ca. 235° T. In Figure 6 hydrodata gleaned from the records of the Belgica and Danmark expeditions have been used to project hydrographic sections onto a plane normal to the current off Shannon Island.

Measuring the areas enclosed by the -1.0° isotherm shows the cross section at the time of Belgica's work in 1905 to be 53% larger than was the case in 1906-08, when the Danmark expedition made its

---

<sup>15</sup> Laktinov et al. (1960) believe the currents to be out of phase, counteracting each other, whereas Timofeyev (1962) has concluded from ice drift observations that they are in phase.




Figure 5. Location of station lines B-05, Belgica expedition 1905; D-06/08, Danmark expedition 1906-08; G-30, Godthaab expedition 1930; and EG-91/92, E. Greenland expedition 1891-92.




Figure 6. Positions of isotherms in July 1905 and August 1906-08. Sections are in same location, normal to East Greenland Current at  $75^{\circ}$ - $76^{\circ}$  N, projected from Belgica and Danmark expedition data.

hydrocasts. The former was in a period of increasing ice cover, the latter in a sequence of years of reversing trend, see Figure 3.

Another telling comparison can be made between the Godthaab section of July 1930 and the East Greenland expedition section of August 1891, Figure 7. These sections are in nearly the same location, in fact intersect each other at about  $16^{\circ}$  W. The difference in size of the PW flows is striking, as is the difference in ice severity: 1930 was one of the lightest ice years on record, whereas Meinardus (1906) reported that 1891 was one of only four years in the 19th century when the ice arrived at Iceland before Christmas.

It is also noticeable that the quantity of water of temperatures  $> 1.0^{\circ}$  C is greater when the PW flow is strong. This seems to indicate a general strengthening of the current with more entrained AIW eddying off the West Spitzbergen Current. The significance of this will be discussed later.

An item of interest is presented by a comparison drawn by Jakhellin (1936) between East Greenland Current volumes in 1905 and in 1931-32. Jakhellin calculated the former to be 1.6 Sv and the 1931-32 average to be 1.32 Sv. It will be seen from Figure 3 that the ice cover in those years deviated from the 1898-1913 mean by +7.2 and  $-14 (\times 10^4 \text{ km}^2)$ , respectively, showing agreement with the change in volume.

The Arlis II current measurements show that, generally, no




Figure 7. Positions of isotherms in August 1891 and July 1930. Sections are located about 74°N, based on Godthaab and East Greenland expedition data.

depth of negligible horizontal motion is found in the East Greenland Current. Although the hydro sections therefore are not translatable into meaningful velocities, it is clear from the foregoing that year-to-year variations in the East Greenland Current do exist.<sup>16</sup> That they are directly linked to the ice occurrence is also empirically indicated.

### The North Atlantic Current

After leaving the vicinity of the American coast, the North Atlantic Current delineates the Polar Front, variable in time and space except at 52° N, 30° W, where it seems permanently fixed over the breakthroughs in the Mid-Atlantic Ridge. Subtropical water, coursing through the current's fingers, maintain secondary fronts in the Norwegian and Greenland Seas and in Denmark Strait. This water of about 8° C and 35.3‰ is a dominant factor in the ice picture, and it is of interest to assess its flows.

---

<sup>16</sup> It is worth noting that the Hansa survivors in 1870 drifted from 75° N to Cape Farewell at an average rate of four miles per day, whereas Irminger reported that the same route was covered by shipwrecked whalers in 1777 at a progress of 11-12 miles per day. Finally, in 1937-38 the ice station NP-1 traveled from 82.8° N, 7.0° W to 70.7° N, 19.3° W, at a net progress of 3.3 miles daily. It is open to speculation whether the differences in drift speeds are due to the different tracks traveled, or to current velocity differences in different years; but these case histories merit recall, as they represent direct measurements.

The North Atlantic Current's potential influence may be placed in proper perspective by comparing the magnitude of a 20% variation in the heat flowing through the Faeroe-Shetland channel in a given month with the heat needed to melt a season's ice in the Greenland Sea. The former can be approximated by using a value of  $2.3 \text{ Sv}^{17}$  at  $10^\circ \text{C}$ , giving a value of  $4.6 \times 10^{12} \text{ cal/sec}$  of such a fluctuation, or  $1.2 \times 10^{19} \text{ cal/month}$ . The latter can be roughly estimated as that heat needed to melt  $500,000 \text{ km}^2$  of ice one meter thick, or  $4 \times 10^{19} \text{ cal}$ . If the current and with it the heat import in fact do fluctuate from year to year, it is obvious that an influence on the melting of the ice cover must be detectable.

In the critical Iceland-Faeroe-Shetlands area, where it gains entrance to the Norwegian and Greenland Seas, the current has been the object of numerous explorations, most recently during the IGY and in the subsequent ICES investigation. Sufficient data have thus been collected on the current to gain several synoptic glimpses of its vagaries. As the core from examination of 68 hydrosections appears centered at about 100 m depth, this level will be analyzed.

It will be assumed that a cross-current temperature gradient exists wherever a major ocean current is located, so that a horizontal gradient will be a both necessary and sufficient reason to infer a

---

<sup>17</sup> This is a value estimated by Mosby (1962).



current's existence. Fuglister (1954) proposed this reasoning as a basis for current analysis, and it has been applied in the following. Between  $60^{\circ}$  and  $65^{\circ}$  N in the Atlantic, seasonal temperature variations by one-degree latitude belts have been computed by planimetry from Hydrographic Office charts of isotherms at the 100 m depth. The variations average  $0.69^{\circ}$  C in the five one-degree belts from  $60^{\circ}$  to  $65^{\circ}$  N, showing a temperature minimum in February and a maximum in August.

For purposes of comparison with this North Atlantic average, the seasonal variation in four one-degree units straddling the Faeroe-Shetland channel has been computed. The area comprising the four units is outlined in red in Figure 9, and the area's mean monthly temperatures have been plotted in Figure 8 together with their standard deviations. A curve representing the variations has been drawn, and it can be seen that the seasonal variation in this area is almost  $\pm 2^{\circ}$  C vs. the  $0.69^{\circ}$  C North Atlantic average for the same latitude. If temperature is equated with current strength, the explanation can be sought in the different current intensities prevalent in different seasons: fall-winter flows exceed spring-summer flows.

Petterson descriptively applied the word "Anschwellung" to the June-September intensities when analyzing data from the turn of the

century,<sup>18</sup> and the phenomenon is reliable, as shown by the small standard deviations in the autumn months.




Figure 8. Monthly temperature averages plus/minus one standard deviation for the area 60°-61° N, 3°-5° W and 61°-62° N, 5°-6° W (framed in red in Figure 8).

Certain longer term variations have been found in the surface layer by Brown and Rodewald (1953), who determined decreases of 0.3°-0.7° C in this area from 1910 to 1950. Rodewald (1966) also discovered cyclic surface temperature variations of the order of

<sup>18</sup>"Zwischen Ende Mai und Mitte September hatte eine mächtige Anschwellung des atlantischen Wassers stattgefunden, begleitet von einer Zunahme des Salzgehalts und der Temperatur" (Pettersen, 1929).

several years, but in the area in question they are well below  $0.5^{\circ}\text{C}$ . A sizeable temperature range in a small area must therefore be due to the presence of a current-induced horizontal gradient in that area, including the effect of current meander.




Figure 9. Temperature ranges in Marsden Squares #217 and #218. Grey area  $> 5^{\circ}$ , hatched area  $2^{\circ} - 5^{\circ}$  range. Area of max currents framed in red. Area enclosed by (— — —) is Fuglister's  $5^{\circ}$  range. Hydrosection line is shown as (— · —).

The material used is from 4115 hydrographic station casts, everything the Oceanographic Data Center in Washington has on file from Marsden Squares 217 and 218. Using temperatures at the 100 m depth, the ranges of observed temperatures have been computed by one-degree unit areas such as  $60^{\circ}00'-60^{\circ}59' N$ ,  $0^{\circ}00'-0^{\circ}59' W$ . Each range has been found by subtracting the minimum observed temperature from the maximum observed, regardless of date of observation.

In Figure 9 the range  $2^{\circ}-5^{\circ}C$  has been hatched, and the range  $>5^{\circ}C$  is shown in grey. Fuglister's outline of the same  $>5^{\circ}C$  range is also shown, and it will be noticed that it is somewhat smaller. The reason must be primarily that he looked at the 200 m depth, whereas the core of the flow as previously mentioned appears centered at the 100 m level.

In order to locate more precisely the currents traversing the section Iceland-Faeroes-Shetlands shown by a heavy stipled line in Figure 9, the mean temperature, the temperature range and the temperatures' standard deviation have been computed for each of the one-degree unit areas the section traverses from Iceland to the Shetlands.

Plotted in Figure 10, the average temperature curve shows that the maximum gradients lie between unit areas centered on  $14^{\circ}30'$ ,  $13^{\circ}30'$  and  $12^{\circ}30' W$ ;  $11^{\circ}30'$  and  $10^{\circ}30' W$ ; and  $5^{\circ}30'$  and  $4^{\circ}30' W$ . Standard deviations are highest in the unit areas centered on  $13^{\circ}W$




Figure 10. Average temperature, maximum range of temperature, and standard deviation by one-degree unit areas covering Iceland-Faeroe-Shetland section, as shown in Figure 8.

and  $3^{\circ}\text{W}$ , indicating major currents to be located at these points. Perusal of the charts in Figures 4 and 9 reveals the current at  $13^{\circ}\text{W}$  to be the Irminger Current, whereas the North Atlantic Current is located between  $5^{\circ}$  and  $2^{\circ}\text{W}$ . The range curve shows high values narrowly peaked at  $13^{\circ}\text{W}$ , meaning that the Irminger Current is fixed in position here. On the other hand, high range values throughout the area  $5^{\circ}$  to  $2^{\circ}\text{W}$  show the North Atlantic Current to assert itself over this entire width, a conclusion which is further supported by the matching high standard deviations.

Knowing the location of the North Atlantic Current, one can evaluate its relative strengths in the years from which data exist, and also the relative magnitude of the heat carried by it into the Norwegian-Greenland Sea region. The current is an upper water phenomenon, defined in depth by the 550 m Wyville Thomson Ridge and breaking surface at times over a wide area. For present purposes estimates have been made of the heat transported in a cross section 100 m deep, 182 km wide and one centimeter thick. The upper 100 m layer may reasonably be expected to influence ice conditions. The 182 km width represents the distance from  $2^{\circ}30'$  to  $5^{\circ}30'\text{W}$  along the section line in Figure 9, which the high range values in Figure 10 showed to be the location of maximum flow.

Analyses have been made of 50 sections deriving from Explorer, Scotia and Dana data from the period 1927-52. The reasoning has

been applied that a strong current through the Faeroe-Shetland channel will manifest itself in a positive temperature anomaly, i. e. the temperature at any point in the stream will be higher than the North Atlantic mean for that latitude. For latitude belts  $60^{\circ}$ - $61^{\circ}$  N and  $61^{\circ}$ - $62^{\circ}$  N the average temperatures as computed from Hydrographic Office charts of 100 m isotherms are  $6.65^{\circ}$  and  $6.54^{\circ}$  C, respectively. Adding about  $2^{\circ}$  C to allow for seasonal variations as determined in Figure 8, a value between  $8^{\circ}$  and  $9^{\circ}$  C results. It can accordingly be stated that the  $9^{\circ}$  isotherm in the section under study must enclose an area of substantial flow.

In each of the 50 sections, 100 m deep, 182 km wide and one centimeter thick, the areas enclosed by the  $9^{\circ}$  and higher isotherms, where present, have been measured planimetrically and their heat contents calculated and summed. The results are listed in Table 5, zero values denoting cases where all temperatures were  $< 9^{\circ}$  C. The figures in parentheses below the heat values are the anomalies by month. Thus for example the mean of all July values has been subtracted from each July value, showing that in 1934 the section's heat content was 44 units below the average for the month. When contemplating Table 5, it should be borne in mind, that a high heat value indicates both the presence in the section of an above average heat content available for transport and that a stronger-than-average current may be inferred. The effect on heat import of a high heat

Table 5. Heat content in cal $\times 10^{10}$  enclosed by the  $9^{\circ}$  isotherm in the section, 182 km wide, 100 m deep and 1 cm thick traversing the peak current intensity in the Faeroe-Shetland channel. Monthly anomalies are listed in parentheses.

	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
1927							211 (-17)				
1929				224 (+159)	240 (+64)						
1931				68 (+3)							
1933					185 (+9)						
1934				59 (-6)		139 (-42)					
1935					192 (+16)						
1936					161 147 (-22)						
1937				48 100 (+9)							
1938				45 (-20)	165 (-11)	210 (+29)					
1939					213 140 (0)	220 (+39)					
1946							280 (+52)				
1947					263 153 (+32)	188 (+7)	235 (+7)				
1948				0 116 (-7)				220 (+1)			
1949		0 (-54)		0 (-65)				227 (+8)		121 (-48)	
1950		161 (+107)			162 (-14)	200 (+19)				161 (-8)	
1951				0 (-65)	130 (-46)	104 (-77)	185 (-43)	211 (-8)	311		195
1952	0	0 (-54)		0 100 82 (+26)	138 (-38)	193 198 (+15)				153 240 (+37)	



value in the section is exponential.

Without reading more into the figures than is reasonably warranted, it may be considered significant that the two largest monthly anomalies are in May-June of 1929, a year when the curve in Figure 3 was plunging toward one of the lightest ice years on record. If the years are classified into four categories of rising, peak, declining and low ice coverage, summation of the monthly anomalies in these categories appears as listed in Table 6. Since the peaks and lows are more pronounced in the years 1927-39, these have been summed separately, but in either case negative totals accrue in the first two categories, positive in the last two. It is interesting that the declining category is more strongly positive than is the low one. It may be explained by the delay of the warm water's arrival at the ice edge. At an average progress of 10 cm/sec, the trip from the channel to the gyre north of Jan Mayen would take just over six months, so that the effect would be evident largely in the succeeding season.

Turning for a moment to the salinity of the current, a paper by Smed (1943) deserves attention. From an analysis of the International Council's records from 1900 to 1940, Smed found a definite increase in surface salinity in several areas of the North Atlantic. For the crucial area between Iceland and Scotland, his results are shown in Figure 11. From 1902/17 to 1919/39, the increase amounted to 0.06‰. Except as a possible indicator of intensity of the flow of

Atlantic water, this salinity change is of little direct consequence in the ice picture; in this case, it lowers the freezing point by a mere  $0.003^{\circ}\text{C}$ . It is nevertheless worth noting that a minimum winter salinity occurred in 1918, the severest ice season of all the years under study here, whereas winter salinity had progressed to a maximum in 1930, an exceptionally light ice year. Smed felt that the onset of the salinity increase began about 1920,<sup>19</sup> and if the graph in Figures 3 and 13 are viewed with this in mind, the coincident decrease in ice cover is striking. Averaging over Smed's periods, the recession of the ice can be computed as  $1.3 \times 10^4 \text{ km}^2$  from 1920/17 to 1910/39.

Table 6. Summation of monthly anomalies in years with ice severity classifiable as rising, peak, declining or low.

	1927-1939	1927-1952
Rising	-14	-20
Peaks	-50	-258
Declining	+239	+215
Lows	+26	+180

<sup>19</sup>Smed stated: "The investigations. . . all point towards the probability of the increase in salinity having commenced about 1920."




Figure 11. Average annual surface salinity of the area  $58^{\circ}$ - $63^{\circ}$  N,  $10^{\circ}$ - $15^{\circ}$  W.

### The Irminger Current

Although the Irminger Current was discovered and described before any other in the Greenland Sea, data from which its behavior may be reconstructed are scanty. During the IGY investigations, Denmark Strait and Irminger Sea charts of geopotential surface topography were produced, revealing rather intricate flow patterns including a large cyclonic whorl southwest of Iceland and a bifurcation to the northwest. From the latter, one branch curves north around Iceland, while another probes westward to join the East Greenland Current. Together the two currents wind their way west

and south, traceable by a band of crowded isotherms as shown in Figure 12, which is constructed from Anton Dohrn data from the summer of 1955. The 500 m and 1000 m isobaths are also shown, as they mark the slope of a steep-sided trough occupying the central part of the strait, with a 650 m sill depth west of Iceland's Cape Horn. The mean ice limits for the months of April, July and August have been superimposed.




Figure 12. Surface isotherms (blue) delineating boundary between Atlantic Water of Irminger Current and Polar Water of East Greenland Current in Denmark Strait, based on Anton Dohrn data from June 1955.

- 8 — surface isotherm, in °C
- 500 — isobath, in meters
- ..... mean monthly ice limit

In the years 1929, 1931 and 1932 detailed ice information was collected in Denmark Strait by Øst and Dana, and a comparison can therefore be made of the ice borders in the months of July of those years. In Figure 13 the three borders have been portrayed, and it is evident that certain features are of a semi-permanent character. It is apparent that the ice border coincides with the position of the Polar Front and that hydrographic conditions creating the front in turn depend on local bathymmetry. The precise bathymmetric charts produced in the 1950s confirm this.<sup>20</sup>

The current-ice relationship in the strait can accordingly be described as follows. The Irminger Current's bifurcation point is marked by a northward bulge of the 33‰ and 34‰ isohalines and the 1°-6° isotherms, which carve a matching bay into the edge of the ice. The point is displaced in different years (see Figures 12 and 13), along a line crossing the sill in a NE-SW direction, and meanders are spawned from it as the western branch of the Irminger Current proceeds together with the East Greenland Current toward the southwest. The meanders pendulate about an axis lying near the edge of the continental shelf and move down-stream at about 1/10th the current velocity. In the process they mold the ice border like the teeth of a giant saw.

---

<sup>20</sup>Thomsen (1938) drew similar conclusions based on successive chartings of the ice border from June through August in 1929.




Figure 13. July ice limits and mean positions of 34‰ isohalines in the years 1929, 1931 and 1932.

— — — — — Ice limit  
34‰ isohaline

It is interesting that the ice recedes in successive summer months to lines which remain parallel to the shelf edge. This takes place under the combined influence of diminishing supply of ice from the north and the July-August swelling of the Atlantic Water flow. Existence of this annual variation has been verified in Figure 14 by using monthly mean temperatures from selected unit areas. Their differences from the annual mean provide the points of the curve, and the autumn temperature rise indicates the same characteristic swelling of the flow as was seen to exist in the Faeroe-Shetland section.




Figure 14. Seasonal variation of temperature at 100 m depth from annual mean in the eight degree unit areas shown.

Year-to-year variations in the position of the ice border in Denmark Strait can then be attributed at least in some measure to the Atlantic Water flow's attaining different strengths in different years. Proof of this is almost totally lacking, as the current in the Irminger Sea generally is very broad and sluggish, not at all amenable to the kind of analysis that was possible on the Faeroe-Shetland channel flow. An attempt at a possible correlation can nevertheless be made by looking at the surface temperatures, the one type of data which is on hand in sufficient quantity.

In a study of long-period trends, Thomsen (1937) compiled surface temperatures at Selvogsbanki in the area framed red in Figure 15 and obtained monthly means for the years 1895-1936. Drawing upon this data series, the mean of the February-May temperature




Figure 15. February-May surface temperature anomaly at Selvogsbanki and ice cover curve (from Figure 3).



anomalies is graphed in Figure 15 together with the ice cover curve for the same period. The correlation coefficient has been found to be -0.30, not an impressive value but with the proper sign. It is here assumed that Selvogsbanki surface water will reach the ice edge through the Irminger Current with a lag time of one month.

It is probably more illustrative to analyze the years in which the ice cover differed significantly from its mean value. This has been done in Table 7, where the surface temperature anomalies have been summed for those years in which ice cover deviated from its regression line by an amount greater than  $s$ . In this manner temperature anomalies totaling  $-2.68^{\circ}\text{C}$  accrue in six years of high (positive) ice cover values, whereas temperature anomalies totaling  $+2.55^{\circ}\text{C}$  accrue in six years distinguished by low ice cover values. An inverse relationship thus seems to exist between intensity of the Irminger Current as indicated by Selvogsbanki surface temperatures on the one hand, and extent of the ice cover on the other.

Table 7. Tabulation of years in which ice cover ( $x$ ) deviated from its mean by a value larger than  $s$  (Figure 3). The February-May surface temperature anomaly is listed under each year.

Years of $x > +s$	1906	1907	1917	1918	1934	Total	
Temperature Anomalies, °C	-.83	-.78	-1.60	+.15	+.38	-2.68	
Years of $x < -s$	1904	1925	1930	1931	1933	1936	Total
Temperature Anomalies, °C	+.35	-.10	+.50	+.30	+.53	+.97	+2.55

## VI. CLIMATIC EFFECTS

### Relevant Weather Stations

Testing the influence of climatic factors is aided by a body of weather information which is rather sizeable--at least if compared with Arctic oceanographic observations. In Figure 16 the stations are shown which have provided the basic data used in the following. Their operational records vary a great deal, Thorshavn, Lerwick and the Norwegian mainland stations being the only ones with practically unbroken records in this century. Observations from points in Iceland are fragmentary, and gaps also exist in those from Angmagssalik.

Spitzbergen was equipped with a complete weather station in 1914,<sup>21</sup> Jan Mayen in 1922, Bear Island in 1928 and Hopen not until 1946. Their records have been reliably continuous except for the WW-2 period.

Some stations have been moved slightly in the period under review. The Tromsø station was in 1927 moved to the town's Geofysiske Institut, changing altitude from 38 meters to 102 meters.

---

<sup>21</sup> A number of observational series mostly of 1- or 2-year duration were obtained from various localities on the island, the earliest dating back to 1894.




Figure 16. Locations of weather stations from which data were used in this study.

Røst data prior to 1907 derive from Skomvaer, some eight miles SE of the present location. The Spitzbergen station was in 1930 moved to Svalbard Radio from Green Harbour, which is 23 miles NE of the present spot. None of the moves should have any measurable effect on the comparative value of the observations as they are used here. Preference is given, where possible, to island stations well off the nearest mainland, so that data sources on the fringe of the area of interest represent oceanic rather than terrestrial climates.

#### Weyl's Precipitation Hypothesis

At this point, it is pertinent to revert momentarily to the hypothesis proposed by Weyl (1966), to wit, that severe ice conditions must be expected after quiet rainy summer seasons. Angmagssalik, Grimsey, Jan Mayen, Spitzbergen, Bear Island and Tromsø should together give a good indication of rainfall in the Greenland Sea. In Figure 17 the precipitation anomalies from these points have been summed by year for the period 1900-39. The curve is assumed to be composed of four station values; when fewer station values are available in a particular year, the sum has been factored up; when more, down.

It is apparent that the curve correlates poorly with the ice cover curve shown above it. If the correlation coefficient




Figure 17. Precipitation anomalies of Angmagssalik, Grimsey, Jan Mayen, Spitzbergen, Bear Island and Tromsø summed by year and juxtaposed with ice cover graph (from Figure 3). Below is graph of mean wind for the period July-October from the same stations.

$$r = \frac{SP}{(SS_x SS_y)^{1/2}}$$

is computed with x as the ice cover's deviation from the 1898-1913 mean in km<sup>2</sup> and y as the cumulative precipitation anomaly in mm as graphed, it is found that  $r = 0.08$ . Hence precipitation per se is of no significant consequence.

The wind strength will now be taken into account as suggested by Weyl by summing from the same stations the mean winds each year for the period July-October, i. e. the four months preceding onset of ice formation. This curve is also graphed in Figure 17, but only from 1914, when the Spitzbergen station started operating, until 1936, when the Norwegian meteorological service ceased reporting wind force and switched to a different reporting criterion.

It is noticeable first of all that the winds are remarkably even in force from year to year, so that one would hesitate to classify any year as decidedly quiet. Secondly, the combination of precipitation and comparative calm is almost non-existent: wind and rain go together. If nevertheless all of the years showing below-average wind force are accepted as calm, it can be seen that two of those years experienced above-average precipitation, namely 1921 and 1924. The effects should have been felt in 1922 and 1925, respectively, but the ice cover was below average in 1922 and very low in 1925.

It is perhaps conceivable that even below-average annual rainfall could suffice for rapid ice formation, if only the fresh water layer were left undisturbed on the surface. In other words, storm activity, or the absence thereof, might be the only governing factor in regard to the precipitation effect. With this in mind the correlation coefficient between wind force and ice cover as portrayed in Figure 17 has been calculated. It turns out that for the 1914-36 period  $r = 0.24$ , an insignificantly small coefficient, and anyway positive, which is contrary to the requirement of Weyl's hypothesis.

All hope of forecasting ice seasons by the handy means of rain and wind reports should thus regrettably be abandoned. Although the effect of course must exist, as pointed out by Weyl, it is too slight to influence the ice regime in a detectable manner.

### Evaporation

Despite scarcity and unreliability of surface evaporation figures from high latitudes, it is possible to establish that variations in heat withdrawal from the ocean surface due to varying evaporation in successive years can exert but negligible influence on the ice cover. The area from which evaporative cooling could be brought to bear is at most  $800,000 \text{ km}^2$ . Assuming 60 cm of annual evaporation, a total of  $4 \times 10^{19}$  cal would be a maximum heat withdrawal.

If a 10% variation in this evaporative heat loss were to occur in



a given year, the amount would be two orders of magnitude below the annual heat import through the Faeroe-Shetland channel as computed earlier.

Change in surface salinity from a 10% variation in evaporation would amount to 0.2‰, changing the freezing point by 0.01° C, if the surface water is uniformly mixed to a depth of 10 m. The change in heat import needed to melt  $10^5 \text{ km}^2$  of ice one meter thick would be only  $10^{15}$  cal, an amount six orders of magnitude below the annual heat import through the Faeroe-Shetland channel. It would seem safe to disregard evaporation as a factor for present purposes.

### Temperature Variations

The gradual climatic warming in northern Europe and the Arctic in the 19th and 20th centuries has received much attention (Ångström, 1939; Scherhag, 1939; Hesselberg, 1940; Eriksson, 1943; Hovmöller, 1947), and a strengthening of the atmospheric circulation is generally surmised to be the cause. The trend is particularly pronounced in the Greenland Sea. Hesselberg (1940) has called attention to the temperature increase at Spitzbergen, where the annual mean has gained 2.5° C from 1899/1908 to 1929/38. Breaking down the increase, it is found that the temperature has gone up by 5.0° C in the winter and only by 0.2° C in the summer, a circumstance Hesselberg ascribes to heat consumption by melting of ice in summer,

which keeps the air temperature near the freezing point.

In regard to the shorter term fluctuations, it is reasonable to expect air temperatures in the Greenland-Norwegian Sea areas to be strongly dependent on the ice regime at any time. In Figure 18 a curve has been drawn of the combined temperature anomaly of four stations in the area, so that the air temperature's relation to the ice cover may be observed. The stations are Grimsey, Spitzbergen, Røst and Thorshavn, of which the first two are in the fringe of the ice, whereas Røst and Thorshavn are separated from it by 500-600 miles of open water.

The correlation coefficient between the combined anomaly curve and the ice is  $-0.512$ . The Grimsey-Spitzbergen anomaly has been drawn separately, and as one would expect its correlation is somewhat better at  $-0.682$ .

The question presents itself whether the correlation varies seasonally, and a 20-year series of monthly temperature means from Spitzbergen has been used to provide a partial answer. It appears that the January-March anomaly relates closely to the ice cover, as the coefficient jumps significantly to  $-0.818$ . On the other hand, calculation of Spitzbergen's May-August means and anomalies turns up a coefficient of only  $-0.292$  when related to the ice cover. Both curves are shown in Figure 18. This casts some doubt upon Hesselberg's argument referred to above. If ice melting so severely




Figure 18. Air temperature anomalies of Spitzbergen and of Grimsey-Spitzbergen-Røst-Thorshavn combined. Spitzbergen's January-March air temperature anomaly is shown separately below.

depresses the summer temperatures, one would expect a rather closer correlation between summer means and ice cover, the more so since ice cover here is defined as that of May-August, the ice melting period.

In Figure 19 both the ice cover graph and the 4-station anomaly have been smoothed by using 5-year moving averages. In an attempt to obtain an indication of the causality, the temperature anomaly has been compared also with previous and subsequent year's ice cover. The results are listed below the graphs, and although the latter correlation is slightly better, it can be said that the temperature anomaly neither manifests itself as a lingering effect of the ice regime nor does it presage the ice variations to any marked degree.

#### Wind-Current Relationships

Weather over the Greenland Sea depends on the momentary location of the atmospheric Arctic front and its generative low pressure cell, the extensive cyclone known as the Icelandic low. Rodewald (1966) has charted the mean pressure distribution in the area, and his findings have been used to show the front's location in Figure 20.




Figure 19. 5-year moving averages of ice cover curve (Figure 3) and annual combined temperature anomaly of stations Grimsey, Spitzbergen, Røst and Thorshavn.

It is of interest to see whether this position has remained statistically fixed during the period in which the ice cover has displayed a decreasing trend, and to this end the pressure changes registered at 11 stations in the area have been calculated and listed in Table 8. Each change has been found as the net difference between the mean pressure in the 1900-19 period and that in the 1920-39 period. The pressure changes have been noted in Figure 20 and they indicate that in the first four decades of this century the Icelandic low has shifted northward, particularly between Spitzbergen and Norway, and that it has become somewhat stronger.




Figure 20. Mean pressure distribution in the European Arctic (after Rodewald). Pressure change from 1900-19 period to 1920-39 period is shown below each station.

Table 8. Change in mean atmospheric pressure at selected stations from the period 1900-19 to 1920-39.

Station	$\bar{P}$ 1900-19	$\bar{P}$ 1920-39	Change	Remarks
Aberdeen	1011.1	1011.1	$\pm 0$	
Angmagssalik	1007.6	1006.9	-0.7	1910-11, 1924 missing
Gjesvaer	1007.0	1007.3	+0.3	1927-39 values from Sletnes
Haparanda	1009.5	1009.5	$\pm 0$	
Kinn	1010.0	1008.5	-1.5	1921-22, 1932-39 missing
Røst	1006.8	1006.8	$\pm 0$	
Spitzbergen	1009.2	1008.3	-0.9	1910-11, 1930 1938-39 missing
Stykkisholm	1006.2	1005.4	-0.8	
Thorshavn	1008.4	1008.3	-0.1	
Tromsø	1003.6	1006.0	+2.4	
Vestmannaeyjar	1005.3	1004.9	-0.4	

To verify this strengthening of the cyclone, a measure of the steepness of the pressure gradient in the key area of the Faeroes can be developed by computing the difference between annual means of atmospheric pressure at Thorshavn and Stykkisholm. This difference is shown in Figure 21, and a slight gradual steepening of the gradient is noticeable from 1900 to 1939, as the difference has increased by 0.195 mb per decade.

Northward displacement of the Arctic front will cause depressions to pass north of Spitzbergen, strengthening the southerly wind

component at Bear Island and Nordøstrundingen. Southward displacement will cause depressions to follow a more southerly track from the Faeroes toward the Baltic, strengthening the southerly component over the Faeroe-Shetland Channel. The wind regime is dominated by these movements of the front and by the increase in atmospheric circulation resulting from the slight strengthening of the low pressure cell. This should be borne in mind in examining the influence on the ice formation, which the variations in prevailing winds can exert by water mass transport.




Figure 21. Difference in annual mean pressure between Thorshavn and Stykkisholm. The regression line, calculated by the method of least squares, indicates a steepening of the gradient amounting to 0.195 mb per decade.

The water mass being created in the gyre northeast of Jan Mayen is the product of admixture of diluted Atlantic Water from the West Spitzbergen Current with Polar Water of the East Greenland



Current. South of Jan Mayen, Polar Water of the Jan Mayen Current mixes with Atlantic Water from the Norwegian Current and possibly with some of the Irminger Current. As we are dealing with surface currents, the wind regime in a given season will influence both directions and intensities of these flows and hence temperature and salinity of the water in the area of ice formation. The importance of the proportion of Atlantic Water advected is obvious.

If the winds at Jan Mayen are between south and west, while those of Tromsø and Røst (see Figure 20) are between south and southwest, the Norwegian Current is accelerated toward the Barents Sea, and mass transport into the Spitzbergen branch of the system is not aided. If on the other hand winds between northeast and southeast predominate, mass transport is diverted toward Spitzbergen, increasing admixture of warmer water in the gyre northeast of Jan Mayen and possibly opposing the East Greenland Current at its exit from the Arctic Ocean.

By analogous reasoning, winds between northwest and northeast at Spitzbergen will increase the admixture, whereas winds between east and south will speed the current on into the Arctic Ocean.

If the cumulative effect of all the winds in a season is expressed quantitatively, its correlation with the ice fluctuations will yield information about the actual importance of the wind regime. Determination of such cumulative effect will be attempted. In order first

to quantify the influence of wind at a certain station, it is practical to define a wind effect index for winds of a given direction as  $WEI = \text{duration} \times \text{strength}$ , where duration is measured as number of observations<sup>22</sup> and strength is expressed as Beaufort force. If the Beaufort scale expresses wind velocity proportional to its force numbers,<sup>23</sup> and if the wind's frictional effect on the surface current is assumed proportional to wind velocity, we have a crude but meaningful measure to work with.

The WEI at a given station will comprise the wind component which tends to aid transport of Atlantic Water to the ice border. The effect of high WEI value should accordingly diminish the ice cover, whereas a low WEI value could be expected to enlarge it. In Table 9 are listed the wind components of which the WEI is composed at six stations. Correlations have been calculated between the WEI and the ice cover by full years and by periods which depend on the estimated travel time of a water parcel from the point of wind influence to the ice border, and which also depend on the season in which the advection of Atlantic Water can be expected to have a maximum effect on

---

<sup>22</sup> Most stations record three daily observations. Where a greater or smaller number is recorded, the result has been factored down or up to become comparable.

<sup>23</sup> This has been achieved in the following by counting wind strengths of Beaufort 3 and less as being of force 3, whereby the scale becomes roughly linear in terms of wind velocity.

Table 9. Correlation coefficients between Greenland Sea ice fluctuations and Wind Effect Indices (WEI) at selected stations in and near the Greenland Sea.

Station	WEI Component	Years used	Full Year Correlation	Period	Correlation
Bear Island	E	1928-39, 46, 48-57	+0.048	Jan-Apr	-0.196
Jan Mayen	E	1922-39, 46-57	+0.099	Jan-Jun	-0.076
Røst	E	1900-39, 46-57	-0.202	Oct-Mar	-0.460
Spitzbergen	N	1912-39, 46-57	+0.043	Oct-Mar	+0.089
Tromsø	E	1900-39, 46-57	+0.481	Oct-Mar	-0.610
Vestmanneyjar	E	1900-19, 24-39, 46-52	+0.238	Dec-May	-0.030

the ice. For example, at Tromsø the E component will aid mass transport toward the ice; the advection will influence ice formation from January through June; and the water's travel time is estimated at three months. Hence at Tromsø the E-component is summed over the period October-March.

The results from the six stations are listed in Table 9, and it will be noticed that the WEI from the periods of estimated maximum effect from Norwegian stations correlate well with the ice cover. At Spitzbergen there is no significant correlation, which can probably be explained by the northerly wind component's dual effects of impeding the outflow of water from the West Spitzbergen Current, shifting it instead into a path contiguous to the East Greenland Current, and of aiding the inflow of Polar Water and ice. This would account for the simultaneous presence of substantial amounts of both PW and diluted AW in the Belgica section in Figure 6. The Jan Mayen correlation is also poor, but the number of years from which data are available from this station is too short to accord them importance comparable to that of the Norwegian series.

The period WEI anomalies are in Table 10 summed over the years in which the ice cover deviated from its mean by an amount greater than  $s$ . The Norwegian stations again show significant results, confirming the expected relationship between wind effect and ice cover.

Table 10. Anomalies of Wind Effect Indices in years the ice cover  $x$  deviated from its mean by an amount greater than  $s$ .

Bear Island	$x > +s$	1905	1906	1917	1918	1934		Total
		--	--	--	--	-3.1		-3.1
	$x < -s$	1904	1925	1930	1931	1933	1936	Total
		--	--	-0.8	+0.1	0	+0.4	-0.3
Jan Mayen	$x > +s$	1905	1906	1917	1918	1934		Total
		--	--	--	--	-0.6		-0.6
	$x < -s$	1904	1925	1930	1931	1933	1936	
		--	+1.5	+2.7	-2.7	+1.2	-0.8	+1.9
Røst	$x > +s$	1905	1906	1917	1918	1934		Total
		-0.9	-0.8	-0.5	-0.5	-0.6		-3.3
	$x < -s$	1904	1925	1930	1931	1933	1936	
		-0.7	+0.8	+1.0	+0.3	-0.4	+1.4	+2.4
Spitzbergen	$x > +s$	1905	1906	1917	1918	1934		Total
		--	--	-0.3	-0.5	--		-0.8
	$x < -s$	1904	1925	1930	1931	1933	1936	
		--	-0.5	-0.4	-0.4	-0.5	-0.3	-2.1
Tromsø	$x > +s$	1905	1906	1917	1918	1934		Total
		-0.1	-0.3	-0.9	-0.7	0		-2.0
	$x < -s$	1904	1925	1930	1931	1933	1936	
		+1.2	+0.1	-0.2	+0.4	-0.2	+0.9	+2.2
Vestmanneyjar	$x > +s$	1905	1906	1917	1918	1934		Total
		--	--	--	--	-2.8		-2.8
	$x < -s$	1904	1925	1930	1931	1933	1936	
		--	-4.0	+0.2	-1.7	+2.0	--	-3.5

## VII. DISCUSSION

The task of evaluating from the foregoing findings the relative importance of the factors influencing the ice regime is complicated by the factors' simultaneous presence, by their variability, and by their having little or no apparent interdependence. It is practical to treat them in reverse order of their estimated importance, thus deleting some at the start.

Variations in surface salinity of the Atlantic water may be induced by local disturbances in the evaporation-precipitation balance equatorward in the Gulf Stream system, or their cause may lie in the varying degrees to which vertical and lateral mixing takes place in the surface layers; but it is evident that they exist as a seasonal occurrence, as a year-to-year phenomenon and as decadal trends. Their direct effects in lowering the freezing point are negligible, however. The  $0.06\text{‰}$  salinity increase from 1902-17 to 1919-39 would only register an effect four orders of magnitude below that of a 20% variation in one month's heat import through the Faeroe-Shetland channel. The salinity fluctuations can therefore be ignored.

Fluctuations in surface temperature as computed by Rodewald might be of greater importance than those of salinity by two orders of magnitude but are still insignificant. Besides, the data give no

indication that the fluctuations extend to deeper parts of the North Atlantic Current. Their influence on the ice can therefore also be considered below the threshold of detectability.

Precipitation falls in the same category as the above two factors. If an anomalous 100 mm of rain water were added at the ideal time to the sea surface in the area of ice formation, and if it were mixed to a depth of only 3 m, the effect would still be an order of magnitude below the effects of known current variations. Hence precipitation can be ignored. As shown in Chapter VI, evaporation is also insignificant.

It bears emphasizing that the above factors certainly all affect the ice regime. The laws of physics compel us to recognize that such effects exist in any case. But in the composite picture of numerous natural forces interacting, the noise level is too high to permit their detection.

Air temperature variations present an enigma. There is no questioning that a pronounced decrease in ice cover over the first four decades of this century has been accompanied by a strong increase in air temperatures of the ice region, and it would indeed have been astounding if such had not been the case. The relationship nevertheless is obscure when year-to-year variations are considered. The best that can be said at this juncture is that available evidence fails to indicate air temperature fluctuations as a cause of the ice

changes.

Wind influence is the most difficult factor to assess quantitatively. Despite extensive records from the perimeter and some shorter series from within the ice region, the effects of wind defy easy analysis. This is not merely due to the multifarious nature of wind's influence through evaporation, precipitation, heat exchange, mass transport, depth of the mixed layer and spreading or compaction of the ice, but it is particularly due to an opposition of effects on the ice by wind components in certain key locations. Some examples will serve to illustrate. A strong easterly component south of Iceland will oppose mass transport of Atlantic Water into the Norwegian Current; but it will simultaneously promote mass transport via the Irminger Current through Denmark Strait toward the ice. A strong southerly component between Iceland and the Faeroes will aid mass transport of Atlantic Water across the sills between the Shetlands and the Faeroes and between the Faeroes and Iceland; but it will at the same time oppose outflow of Polar Water in the Irminger Current encircling Iceland. A strong northerly component between Nordøstrundingen and Spitzbergen will transport both ice, Polar Water and Atlantic Water from the West Spitzbergen Current into Greenland Sea. Conversely, absence of compound influence may well explain the rather good correlation between the ice cover and the easterly wind component at Norwegian coastal stations. From



there, an east wind can but transport water northwestward, toward the ice.

Detailed synoptic wind data and much additional information about the precise current structures are needed from the areas of oppositional influence before a clear picture can be developed of the winds' effects. At present, the effects can only be termed verifiable, but not dominant.

It is realized, of course, that this discussion has been confined to the wind-induced mass transports within the Greenland-Norwegian Sea area. It has been practical to deal with fluctuations in the volume of water entering from the North Atlantic Current under the heading of current effects, although the unknown, distant causes for this phenomenon probably lie in winds of the lower latitudes.

Year-to-year fluctuations in the flows of Polar Water and Atlantic Water into the Greenland Sea remain as the primary cause for year-to-year ice variations. This is concluded first on available evidence: 1) as calculated from hydrosections, the currents transport such quantities of heat (the southern flows) and ice (the northern flow) that their variations have the physical capacity to influence the ice cover; 2) their intensities in fact do vary, both seasonally and from one year to another; and 3) a correlation has been shown between high intensity and ice cover, negative for the southern flows and positive in the case of the northern. Second, the conclusion is

supported on deductive grounds: the currents are the only factor analyzed which has the physical capacity to influence the ice cover.

In summary, the causes for the short-term ice fluctuations are to be sought in the relatively easily monitored flows of water and air into and out of the Greenland Sea. The much more complex air-sea energy exchanges and evaporation-precipitation relationships, difficult or impossible to measure synoptically, are unimportant by comparison. One can therefore anticipate the day when a suitable system of recording buoys together with shore-based installations will provide the information for reliable forecasts of ice conditions and their concomitant weather effects. By then, the bafflement at the ice phenomena noted by the nameless explorer seven centuries ago in Konungs Skuggsjá will finally vanish.

## BIBLIOGRAPHY

- Aagaard, Knut. 1966. The East Greenland Current north of Denmark Strait. Ph. D. thesis. Seattle, University of Washington. 122 numbered leaves.
- Alekseev, A. P. and B. V. Istoshin. 1960. Some recent results of oceanographic research in the Norwegian and Greenland Seas. Soviet Fisheries Investigations in North European Seas. Moscow. Transl. by Fisheries Laboratory, Lowestoft, Suffolk. 26 p.
- Amdrup, G. 1902. Skibsekspeditionen til Grønlands Østkyst. VII. Hydrografi. Meddelelser om Grønland 27:341-52.
- Ångström, Anders. 1939. Temperaturklimaets ändring i nuvarande tid och dess orsak. Ymer, 1939, p. 119-131.
- Beechey, F. W. 1843. A voyage of discovery towards the north pole, performed in His Majesty's ships Dorothea and Trent, under the command of Captain David Buchan, R.N. London, Bentley. 351 p.
- Brennecke, W. 1904. Beziehungen zwischen der Luftdruckverteilung und den Eisverhältnissen des Ostgrönländischen Meeres. Annalen der Hydrographie und Maritimen Meteorologie 32:49-62.
- Brooks, C. E. P. and W. A. Quennell. 1928. The influence of Arctic ice on the subsequent distribution of pressure over the eastern north Atlantic and western Europe. London, Air Ministry Meteorological Committee. 110 p. (Geophysical Memoirs no. 41)
- Brown, P. R. 1953. Climatic fluctuations in the Greenland and Norwegian Seas. Quarterly Journal of the Royal Meteorological Society 79:272-281.
- Chaplygin, I. E. 1960. O dinamike vod vostochno-grenlandskogo techeniya. Problems of the Arctic and Antarctic 5:15-19.
- Chavanne, Josef. 1875. Die Eisverhältnisse im arktischen Polarmeere und ihre periodischen Veränderungen. Geographie und Erforschung der Polar-Regionen 108:134-280.

- Conseil Permanent International pour l'Exploration de la Mer.  
Septembre, 1961. Rapports et Proces-Verbaux des Reunions.  
Vol. 149. Contributions to Special IGY Meeting 1959.  
Copenhagen, Andr. Fred. Høst & Fils.
- Defant, A. 1930. Bericht über die ozeanographische Untersuchungen  
des Vermessungsschiffes "Meteor" in der Dänemarksstrasse  
und in der Irmingersee. Physikalisch-Mathematische Klasse  
16:230-35.
- Det Danske Meteorologiske Institut. 1899-1957. Meteorologisk  
Aarbog, 2den Del: Grønland. Kjøbenhavn, Centraltrykkeriet.
- Det Norske Meteorologiske Institut. 1900-57. Jahrbuch. Kristiania,  
Grøndahl & Søn.
- Dietrich, G. 1961. On the relationship between the distribution of  
redfish and redfish larvae and the hydrographical conditions in  
the Irminger Sea. Conseil Permanent International pour  
l'Exploration de la Mer. Rapports et Proces Verbaux 150:124-  
39.
- Dietrich, G. 1963. General oceanography. New York, John Wiley  
& Sons, Inc. 588 p.
- Dietrich G. 1964. Oceanic polar front survey in the North Atlantic.  
Cambridge, The MIT Press. 595 p. (Research in Geophysics,  
Vol. 2)
- Dunbar, Moira. 1963. Some features of ice movement in the Arctic  
Basin. In: Proceedings of the Arctic Basin Symposium, October  
1962. Hershey, Pa., Tidewater Publishing Corp. 288 p.
- Eis um Island und Ost-Grønland. 1940. Oberkommando der  
Kriegsmarine, neubearbeitet von der Deutschen Seewarte.  
Hamburg-Altona, Hammerich & Lesser.
- England. Department of State. Meteorological Office. 1892-1957.  
Quarterly Summary of Weekly Weather Reports.
- Eriksson, B. E. 1943. Till kännedom om den nutida  
klimatändringen inom områdena kring nordligaste Atlanten.  
Geografiske Annaler 25:170-201. Stockholm.

- Ewing, Maurice and William L. Donn. 1956. A theory of ice ages. *Science* 123:1061-66.
- Ewing, Maurice and William L. Donn. 1958. A theory of ice ages. *Science* 127:1159-62.
- Eythorsson, Jon. 1953-1961. Report on sea ice off the Icelandic coasts. Jökull, Reykjavik.
- Fletcher, J. O. 1965. The heat budget of the Arctic Basin and its relation to climate. Santa Monica, California, The Rand Corporation. 179 p. (memo R-444-PR)
- Fletcher, J. O. 1966. Proceedings of the symposium on the Arctic heat budget and atmospheric circulation. Memo RM-5233-NSF. Santa Monica, California, The Rand Corporation. 567 p. (memo RM-5233-NSF)
- Fletcher, J. O., B. Keller and S. M. Olencoff. 1966. Soviet data on the Arctic heat budget and its climatic influence. Santa Monica, California, The Rand Corporation. 205 p. (memo RM-5003-PR)
- Frommeyer, M. 1928. Die Eisverhältnisse um Spitzbergen und ihre Beziehungen zu klimatischen Faktoren. *Annalen der Hydrographie und der maritimen Meteorologie* 98:114-16.
- Fuglister, F. C. 1954. Average temperature and salinity at a depth of 200 meters in the North Atlantic. *Tellus* 6:46-58.
- Fuglister, F. C. 1960. Atlantic Ocean atlas of temperature and salinity profiles and data from the International Geophysical Year of 1957-1958. Woods Hole, Mass., Woods Hole Oceanographic Institution. 209 p.
- Garde, V. 1899. Isforholdene i Farvandene Øst og Vest for Grønland 1899. Særtryk of det danske meteorologiske Instituts nautisk-meteorologiske Aarbog. Copenhagen. 210 p.
- Hamilton, R. A. and G. Rollitt. 1957. Climatological tables for the site of the expeditions base at Britannia Sø. British North Greenland Expedition 1952-54. *Meddelelser om Grønland*. 158:1-82.
- Hann, J. 1904. Die Anomalien des Witterung auf Island in dem Zeitraum 1851-1900 usw. *Sitzungs-Bericht Wiener Akten*. Bd. 113:1-35.

- Hansen, G. J. A. and P. P. Sveistrup. 1943. Arctic ice fluctuations in Julianehaab Bay 1901-1937. *Meddelelser om Grønland* 131: 1-36.
- Helland-Hansen, Bjørn and Fridtjof Nansen. 1920. *Temperaturschwankungen des Nordatlantischen Ozeans und in der Atmosphäre. Einleitende Studien über die Ursachen der klimatologischen Schwankungen.* *Smithsonian Miscellaneous Collections* 70(4):1-266.
- Henry, A. J., C. E. P. Brooks and W. Quennell on the influence of Arctic ice on subsequent distribution of pressure over eastern North Atlantic and western Europe. 1929. *Monthly Weather Review* 57(3):99-102.
- Hesselberg, Th. 1940. Klimavariasjoner i Norge i vår tid. *Naturen* 64:289-303.
- Hill, H. W. and A. J. Lee. 1958. The effect of wind on water transport in the region of the Bear Island Fishery. *Proceedings of the Royal Society, Series B*, 148:104-16.
- Hovmöller, E. 1947. Climate and weather over the coast-land of northeast Greenland and the adjacent sea. *Meddelelser om Grønland* 144:I-1-I-148.
- Irminger, E. 1856. The Arctic current around Greenland. *Journal of the Royal Geographical Society* 26:36-43.
- Irminger, E. 1861. Die Strömungen und das Eistreiben bei Island. *Zeitschrift für Allgemeine Erdkunde* 11:191-211.
- Jacobsen, J. P. 1943. The Atlantic current through the Faroe-Shetland channel and its influence on the hydrographical conditions in the northern part of the North Sea, the Norwegian Sea, and the Barents Sea. *Conseil Permanent International pour l'Exploration de la Mer*, 48.
- Jakhellin, A. 1936. Oceanographic investigation of east Greenland waters. *Norges Svalbard- og Ishavsundersøgelser* 67:1-34.
- Kiilerich, A. B. 1945. On the hydrography of the Greenland Sea. *Meddelelser om Grønland* 144(2):1-63.

- Knudsen, M. 1905. On the influence of the east Icelandic polar stream on the climatic changes of the Faeroes Isles, the Shetlands, and the north of Scotland. Conseil Permanent International pour l'Exploration de la Mer, Rapp. Proces-Verbaux 3:1-8.
- Koch, Lauge. 1945. The east Greenland ice. Meddelelser om Grønland 130(2):1-368.
- Krauss, Wolfgang. 1955. Zum System der Meeresströmungen in den höheren Breiten. Deutsche Hydrographische Zeitschrift 8(3):102-111.
- Laktinov, A. F., V. A. Shamontev and A. V. Yanes. 1960. Okeanograficheskii ocherk severnoi chasti Grenlandskogo morya. pp. 51-65. In: Sovjetskie rybokhozyaistvennye issledovaniya v moryakh Evropeiskogo severa. Moscow, VNIRO PINRO. 468 p.
- Larson, L. M. (transl.). 1917. The king's mirror (speculum regale - konungs skuggsjá). American-Scandinavian Foundation, New York. 388 p.
- Lee, A. J. 1962. The effect of the wind on water movements in the Norwegian and Greenland Seas. In: Proceedings of the Symposium on Mathematical-Hydrodynamical Methods of Physical Oceanography, Institute of Oceanography, University of Hamburg. p. 353-73.
- Lee, A. J. 1963. The hydrography of the European Arctic and Subarctic Seas. Oceanographic Marine Biological Annual Review 1:47-76.
- Lyell, Sir Charles. 1866. Principles of geology. Rev. ed. New York, D. Appleton & Co. 834 p.
- Malmgren, F. 1928. On the properties of sea ice. In: The Norwegian North Polar Expedition with the Maud, 1918-1925: Scientific Results. Vol. 1, No. 4. Bergen. p. 1-67.
- Meinardus, Wilhelm. April 1906. Periodische Schwankungen der Eistrift bei Island. Annalen der Hydrographie und Maritimen Meteorologie 34:148-162, 227-39, 278-85.

- Mikkelsen, E. 1922. Alabama-Ekspeditionen til Grønlands Nordøstkyst 1909-1912. Meddelelser om Grønland 52:1-395.
- Montgomery, R. B. 1940. Discussion of some theories on temperature variations in the north Atlantic Ocean and in the atmosphere. Monthly Weather Review, Vol. 39, sup., p. 52-57.
- Mosby, Hakon. 1937. Svalbard waters. Geofysiske Publikationer 12: 1-85.
- Mosby, Hakon. 1963. Water, salt and heat balance in the north Polar Sea. In: Proceedings of the Arctic Basin Symposium October 1962. Hershey, Pa., Tidewater Publishing Corp. p. 33-58.
- Nansen, Fridtjof. 1928. The oceanographic problems of the still unknown Arctic regions. A series of papers. New York, American Geographical Society. 112 p. (Special Publication no. 7)
- Nielsen, J. N. 1928. Greenland. In: The waters around Greenland, ed. by The Commission for the Direction of Geological and Geographical Investigations in Greenland. Copenhagen, C. A. Reitzel. p. 185-229.
- Petterson, Otto. 1929. Changes in the oceanic circulation and their climatic consequences. The Geographical Review 19(1):121-131.
- Rodewald, M. 1966. Abkühlungstrend im Oberflächenwasser des Nordatlantischen Ozeans. Umschau 23:777.
- Ryder, C. 1895. Den østgrønlandske Ekspedition udført i Aarene 1891-92. V. Hydrografi. Meddelelser om Grønland 17:189-279.
- Schell, I. I. 1940. Polar ice as a factor in seasonal weather. Monthly Weather Review, Vol. 39, sup., p. 27-51.
- Schell, I. I. 1955. Interrelations of Arctic ice with the atmosphere and the ocean in the north Atlantic-Arctic and adjacent areas. Journal of Meteorology 13:46-58.



- Scherhag, R. 1939. Die Erwärmung des Polargebiets. *Annalen der Hydrographie und der maritimen Meteorologie* 67:57-67.
- Schott, G. 1904. Über die Grenzen des Treibeises bei der Neufundlandbank sowie über eine Beziehung zwischen neufundländischen und ostgrönländischen Treibeises. *Annalen der Hydrographie und der maritimen Meteorologie* 32:305-309.
- Scoresby, W. 1820. An account of the Arctic regions, with a history and description of the northern whale-fishery. Vol. I. Edinburgh, Constable. 551 p.
- Sestoft, Ingolf. 1956. Færøernes Vejr - og Klimaforhold. Særtryk af Den Færøske Lods. 6th ed. København, Det Kongelige Søkort-Arkiv. 8 p.
- Shaw, Napier. 1942. Manual of meteorology. Vol. II. Cambridge, At The University Press. 317 p.
- Smed, Jens. 1943. Annual and seasonal variations in the salinity of the north Atlantic surface water. *Conseil Permanent International Pour l'Exploration de la Mer, Rapports et Proces-Verbaux des Réunions* 12:77-94.
- Smith, Edward H. 1931. The Marion Expedition to David Strait and Baffin Bay. Washington, D. C., U. S. Government Printing Office. 219 p. (U. S. Coast Guard Bulletin no. 19)
- Sverdrup, H. U., Martin W. Johnson and Richard H. Fleming. 1962. The oceans. 12th ed. Englewood Cliffs, N. J., Prentice-Hall, Inc. 1087 p.
- Tait, John B. 1956. Recent oceanographical investigations in the Faeroe-Shetland Channel. *Proceedings of the Royal Society of Edinburgh, Section A*, 14:239-89.
- Tait, John B. 1957. Hydrography of the Faeroe-Shetland Channel 1927-1952. Edinburgh, Her Majesty's Stationery Office. 309 p. (Scottish Home Department Marine Research no. 2)
- Thomsen, H. 1937. Variation of the surface temperature at Selvogsbanki, Iceland 1895-1935. *Conseil Permanent International Pour l'Exploration de la Mer. Extrait au Rapports et Proces-Verbaux* 150:1-311.

- Thomsen, H. 1938. Hydrography of Icelandic waters. The Zoology of Iceland. Vol. 1, part 4. Copenhagen, Levin & Munksgaard. 220 p.
- Thoroddsen, Th. 1884. Den grønlandske drifisen vid Island. Ymer 4:145-160.
- Timofeyev, V. T. July 1962. The movement of Atlantic water and heat into the Arctic Basin. Deep Sea Research 9:358-61.
- Transehe, N. A. 1928. The ice cover of the Arctic Sea with a genetic classification of sea ice. New York, American Geographical Society. 311 p. (Special Publication no. 7)
- Treshnikov, A. F. 1959. Oceanography of the Arctic Basin. In: International Oceanographic Congress, 1959. Preprints. Washington, D. C. p. 522-524.
- Trolle, A. 1913. Hydrographical observations from the Danmark Expedition. Meddelelser om Grønland 41:271-426.
- U. S. Navy Hydrographic Office. 1958. Mean Temperatures at 100 Meters. Washington, D. C. 84 p. (Publication no. 700, sec. II)
- U. S. Navy Hydrographic Office. 1958. Oceanographic Atlas of the Polar Seas. Washington, D. C. 139 p. (Publication no. 705)
- Veðráttan, Manaðaryfirlit Samin A Veðurstoffunni. 1921-1957. Reykjavik.
- Vowinckel, E. and S. Orvig. 1961. Water balance and heat flux of the Arctic Ocean. Toronto, McGill University. 35 p. (Publications in Meteorology, no. 44)
- Wagner, Artur. 1940. Klimaänderungen und Klimaschwankungen. Friedrich Vieweg & Sohn, Braunschweig. (De Wissenschaft Band 92)
- Weyl, Peter K. 1966. The role of the oceans in climatic change: a theory of the ice ages. Unpublished paper. Corvallis, Oregon, Oregon State University.
- Wiese, W. 1924. Polareis und atmosphärische Schwankungen. Geografiske Annaler 6:273-229.

World weather records. 1927-34. Washington, D. C. 12 Vols.  
(Smithsonian Miscellaneous Collection, Vol. 79-90)

Wüst, Georg. 1937. Recent conceptions of the nature of the gulf  
stream system and the nomenclature of its parts. Der Seewart  
10:359-67.