

AN ABSTRACT OF THE THESIS OF

Lois F. Alexander for the degree of Master of Science in
Wildlife Science presented on 31 March 1994.

Title: A Morphometric Analysis of Geographic Variation Within *Sorex monticolus*.

Redacted for privacy

Abstract approved: _____

Shrews previously recognized as *Sorex monticolus* were classified into two species (one with 14 subspecies, the other monotypic) on the basis of a morphometric analysis of 3610 individuals from throughout their range. *Sorex m. neomexicanus* has been recognized previously as a subspecies of *Sorex monticolus* but is recognized herein as a distinct species. This taxon occurs in the Sacramento and Capitan mountains of New Mexico. This region possibly acted as a boreal-forest refugium for *S. monticolus*-type shrews during the Pleistocene glaciation, and during the warmer interglacial period, after the most recent glaciation, the valleys became too arid for survival and these shrews survived in forested, montane regions of New Mexico. These mountains are sufficiently isolated from other mountainous regions in the state to reduce or eliminate gene flow between these populations of shrews. *S. monticolus* as defined herein exhibits relatively little morphometric variation. Even among nominate races, the only obvious morphometric variation is a north-south cline in greatest length of skull. There is a general

trend of increasing size from south to north. The southern subspecies restricted to isolated mountains (S. m. monticolus and S. m. parvidens) have the shortest skull lengths of all S. monticolus. The subspecies found in the northern coastal and insular areas of southeast Alaska and British Columbia (S. m. longicaudus, S. m. prevostensis, S. m. malitiosus, S. m. insularis, S. m. calvertensis, S. m. alascensis, and S. m. elassodon) have the longest skulls. S. m. setosus, S. m. isolatus, S. m. soperi, S. m. obscurus and S. m. shumaginensis all have skulls of intermediate length. Insular and coastal populations of S. monticolus have longer skulls than the S. monticolus that occupy the mainland. Shrews with long skulls that occur on the mainland (S. m. longicaudus and S. m. alascensis) also occur on some islands, and the mainland portion of their distributions are restricted to a narrow band along the coasts of Alaska and British Columbia. The southernmost subspecies of S. monticolus with short skulls are restricted to small montane islands. The morphometric variation among nominate races is sufficient to warrant continued separation at the subspecies level of all taxa except S. m. calvertensis and S. m. elassodon. Were it not for differences in pelage color, based on my morphometric analysis, S. m. calvertensis and S. m. elassodon should be synonymized.

A Morphometric Analysis of Geographic Variation
Within Sorex monticolus.

by

Lois F. Alexander

A THESIS

submitted to

Oregon State University

in partial fulfillment of
the requirements for the
degree of

Master of Science

Completed 31 March 1994

Commencement 12 June 1994

APPROVED:

Redacted for privacy

~~Professor of Wildlife Ecology in charge of major~~

Redacted for privacy

~~Interim Head of Department of Fisheries and Wildlife~~

Redacted for privacy

~~Dean of Graduate School~~

Date thesis is presented 31 March 1994

Typed by researcher for Lois F. Alexander

ACKNOWLEDGMENTS

I thank B. J. Verts for his encouragement and patience while serving as my major professor. R. A. Pimentel provided assistance with BIOSTAT. I thank P. M. Harris, C. W. Alexander, L. N. Carraway, R. A. Pimentel, W. P. Stephen, and D. Nagorsen for many valuable discussions, and David, Nancy, and Daphne Nagorsen for their hospitality. I also thank my parents for providing financial support during the years of this research.

For the loan of Sorex monticolus specimens, I thank the following curators: C. L. Pritchett, Brigham Young University, Life Sciences Museum (BYU); D. G. Huckaby, California State University, Long Beach, Ross Hardy Mammal Collection (CSLB); C. G. van Zyll de Jong, Canadian Museum of Nature (CMN); S. B. McLaren, Carnegie Museum of Natural History (CMNH); B. D. Patterson, Field Museum of Natural History (FMNH); R. D. Fisher, National Museum of Natural History (USNM); S. B. George and L. Barkley, Natural History Museum of Los Angeles County (LACM); B. J. Verts, Oregon State University (OSUFW); G. W. Hughes, Royal British Columbia Museum (RBCM); M. Engstrom, Royal Ontario Museum (ROM); P. Unitt, San Diego Natural History Museum (SDNHM); G. L. Kirkland, Jr., Shippensburg University Vertebrate Museum (SUVN); G. D. Baumgardner, Texas A&M University, Texas Cooperative Wildlife Collection (TCWC); J. A. Cook, University of Alaska Museum (UAM); W. Roberts, University of Alberta, Museum of Zoology (UAMZ); B. R. Stein, University of California, Berkeley, Museum of Vertebrate Zoology (MVZ); D. Blake, University of

Illinois, Museum of Natural History (UIMNH); P. Myers, University of Michigan, Museum of Zoology (UMMZ); W. L. Gannon, University of New Mexico, Museum of Southwestern Biology (MSB); E. A. Rickart, University of Utah, Museum of Natural History (UUMNH); and R. E. Johnson, Washington State University, Charles R. Conner Museum (CRCM).

For their help during museum visits, I thank J. R. Choate, Fort Hays State University (FHSU); R. D. Fisher, National Museum of Natural History (USNM); D. W. Nagorsen, Royal British Columbia Museum (RBCM); R. Cannings, University of British Columbia, Cowan Vertebrate Museum (UBC); B. R. Stein, University of California, Berkeley, Museum of Vertebrate Zoology (MVZ); R. M. Timm and T. Holmes, University of Kansas, Museum of Natural History (KU); G. Shugart and D. Paulson, University of Puget Sound, Museum of Natural History (PSM); M. S. Boyce, University of Wyoming (UW); and M. A. Bogan, U.S. Fish and Wildlife Service, Biological Surveys Collection/Fort Collins (BS/FC).

This research was supported in part by Oregon Agricultural Experiment Station project 902. This is Technical Paper No. 10448, Oregon Agricultural Experiment Station.

TABLE OF CONTENTS

INTRODUCTION	1
METHODS	8
RESULTS	14
SYSTEMATICS AND TAXONOMY	34
<u>Sorex neomexicanus</u> Bailey	34
<u>Sorex bairdi</u> Merriam	42
<u>Sorex monticolus</u> Merriam	46
<u>Sorex monticolus alascensis</u> Merriam	47
<u>Sorex monticolus calvertensis</u> Cowan	53
<u>Sorex monticolus elassodon</u> Osgood	56
<u>Sorex monticolus insularis</u> Cowan	60
<u>Sorex monticolus isolatus</u> Jackson	62
<u>Sorex monticolus longicaudus</u> Merriam	67
<u>Sorex monticolus malitiosus</u> Jackson	73
<u>Sorex monticolus monticolus</u> Merriam	74
<u>Sorex monticolus obscurus</u> Merriam	78
<u>Sorex monticolus parvidens</u> Jackson	107
<u>Sorex monticolus prevostensis</u> Osgood	108
<u>Sorex monticolus setosus</u> Elliot	110
<u>Sorex monticolus shumaginensis</u> Merriam	116
<u>Sorex monticolus soperi</u> Anderson and Rand	119
DISCUSSION	121
LITERATURE CITED	126

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1. Distribution of (a) 791 unique localities for specimens used to establish current taxonomy of <u>Sorex monticolus</u> , based on Carraway (1990), George and Smith (1991), Hall (1981), and Hennings and Hoffmann (1977) and (b) 1336 unique localities for <u>Sorex monticolus</u> specimens included in this morphometric analysis.	5
2. Camera lucida tracing of a <u>S. monticolus</u> (MSB 1944) illustrating skull and mandible dimensions measured.	9
3. Map of marginal-record polygons for 14 subspecies of <u>Sorex monticolus</u> used as a priori groups in the present morphometric analysis. Marginal records based on Hall (1981) and Hennings and Hoffmann (1977).	11
4. Means, 95% confidence intervals, and ranges of discriminant scores for two-group discriminant analyses with (a) <u>S. bairdi</u> and <u>S. monticolus</u> , (b) <u>S. m. neomexicanus</u> and the remaining subspecies of <u>S. monticolus</u> , and (c) <u>S. bairdi</u> and <u>S. m. setosus</u> with three characters.	15
5. Confidence ellipses (95%) for all 14 a priori <u>S. monticolus</u> groups and <u>S. bairdi</u> .	17
6. A dendrogram generated by a cluster analysis based on the generalized-distance matrix derived from the 15-group discriminant analysis for <u>S. bairdi</u> and <u>S. monticolus</u> .	18
7. Confidence ellipses (95%) for (a) the remaining 13 a priori <u>S. monticolus</u> groups, (b) <u>S. m. obscurus</u> and <u>S. m. obscuroides</u> , (c) <u>S. m. calvertensis</u> and <u>S. m. elassodon</u> , and (d) the 12 remaining groups after all specimens had been placed in appropriate groups.	22
8. Dendograms generated by cluster analyses based on the generalized-distance matrices derived from the multigroup discriminant analyses for (a) the 13 a priori groups, and (b) the 12 remaining groups with all specimens in groups.	23
9. Distribution of localities for <u>S. neomexicanus</u> specimens included in this analysis.	36
10. Camera lucida tracing of skull and mandible of <u>S. neomexicanus</u> (MSB 41081).	38

11. Confidence ellipses (95%) for S. bairdi, S. neomexicanus, and the remaining S. monticolus. 40
12. A dendrogram generated by a cluster analysis based on the generalized-distance matrix derived from the multigroup discriminant analysis for S. bairdi, S. neomexicanus, and the remaining S. monticolus. 41
13. Distribution of localities for (a) S. m. alascensis (squares) and S. m. shumaginensis (circles) specimens included in this analysis representing 63 and 52 unique localities, respectively. 49
14. Distribution of localities for S. m. calvertensis (triangles and solid arrows), S. m. insularis (squares and open arrow), S. m. elassodon (circles), and S. m. prevostensis (asterisks and hatched arrow) specimens included in this analysis representing 5, 3, 55, and 3 unique localities, respectively. 54
15. Distribution of localities for S. m. isolatus (squares) and S. m. setosus (circles) specimens included in this analysis representing 59 and 150 unique localities, respectively. 64
16. Distribution of localities for S. m. longicaudus (circles) and S. m. malitiosus (squares and solid arrow) specimens included in this analysis representing 57 and 2 unique localities, respectively. 69
17. Distribution of localities for S. m. monticolus (squares and solid arrows), S. m. obscurus (circles), and S. m. parvidens (triangles and open arrow) specimens included in this analysis representing 39, 5, and 794 unique localities, respectively. 76
18. Mean and range of greatest length of skull (mm) for all subspecies of Sorex monticolus and the approximate distribution of each subspecies demonstrating the trend of increasing length of the skull from south to north. 122

LIST OF TABLES

<u>Table</u>	<u>Page</u>
1. Mean $\pm$ <u>SE</u> , range, and CV of 18 cranial and mandibular characters (in mm) for all taxa included in analyses. Sample size (<u>n</u>) for each group in parentheses.	20
2. Results of two-group discriminant analyses for adjacent subspecies of <u>Sorex monticolus</u> used to diagnose specimens from localities between a priori polygons. All pairs of group centroids were significantly different (<u>P</u> < 0.05).	26

PREFACE

This thesis was prepared in the style of an Occasional Paper, The Museum of Natural History, The University of Kansas, in anticipation of submitting it for consideration for publication in that series.

A MORPHOMETRIC ANALYSIS OF GEOGRAPHIC VARIATION

WITHIN SOREX MONTICOLUS

INTRODUCTION

Sorex monticolus ranges from northern Alaska south to northern Mexico and from the Pacific Coast east to northwestern Saskatchewan and eastern Alberta, Montana, Wyoming, Colorado, and New Mexico. Trinomials have been applied to insular populations of coastal Alaska and Canada (including elassodon, malitiosus, prevostensis, calvertensis, insularis, and isolatus), and to relictual montane populations in southwestern United States and Mexico (monticolus, parvidens, and neomexicanus) and central Saskatchewan and western Manitoba (soperi). The mainland of North America includes one primary taxon, S. monticolus obscurus. S. m. setosus occurs in western Oregon, Washington, and British Columbia; S. m. shumaginensis, S. m. alascensis, and S. m. longicaudus occur in western Alaska, the south coast of Alaska, and the southeast coast of Alaska and coastal British Columbia, respectively. All of the isolated populations probably were derived from an obscurus-like ancestor in the Rocky Mountains (Hennings and Hoffmann, 1977). Findley (1955) also indicated that these were offshoots from the main Rocky Mountain population but he considered them all subspecies of S. vagrans. Findley (1955) included longiquus and obscuroides as separate subspecies (of S. vagrans) but Hennings and Hoffmann

(1977:15) classified these as "no more than a cluster of ecotypic variants of S. monticolus obscurus."

The nominate form, Sorex monticolus monticolus, was described by Merriam (1890) as Sorex monticolus. Without comment, Merriam (1895) renamed it S. vagrans monticola. This taxon is restricted to the southwestern United States; the type specimen was collected in Arizona. Sorex monticolus obscurus was described by Merriam (1891) as Sorex vagrans similis; however, the name similis was preoccupied by Sorex similis Hensel, 1855 (= Neomys similis). In 1895, Merriam renamed it Sorex obscurus. The type specimen for this taxon was collected in Idaho. Sorex monticolus setosus was described originally as a separate species (Sorex setosus) from Washington, by Elliot (1899); Jackson (1918) renamed it S. obscurus setosus without comment. Currently recognized subspecies S. m. calvertensis, S. m. insularis, S. m. malitiosus, S. m. neomexicanus, S. m. soperi, S. m. isolatus, S. m. mixtus, S. m. parvidens, S. m. alascensis, and S. m. longicaudus all were described originally as subspecies of S. obscurus (Anderson and Rand, 1945; Bailey, 1913; Cowan, 1941; Hall, 1938; Jackson, 1919, 1921, 1922; Merriam, 1895). Both S. m. elassodon and S. m. prevostensis, however, were described originally as subspecies of Sorex longicauda by Osgood (1901). Merriam (1895, 1900) referred to S. obscurus longicauda in text as S. longicauda, presumably shorthand for S. o. longicauda. Osgood (1901) seemingly interpreted Merriam's (1895, 1900) references to mean S. o. longicauda had been raised to the specific level and proceeded to describe S. m. elassodon and S. m. prevostensis as subspecies of S.

longicauda. In 1905, Elliot renamed S. m. elassodon and S. m. prevostensis as subspecies of S. obscurus. Similarly, S. m. alascensis was described originally by Merriam (1895) as a subspecies of S. obscurus, but S. m. shumaginensis was described by Merriam (1900) as a subspecies of Sorex alascensis. S. o. alascensis was referred to in text by Merriam (1895) as S. alascensis, thereby raising S. o. alascensis to the specific level; in 1900, he named S. alascensis shumaginensis. Allen (1902) renamed the latter taxon S. obscurus shumaginensis. By 1945, after the description of S. o. soperi by Anderson and Rand (1945), what is recognized currently as Sorex monticolus consisted of S. obscurus (with 15 subspecies), and a subspecies of S. vagrans (S. vagrans monticola).

A taxonomic revision of the shrews in the "Sorex vagrans species complex" changed all of the shrews in this complex, which includes S. monticolus, to subspecies of S. vagrans (Findley, 1955). Findley (1955) believed that S. vagrans and S. monticolus were not reproductively isolated throughout the range, but where sympatric, did not interbreed because they were the overlapping ends of a "Rassenkreis." With this revision, S. monticolus monticolus remained S. vagrans monticola, S. obscurus became S. vagrans obscurus, and all the taxa described as subspecies of S. obscurus kept their trinomial but became subspecies of S. vagrans. Johnson and Ostenson (1959) disagreed with Findley's (1955) revision and stated that until a more detailed study was accomplished, the taxonomy should remain as described by Jackson (1928). Jackson

(1928) separated S. vagrans and S. obscurus (=monticolus) on the basis of size (total length >110 mm in obscurus and <110 mm in vagrans). Hennings and Hoffmann (1977) separated S. vagrans and S. monticolus on the basis of the structure of the medial tines on the first upper incisors (I1), the height of red pigmentation on the anterior face of I1 in relation to the medial tines, and by limited morphometrics (they included only three variables--condylobasal length, least interorbital breadth, and palatal length). These authors considered S. vagrans monticola to be S. obscurus monticolus. Thus, Hennings and Hoffmann (1977) resurrected the name with priority, S. monticolus. Without explanation of differences among them, Hennings and Hoffmann (1977) further divided S. monticolus into 18 subspecies, seemingly based on geography and acceptance of previous trinomials. A multivariate morphometric analysis of the "Sorex vagrans species complex" in the Pacific Coast region indicated to Carraway (1990) that S. m. bairdi and S. m. permiliensis were specifically different from S. m. setosus and S. m. obscurus; she resurrected S. bairdii Merriam. George and Smith (1991) considered S. m. mixtus to be synonymous with S. m. setosus; they recognized S. m. setosus and S. m. isolatus as distinct taxa. Distribution of localities for specimens examined in these three studies combined included 791 distinct localities (Fig. 1a).

Sorex monticolus commonly occurs in high altitude spruce (Picea)-fir (Abies) forests and alpine tundra, and mid-altitude Douglas-fir (Pseudotsuga menziesii), lodgepole pine (Pinus contorta), western larch (Larix sp.), and grand fir (Abies grandis)





Fig. 1.--Distribution of (a) 791 unique localities for specimens used to establish current taxonomy of Sorex monticolus, based on Carraway (1990), George and Smith (1991), Hall (1981), and Hennings and Hoffmann (1977) and (b) 1336 unique localities for Sorex monticolus specimens included in this morphometric analysis.

forests (Hennings and Hoffmann, 1977). These habitat types occur in patchy areas possibly explaining the apparently patchy distribution of S. monticolus. Hennings and Hoffmann (1977) indicated that S. monticolus moved through continuous boreal-forest corridors during the Wisconsin glacial period and gained access to isolated mountainous regions such as the Sierra Nevada, San Bernardino, Piute, and Tehachapi mountains, and mountains in northern Mexico. Today, however, many of these regions (therefore, populations of shrews) are completely isolated in terms of appropriate habitat for S. monticolus.

Findley's (1955) classification of the "Sorex vagrans species complex" was based on overall size and color. Previous studies, including that of Hennings and Hoffmann (1977), were based on qualitative characters or on univariate to trivariate quantitative comparisons. From color and cranial characters, Jackson (1928) indicated that Sorex monticolus contains many intergrades, however, this also was based on qualitative observations. The geographic variation of Sorex monticolus has not been assessed previously by quantitative multivariate analyses.

The objectives of my research were to evaluate and describe the geographic variation within Sorex monticolus in the western United States, Canada, and Mexico, and to consider the appropriateness of current taxonomy applied to geographic variants. My research represents the first comprehensive quantitative analysis of geographic variation in Sorex monticolus. Many specimens have become available since Hennings and Hoffmann (1977) conducted their

research. A better understanding of the systematic relationships of these shrews is essential for further studies of ecological relationships in small-mammal communities and studies of relictual mammal populations.

METHODS

Skulls (and skins when available) of 3610 Sorex monticolus (sensu Hennings and Hoffmann, 1977, which includes Sorex obscurus, Sorex vagrans monticola, Sorex vagrans obscurus, and Sorex bairdi) from 1336 unique localities were examined and measured (Fig. 1b). Specimens were identified to species by tooth characteristics as described by Carraway (1990) and Hennings and Hoffmann (1977). Only undamaged skulls of adults, having slight but not excessive tooth wear (Jackson, 1928), were measured.

An ocular micrometer mounted in a Bausch and Lomb binocular microscope was used to measure 17 cranial and mandibular characters (Fig. 2). A Fowler Max-Cal electronic caliper was used to measure cranial depth and greatest length of skull to the nearest 0.01 mm. These 19 characters were selected because they are conventional dimensions found useful in previous studies of soricids (Carraway, 1990; Choate, 1972; Diersing, 1980; van Zyll de Jong, 1980) and because they have objective endpoints and are consistently repeatable. Values for characters measured with the ocular micrometer were retained as number of ocular lines for analysis; values subsequently were converted to millimeters and are reported as such herein. Gender in Sorex does not seem to effect cranial morphology (Rudd, 1955; van Zyll de Jong, 1980); in addition, many ($\geq 10\%$) museum specimens of shrews are misclassified to sex (L. Carraway, pers. comm.) thus biasing results claiming a difference. For these reasons sexes were combined in this analysis.




Fig. 2.--Camera lucida drawing of a *Sorex monticolus* (MSB 1944) illustrating skull and mandible dimensions measured (after van Zyll de Jong, 1980:67, fig. 1 and Carraway, 1990:11, fig. 4): 1, greatest length of skull (GLSKL); 2, maxillary breadth (MXBR); 3, least interorbital breadth (LIOB); 4, width across I1-I1 (I1I1); 5, breadth of zygomatic plate (ZYGP); 6, cranial breadth (CRBR); 7, cranial depth (CRDP); 8, breadth at M2-M2 (M2M2); 9, breadth at U4-U4 (U4U4); 10, breadth at U1-U1 (U1U1); 11, length of unicuspid toothrow (LUTR); 12, length of P4-M3 (P4M3); 13, width of upper condylar facet (WUCF); 14, width lower condylar facet (WLCF); 15, coronoid process-condylar length (CCLG); 16, length of mandibular toothrow (LMTR); 17, height of coronoid process (CORH); 18, greatest condylar depth (GCDP); 19, length of mandible (LGMN).

Sorex bairdi has been recognized previously as a subspecies of S. monticolus (S. monticolus bairdi and S. monticolus permiliensis--Hennings and Hoffmann, 1977). Subsequently, Carraway (1990) indicated that these taxa are specifically different from S. monticolus. S. bairdi and S. monticolus were analyzed for species-level differences by use of multigroup discriminant-functions analysis (BIOSTAT II--Pimentel and Smith, 1986).

A priori groups within Sorex monticolus were established primarily by use of currently accepted taxonomy. A priori groups consisted of alascensis, calvertensis, elassodon, insularis, isolatus, longicaudus, malitiosus, monticolus, neomexicanus, obscurus, prevostensis, setosus, and shumaginensis. I was unable to obtain a sufficiently large sample to include S. m. parvidens ($n = 6$) or S. m. soperi ($n = 1$) as a priori groups, thus, these groups were excluded from the initial analyses. I included S. m. obscuroides ($n = 116$) as a separate a priori group because this taxon was recognized previously as a subspecies of S. vagrans (= S. monticolus) by Findley (1955) and is geographically isolated. A reexamination of its relationship to the remaining taxa within the group was considered appropriate.

Straight-line polygons were drawn around the published marginal records (Hall, 1981; Hennings and Hoffmann, 1977) for each subspecies to form the 14 a priori groups (Fig. 3). Specimens from localities within a polygon were considered to belong to the taxon for which marginal records were used to produce that polygon. Specimens from localities outside of polygons ($n = 107$) were not




Fig. 3.--Map of marginal-record polygons for 14 subspecies of *Sorex monticolus* used as a priori groups in the present morphometric analysis. Marginal records based on Hall (1981) and Hennings and Hoffmann (1977).

included in the initial analyses, but were assigned to their appropriate groups by use of diagnosis files created with two- or three-group discriminant analyses between geographically adjacent a priori groups. S. bairdi ($n = 110$) was used as an outgroup.

Univariate summary statistics for all measurements were calculated for each group by use of basic descriptive statistics available in STATGRAPHICS (Statistical Graphics Corporation, 1987), and the corresponding coefficients of variation for each variable were compared. Multivariate analyses were performed on cranial and mandibular data for each group by use of the programs for principal-components analysis, multigroup discriminant-functions, and principal-coordinate analyses in BIOSTAT II (Pimentel and Smith, 1986); and multigroup discriminant-functions, covariance, and eigen-analyses were performed in STATGRAPHICS (Statistical Graphics Corporation, 1987). Principal-coordinate analysis was used to validate the results of the discriminant-functions analysis. Generalized-distance matrices derived from character means for each group, generated by the discriminant-functions analyses, were used in cluster analyses (unweighted pair-group method by use of arithmetic average) of the 14 a priori groups and in the three-group species-level analysis by use of BIOSTAT II (Pimentel and Smith, 1986). Plotting options were performed by use of STATGRAPHICS (Statistical Graphics Corporation, 1987), Corel Draw (Corel Corporation, 1992), and QuickBASIC graphics options (Microsoft Corporation, 1988). A probability level of $P \leq 0.05$ was accepted as statistically significant for all analyses.

Covariance and eigen-analyses were performed on the canonical-variate scores for each group to generate eigenvectors and eigenvalues (Statistical Graphics Corporation, 1987) used to calculate 95% confidence ellipses around group centroids. The first and second eigenvalues and n were used to calculate the half-lengths for the major and minor axes. The focal length for each ellipse is the square root of the difference between the two squared half-lengths. The arc cosine of the first element of the first eigenvector equals the slope of the major axis (Owen and Chmielewski, 1985).

Systematic accounts contain the original taxonomic designation, new name combinations, and junior synonyms where appropriate. In specimens-examined lists, specimens are ordered alphabetically by state, county, major geographic point in specific locality, and museum acronym, and numerically by catalog number. Specimens from localities not marked on distribution maps are marked with an asterisk in the appropriate specimens-examined list. On distribution maps, many locality symbols cover more than one unique locality. Instead of presenting subspecies as wholly separate units, I have attempted to describe the regions of intergradation between adjacent subspecies. Regions where two taxa come together were considered zones of intergradation between subspecies because the shrews from these regions typically had weak classifications. Such individuals are included in the specimens-examined list of the taxon with which they had the strongest affinities.

RESULTS

A comparison of all 19 cranial and mandibular characters for each a priori group indicated that for width of the upper condylar facet the coefficient of variation was nearly twice that of all other characters. Because this character exhibited so much variation within each group, it seemed of limited value for distinguishing between groups and was dropped from all further analyses.

SPECIES-LEVEL ANALYSIS

Univariate Student-Newman-Keuls multiple-range tests for Sorex bairdi ($n = 110$) and S. monticolus ($n = 3500$) resulted in a significant difference between means of all characters analyzed. Sorex bairdi averaged larger than S. monticolus in all characters measured; however, there was significant overlap of ranges. In a discriminant analysis between these two taxa, the group centroids were significantly different (F -ratio = 145.191, Wilks Lambda = 0.5787787; $d.f. = 18, 3591$) and 99% of the individuals were correctly classified into their a priori groups (Fig. 4a). Twenty-two S. bairdi and eight S. monticolus were misclassified. Characters with the highest positive coefficients of correlation to the first canonical-variate were height of coronoid process (0.792), breadth at U4-U4 (0.701), length of mandible (0.644), and coronoid process-condylar length (0.610). The Mahalanobis' distance between these two groups was 4.962.




Fig. 4.--Means, 95% confidence intervals, and ranges of discriminant scores for two-group discriminant analyses with (a) *S. bairdi* and *S. monticolus*, (b) *S. m. neomexicanus* and the remaining subspecies of *S. monticolus*, and (c) *S. bairdi* and *S. m. setosus* with three characters.

VARIATION WITHIN SOREX MONTICOLUS

Principal-components analysis was performed on each of the a priori groups within Sorex monticolus to determine if variation within any one group was greater than expected for one taxon. The first two components for each group were examined for possible outliers and for an indication that any of the a priori groups should be separated into more than one group. All a priori groups remained as grouped.

A 15-group discriminant analysis with the 14 a priori S. monticolus groups and S. bairdi resulted in a 67% overall correct classification of individuals. The first two canonical variates accounted for 73.87% (46.59% and 27.28%, respectively) of the among-group variation within the 18 remaining cranial and mandibular characters examined (Fig. 5). Characters with the highest positive coefficients of correlation to the first canonical-variate axis were greatest length of skull (0.847) and length of unicuspid toothrow (0.518). Characters with the highest positive coefficients of correlation to the second canonical-variate axis were height of the coronoid process (0.845), breadth at U1-U1 (0.688), breadth at U4-U4 (0.639), and coronoid process-condylar length (0.616). A principal-coordinate analysis of the generalized-distance matrix derived from character means resulted in a distribution of group means on the first two coordinates consistent with the results of the discriminant analysis. A cluster analysis of the generalized-distance matrix resulted in a distance phenogram (Fig. 6) that indicated that S. m. neomexicanus is morphologically distinct from




Fig. 5.--Confidence ellipses (95%) for all 14 a priori *S. monticolus* groups and *S. bairdi*.




Fig. 6.--A dendrogram generated by a cluster analysis based on the generalized-distance matrix derived from the 15-group discriminant analysis for S. bairdi and S. monticolus.

the other 13 a priori groups at Mahalanobis' distance 4.836. Both the 95% confidence ellipses (Fig. 5) and the distance phenogram (Fig. 6) indicated that S. m. neomexicanus is at least as distinct morphometrically as S. bairdi from S. monticolus. In addition, in the S. bairdi and S. monticolus discriminant analysis, all eight of the misclassified S. monticolus individuals were from the subspecies S. m. neomexicanus. Another four (8.7%) of the 46 S. m. neomexicanus individuals had Geisser-probability classifications <80%; 10 (22%) had classification probabilities <60%. Of all remaining S. monticolus ($n = 3454$) only 10 individuals had a classification probability <90%. The poor classification of individuals within the S. m. neomexicanus a priori polygon indicated that further analysis of S. m. neomexicanus was appropriate. Another species-level discriminant-functions analysis was performed for S. m. neomexicanus ($n = 46$) and the other subspecies of S. monticolus ($n = 3454$).

Univariate Student-Newman-Keuls multiple-range tests for Sorex m. neomexicanus and the other subspecies of S. monticolus resulted in a significant difference between means for all characters analyzed. Sorex m. neomexicanus averaged larger than other subspecies of S. monticolus in all characters measured; however, ranges overlapped significantly (Table 1). In a discriminant analysis between S. m. neomexicanus and the remaining subspecies of S. monticolus, group centroids were significantly different (F -ratio = 66.537, Wilks Lambda = 0.7440163; $d.f. = 18, 3481$). The Geisser-classification probabilities resulted in the correct classification

Table 1. Mean $\pm$ SE for all taxa included in analyses.
Sample size (n) for

Character	<i>alascensis</i> ($n = 252$)	<i>calverti</i> ($n = 2$)	<i>cecostensis</i> ($n = 25$)	<i>setosus</i> ($n = 506$)	<i>shumaginensis</i> ($n = 194$)	<i>soperi</i> ($n = 1$)	<i>neomexicanus</i> ($n = 46$)	<i>Sorex</i> <i>bairdi</i> ($n = 110$)
Maxillary breadth	5.07 $\pm$ 0.01 4.57-5.71 4.28	4.85 $\pm$ 0.01 4.57-5.13 2.88	5.21 $\pm$ 0.05 4.71-5.86 4.48	4.90 $\pm$ 0.01 4.43-5.43 3.96	4.82 $\pm$ 0.01 4.43-5.29 3.02	4.71	5.24 $\pm$ 0.02 5.00-5.57 3.29	5.49 $\pm$ 0.03 4.71-6.00 5.47
Least interorbital breadth	3.68 $\pm$ 0.01 3.43-4.00 3.95	3.63 $\pm$ 0.01 3.29-3.97 3.37	3.85 $\pm$ 0.02 3.57-4.00 2.38	3.58 $\pm$ 0.01 3.14-4.14 4.04	3.52 $\pm$ 0.01 3.29-3.86 3.28	3.43	3.80 $\pm$ 0.02 3.57-4.00 3.20	3.91 $\pm$ 0.02 3.43-4.43 4.90
Cranial breadth	8.81 $\pm$ 0.02 8.14-9.86 3.97	8.31 $\pm$ 0.01 8.00-8.62 1.93	8.78 $\pm$ 0.05 8.00-9.14 2.82	8.45 $\pm$ 0.01 7.43-9.43 3.73	8.56 $\pm$ 0.02 8.14-9.14 2.50	8.00	8.81 $\pm$ 0.04 8.14-9.29 3.50	9.26 $\pm$ 0.04 8.29-9.86 4.59
Breadth of zygomatic plate	2.08 $\pm$ 0.01 1.86-2.57 6.23	1.93 $\pm$ 0.01 1.86-2.17 4.80	1.97 $\pm$ 0.02 1.71-2.14 5.53	1.90 $\pm$ 0.01 1.57-2.14 5.67	1.98 $\pm$ 0.01 1.71-2.14 4.89	2.00	2.15 $\pm$ 0.01 2.00-2.43 4.19	2.11 $\pm$ 0.01 1.86-2.43 6.01
Breadth at M2-M2	4.63 $\pm$ 0.01 4.29-5.14 4.03	4.44 $\pm$ 0.01 4.29-5.14 7.05	4.62 $\pm$ 0.02 4.43-4.86 2.66	4.46 $\pm$ 0.01 3.86-5.14 3.96	4.35 $\pm$ 0.01 4.00-4.86 2.90	4.14	4.89 $\pm$ 0.03 4.57-5.14 3.59	4.97 $\pm$ 0.02 4.43-5.43 4.85
Breadth at U1-U1	1.85 $\pm$ 0.01 1.57-2.14 5.60	1.70 $\pm$ 0.01 1.57-1.86 2.28	1.79 $\pm$ 0.01 1.71-1.86 4.03	1.76 $\pm$ 0.01 1.57-2.00 5.89	1.74 $\pm$ 0.01 1.57-1.86 3.98	1.71	2.15 $\pm$ 0.01 2.00-2.29 3.81	2.07 $\pm$ 0.01 1.86-2.29 6.46
Breadth at U4-U4	2.32 $\pm$ 0.01 2.00-2.57 4.71	2.21 $\pm$ 0.01 2.00-2.43 4.18	2.35 $\pm$ 0.01 2.29-2.43 3.09	2.28 $\pm$ 0.01 2.00-2.57 5.31	2.18 $\pm$ 0.01 2.00-2.43 3.69	2.14	2.60 $\pm$ 0.02 2.43-2.71 4.23	2.70 $\pm$ 0.01 2.43-3.00 5.79
Width across I1-I1	1.57 $\pm$ 0.01 1.29-1.86 7.18	1.48 $\pm$ 0.01 1.29-1.71 6.14	1.55 $\pm$ 0.02 1.43-1.71 5.51	1.50 $\pm$ 0.01 1.29-1.71 7.36	1.49 $\pm$ 0.01 1.29-1.71 5.23	1.57	1.74 $\pm$ 0.01 1.57-1.86 5.15	1.79 $\pm$ 0.01 1.57-2.00 6.56
Length of P4-M3	4.20 $\pm$ 0.01 3.86-4.71 4.54	4.14 $\pm$ 0.01 4.00-4.29 2.18	4.29 $\pm$ 0.02 4.14-4.57 2.45	4.03 $\pm$ 0.01 3.57-4.57 3.34	3.97 $\pm$ 0.01 3.71-4.29 3.10	4.00	4.30 $\pm$ 0.02 4.00-4.57 3.46	4.39 $\pm$ 0.03 2.86-4.71 6.17
Length of unicuspid tooththrow	2.63 $\pm$ 0.01 2.29-3.14 5.30	2.44 $\pm$ 0.01 2.29-2.71 2.83	2.51 $\pm$ 0.02 2.29-2.71 4.34	2.45 $\pm$ 0.01 2.14-2.86 5.29	2.49 $\pm$ 0.01 2.29-2.71 4.04	2.43	2.85 $\pm$ 0.03 2.57-3.29 6.55	2.80 $\pm$ 0.01 2.43-3.29 5.56
Cranial depth	5.16 $\pm$ 0.02 4.38-6.03 5.36	4.99 $\pm$ 0.01 4.41-5.57 4.84	5.23 $\pm$ 0.06 4.80-5.79 5.33	4.99 $\pm$ 0.01 4.19-5.71 5.88	5.13 $\pm$ 0.02 4.22-5.67 4.93	4.94	5.12 $\pm$ 0.04 4.45-5.67 5.33	5.27 $\pm$ 0.02 4.72-5.86 4.30
Greatest length of skull	17.81 $\pm$ 0.02 16.43-18.89 2.08	17.86 $\pm$ 0.01 17.25-18.89 1.76	18.34 $\pm$ 0.07 17.77-19.15 1.85	17.43 $\pm$ 0.02 15.76-18.65 2.41	17.20 $\pm$ 0.02 16.31-18.18 2.07	17.22	17.97 $\pm$ 0.03 17.42-18.65 1.31	18.66 $\pm$ 0.04 17.49-19.58 2.52
Length of mandible	8.30 $\pm$ 0.02 7.43-9.29 4.18	7.81 $\pm$ 0.01 7.43-8.62 2.69	8.41 $\pm$ 0.06 8.00-9.14 3.39	7.95 $\pm$ 0.01 6.71-9.14 4.21	7.79 $\pm$ 0.02 7.29-8.86 2.90	7.71	8.52 $\pm$ 0.05 7.71-9.14 3.92	9.03 $\pm$ 0.05 7.86-9.86 5.98
Length of mandibular tooththrow	5.19 $\pm$ 0.01 4.86-5.71 3.90	4.97 $\pm$ 0.01 4.86-5.57 2.04	5.20 $\pm$ 0.02 5.00-5.43 2.38	4.91 $\pm$ 0.01 4.29-5.43 3.41	4.85 $\pm$ 0.01 4.57-5.29 2.63	4.86	5.46 $\pm$ 0.03 5.14-5.86 3.75	5.37 $\pm$ 0.03 4.71-5.86 5.06
Height of coronoid process	4.05 $\pm$ 0.01 3.71-4.57 5.01	3.72 $\pm$ 0.01 3.57-3.86 2.85	3.91 $\pm$ 0.03 3.57-4.14 3.49	3.84 $\pm$ 0.01 3.29-4.29 5.37	3.77 $\pm$ 0.01 3.57-4.14 3.17	3.71	4.47 $\pm$ 0.03 4.14-4.86 4.18	4.70 $\pm$ 0.03 4.00-5.29 7.06
Greatest condylar depth	2.12 $\pm$ 0.01 1.86-2.43 5.35	2.03 $\pm$ 0.01 1.86-2.17 4.46	2.18 $\pm$ 0.02 2.00-2.57 5.83	2.05 $\pm$ 0.01 1.71-2.43 5.57	2.04 $\pm$ 0.01 1.86-2.29 4.15	1.86	2.21 $\pm$ 0.01 2.00-2.43 4.46	2.33 $\pm$ 0.01 2.00-2.71 6.67
Width of lower condylar facet	1.24 $\pm$ 0.01 1.00-1.43 7.09	1.20 $\pm$ 0.01 1.14-1.26 5.98	1.30 $\pm$ 0.01 1.14-1.43 5.43	1.20 $\pm$ 0.01 1.00-1.43 7.04	1.17 $\pm$ 0.01 1.00-1.29 5.25	1.14	1.32 $\pm$ 0.01 1.14-1.43 5.31	1.42 $\pm$ 0.01 1.14-1.71 8.15
Coronoid process-condylar length	2.88 $\pm$ 0.01 2.43-3.29 5.27	2.68 $\pm$ 0.01 2.43-2.97 4.15	2.85 $\pm$ 0.02 2.57-3.00 3.68	2.78 $\pm$ 0.01 2.43-3.14 5.54	2.70 $\pm$ 0.01 2.43-3.00 4.41	2.71	3.10 $\pm$ 0.02 2.86-3.43 4.65	3.26 $\pm$ 0.02 2.71-3.71 7.61

into their a priori groups of all but two S. m. neomexicanus, and one of the remaining S. monticolus (Fig. 4b). Characters with the highest positive coefficients of correlation to the first canonical-variate axis were breadth at U1-U1 (0.732), height of coronoid process (0.594), length of unicuspid tooththrow (0.565), and breadth at U4-U4 (0.556). The Mahalanobis' distance between these two groups was 5.149. S. m. neomexicanus was not included in subsequent analyses of geographic variation within S. monticolus.

A multigroup discriminant analysis with the 13 remaining a priori S. monticolus groups resulted in a 67% overall correct classification of individuals. The first two canonical variates accounted for 76.17% (60.94% and 15.23%, respectively) of the among-group variation within the 18 remaining cranial and mandibular characters examined (Fig. 7a). The character with the highest positive coefficient of correlation to the first canonical-variate axis was greatest length of skull (0.760); all remaining characters had correlation coefficients of <0.400. Characters with the highest positive coefficients of correlation to the second canonical-variate axis were length of unicuspid tooththrow (0.720), cranial breadth (0.565), breadth of zygomatic plate (0.525), and length of mandibular tooththrow (0.504). The first canonical variate apparently represents overall size variation, whereas the second canonical variate represents variation in shape. A cluster analysis of the generalized-distance matrix derived from character means resulted in a distance phenogram (Fig. 8a) that demonstrates a separation of the 13 a priori groups into two large clusters at Mahalanobis' distance




Fig. 7.--Confidence ellipses (95%) for (a) the remaining 13 a priori *S. monticolus* groups, (b) *S. m. obscurus* and *S. m. obscuroides*, (c) *S. m. calvertensis* and *S. m. elassodon*, and (d) the 12 remaining groups after all specimens had been placed in appropriate groups.





Fig. 8.--Dendograms generated by cluster analyses based on the generalized-distance matrices derived from the multigroup discriminant analyses for (a) the 13 a priori groups, and (b) the 12 remaining groups with all specimens in groups.

3.855. S. m. obscurus and S. m. obscuroides clustered together at Mahalanobis' distance 1.472 and seem to be similar morphologically (Fig. 7b). S. m. obscuroides was grouped with S. m. obscurus in further analyses. S. m. calvertensis and S. m. elassodon cluster together at Mahalanobis' distance 1.226; these groups also seem to be similar morphologically (Fig. 7c). Both summer and winter pelages of S. m. calvertensis (from both Calvert and Banks islands, British Columbia), however, are much paler than those of either S. m. elassodon or S. m. longicaudus (Cowan, 1941). Pelage color was observed but not measured quantitatively in my study. Because of the small sample size for S. m. calvertensis ($n = 26$) it seemed prudent to be conservative, so S. m. calvertensis was retained as a separate a priori group for the remaining analyses. A biochemical analysis of the shrews currently designated S. m. calvertensis, S. m. elassodon, and S. m. longicaudus may be needed to decipher these relationships.

Many misclassified individuals in the 13-group discriminant analysis were classified with groups from geographically distant areas, consequently these affiliations probably have no biological meaning (e.g., specimens from British Columbia being more similar to groups in Arizona than to groups adjacent geographically). Because of the number of groups involved and the similarity of some geographically distant groups, the overall classification of individuals was expectedly poor. Thus, two- or three-group discriminant analyses with geographically adjacent a priori groups were performed to obtain diagnosis files to increase the probability

of assigning specimens not in a priori groups to their proper groups (Table 2).

Of those specimens outside polygons for S. m. shumaginensis, S. m. alascensis, and S. m. obscurus one individual was approximately equidistant between them. A three-group discriminant analysis was performed to obtain a diagnosis file. The discriminant analysis resulted in 89% correct classification of individuals into their a priori groups and indicated that the unknown individual should be classified as S. m. obscurus.

Some individuals misclassified in the two- and three-group discriminant analyses were examined more closely to determine if their group designation should be changed. In the S. m. monticolus and S. m. obscurus discriminant analysis, 16 of 18 individuals in the S. m. monticolus a priori group from New Mexico and all six of the individuals from Mexico were misclassified as S. m. obscurus. The Mexico specimens had classification probabilities $\geq 83\%$ and 10 of 16 misclassified individuals from New Mexico had classification probabilities of $\geq 80\%$. Within the Arizona specimens from the S. m. monticolus a priori group, 12 of 13 individuals from the northeastern portion of the state (northern Apache Co. and Coconino Co.) were misclassified as S. m. obscurus; eight of these 12 had classification probabilities of $\geq 72\%$. These misclassifications indicated that individuals in northeastern Arizona, western New Mexico, and Mexico should be considered S. m. obscurus. These individuals were regrouped for further analyses.

Table 2. se specimens from localities between a priori polygons.

rs Subspecies 1 significantly nt from Subspecies 2 ^{ab}	
Subspecies ^a aller	Larger
<u>S. m. shum</u> IOB, ZYGP, M2M2, 4U4, MNTR, CORH, CLG	
<u>S. m. shum</u> pt CRDP	
<u>S. m. obs</u> racters	
<u>S. m. long</u> YGP, M2M2, U1U1, GMN, CORH, CCLG	LIQB, U4U4, P4M3, GLSKL, WLCF
<u>S. m. obs</u> racters	
<u>S. m. set</u> cept CCLG	
<u>S. m. iso</u> racters	
<u>S. m. obs</u> rBR, U4U4, LUTR, LSKL, LGMN, LMTR,	ZYGP, M2M2, U1U1, P4M3, CORH, WLCF, CCLG
<u>S. m. mont</u> pt IIII	
^a Based on	
^b Acronyms	

In the S. m. alascensis and S. m. obscurus discriminant analysis, all 11 individuals from Tats Lake, British Columbia, were misclassified as S. m. obscurus; 10 of 11 had classification probabilities $\geq 74\%$ and one individual had a classification probability of 63%. Tats Lake is on the northern edge of the previously recognized distribution of S. m. alascensis and is adjacent to the range of S. m. obscurus. High Geisser classifications and absence of individuals classified as S. m. alascensis from this locality indicate that instead of being a zone of intergradation, Tats Lake should be included within the range of S. m. obscurus. One individual each from Chilkoot Lake and Wells, and two from Skagway also were misclassified as S. m. obscurus (classification probabilities of 91%, 99%, 68%, and 95%, respectively). The area including all three of these localities also is near the zone of intergradation for S. m. alascensis and S. m. obscurus; dusky shrews from these localities should be considered S. m. obscurus. These 15 individuals were considered S. m. obscurus for further analyses.

In the S. m. longicaudus and S. m. obscurus discriminant analysis, the individual from Eutsuk Lake and one of two from Meziadin Lake, British Columbia, were misclassified as S. m. obscurus. These lakes are on the eastern edge of the previously recognized distribution of S. m. longicaudus. The same two shrews from Meziadin Lake were misclassified in the discriminate analysis between S. m. setosus and S. m. longicaudus and had poor classification probabilities ($<60\%$) in the S. m. alascensis and S.

m. longicaudus discriminant analysis. This indicated that these shrews do not belong with the S. m. longicaudus group and should be considered S. m. obscurus. All three individuals were added to the S. m. obscurus group for further analyses. In the same discriminant analysis both individuals from Lakelse Lake, British Columbia, were misclassified as S. m. longicaudus (having classification probabilities of 96% and 100%). This lake is near the point of parapatry for S. m. obscurus and S. m. longicaudus; Lakelse Lake should be included in the range of S. m. longicaudus resulting in an eastern shift in the distribution of S. m. longicaudus. Both of these individuals were grouped with S. m. longicaudus for further analyses.

In the S. m. longicaudus and S. m. setosus discriminant analysis, 12 of 21 individuals from Goose Island, British Columbia, were misclassified as S. m. setosus. Ten of these 12 misclassifications had probability classifications of $\geq 71\%$, whereas five of the nine "correct" classifications were $< 70\%$. The mainland range of S. m. setosus extends from approximately Rivers Inlet, northeast to Hagensborg and Stuie, British Columbia. Goose Island is northwest of the previously accepted distribution of S. m. setosus; however, my results indicate that the shrews from Goose Island should be considered S. m. setosus. Similarly, the individual from Hecate Island, slightly northwest of Rivers Inlet, was misclassified as S. m. setosus. The classification probability was weak (63%) but this specimen was misclassified in three of five other discriminant analyses involving S. m. longicaudus. The degree

of isolation could explain the poor classification probabilities of shrews from both Hecate Island and Goose Island. All 22 shrews were grouped with S. m. setosus for further analyses.

A series of three-group discriminant analyses was used in an attempt to clarify the relationships of the shrews from the coastal islands of British Columbia and Alaska. The Estevan group of islands, particularly Dewdney Island, British Columbia, is within the a priori group of S. m. longicaudus; however, it is isolated (ca. 10 km) from all other islands. In a discriminant analysis including S. m. alascensis, S. m. longicaudus, and S. m. elassodon, four of six shrews from Dewdney Island had classification probabilities of $\geq 70\%$ as S. m. elassodon; two had weak classification probabilities (50% and 51% as S. m. longicaudus). Similarly, in a discriminant analysis including S. m. longicaudus, S. m. elassodon, and S. m. malitiosus, the same shrews again were classified primarily as S. m. elassodon; four of six had classification probabilities of $\geq 80\%$ as S. m. elassodon, and two had weak classification probabilities (56% as S. m. elassodon and 51% as S. m. longicaudus). In spite of the geographic distance between Dewdney Island and the nearest S. m. elassodon population, these shrews are more closely allied with S. m. elassodon than with S. m. longicaudus. Similarly, shrews from Porcher Island, British Columbia, ($n = 4$) previously were considered to be S. m. longicaudus; however, in the discriminant analysis including S. m. alascensis, S. m. longicaudus, and S. m. elassodon three of four individuals had classification probabilities of $\geq 80\%$ as S. m.

elassodon. The classification probability of the fourth individual was weak (47%) but again was classified as S. m. elassodon. In a discriminant analysis including S. m. longicaudus, S. m. elassodon, and S. m. malitiosus all four shrews from Porcher Island had classification probabilities of $\geq 81\%$ as S. m. elassodon. In a discriminant analysis including S. m. longicaudus, S. m. insularis, and S. m. calvertensis, neither the shrews from Porcher Island nor from Dewdney Island were at all similar to S. m. insularis or S. m. calvertensis. These 10 shrews were grouped with S. m. elassodon for further analyses.

Admiralty Island, Alaska, previously was considered to be within the range of S. m. elassodon but is near the zone of intergradation with S. m. alascensis. In the discriminant analysis including S. m. alascensis, S. m. longicaudus, and S. m. elassodon the individual from Admiralty Island had a strong classification probability (97%) as S. m. alascensis. In the discriminant analysis including S. m. longicaudus, S. m. elassodon, and S. m. malitiosus (but not including S. m. alascensis) this individual was classified as S. m. longicaudus. In the original 13-group discriminant analysis this individual had classification probabilities of 0% as S. m. elassodon and 10% as S. m. alascensis. These results taken together indicated that this individual does not have close affinities with S. m. elassodon and should be considered S. m. alascensis. This shrew was grouped with S. m. alascensis for further analyses.

Forrester Island is nearest to the distribution of S. m. elassodon; however, in the discriminant analysis including S. m.

alascensis, S. m. longicaudus, and S. m. elassodon and in the discriminant analysis including S. m. longicaudus, S. m. elassodon, and S. m. malitiosus, the one individual from Forrester Island had a weak classification probability (78% and 74%, respectively) as S. m. longicaudus. Because the classification probabilities were relatively poor and because only one individual was examined from Forrester Island, this shrew was retained in the S. m. elassodon group. Shrews from the coastal islands of southeast Alaska and British Columbia, including Forrester Island, need further analysis to clarify these relationships.

Campania Island is within the range of S. m. longicaudus. In the discriminant analysis including S. m. alascensis, S. m. longicaudus, and S. m. elassodon, the shrews from Campania Island ($n = 2$) had weak classification probabilities as S. m. elassodon (65% and 75%). In the discriminant analysis including S. m. longicaudus, S. m. elassodon, and S. m. malitiosus, these shrews again were classified as S. m. elassodon (classification probabilities of 69% and 88%). In the discriminant analysis including S. m. longicaudus, S. m. insularis, and S. m. calvertensis, these shrews were classified as S. m. longicaudus (both had classification probabilities of 65%). Because classifications were poor and because only two individuals were examined from Campania Island, these shrews were retained in the S. m. longicaudus group. Shrews in the area including Campania Island also need further analysis to decipher their relationships.

A discriminant analysis for the remaining 12 groups (reflecting all group affiliation changes and the placement of diagnosed individuals) resulted in a 72% correct classification of individuals into their a priori groups. This reflects a slight improvement in the classification probabilities from the original a priori group affiliations. The first two canonical-variate axes represent 76.35% of the among-group variation (62.44% and 13.91%, respectively; Fig. 7d). The character with the highest positive coefficient of correlation to the first canonical-variate axis was greatest length of skull (0.768); all of the remaining characters had correlation coefficients of <0.400 . Characters with the highest positive coefficients of correlation to the second canonical-variate axis were length of unicuspid toothrow (0.704), cranial breadth (0.604), breadth of zygomatic plate (0.584), length of mandibular toothrow (0.554), and height of coronoid process (0.513).

A cluster analysis of the generalized-distance matrix derived from character means resulted in a distance phenogram (Fig. 8b) that demonstrates the separation of the 12 groups into two large clusters at Mahalanobis' distance of 3.935. Changes made in the group affiliations did not alter the strong morphometric similarity of S. m. calvertensis and S. m. elassodon; these two taxa clustered together at a Mahalanobis' distance of 1.227. S. m. isolatus and S. m. setosus clustered together at Mahalanobis' distance of 1.529. However, the two-group discriminant analysis for S. m. isolatus and S. m. setosus indicated that these two taxa were significantly different. In the first discriminant analysis (including all 13 a

priori groups before evaluation of group affiliation and addition of diagnosed individuals) the Mahalanobis' distance between S. m. obscurus and S. m. monticolus was 1.789 (Fig. 8a). After changing the group affiliation of the shrews from Mexico, western New Mexico, and northeastern Arizona from S. m. monticolus to S. m. obscurus, these two taxa were linked together at Mahalanobis' distance of 2.082 (Fig. 8b).

The primary difference between the results of the discriminant analyses including 12 and 13 subspecies is in the placement of S. m. longicaudus. In the original analysis (including 13 a priori groups), S. m. longicaudus grouped closely with S. m. calvertensis and S. m. elassodon (Fig. 8a). After removing the shrews from Goose and Hecate islands and Eutsuk and Meziadin lakes from the S. m. longicaudus group, this taxon no longer clusters closely with S. m. elassodon or with S. m. calvertensis (Fig. 8b).

SYSTEMATICS AND TAXONOMY

Order Insectivora

Superfamily Soricoidae

Family Soricidae

Subfamily Soricinae

Tribe Soricini

Genus SorexSubgenus OtisorexSorex neomexicanus Bailey

New Mexico Shrew

1913. Sorex obscurus neomexicanus Bailey, Proc. Biol. Soc.
Washington, 26:133, May.
1955. Sorex vagrans neomexicanus Findley, Univ. Kansas Publ., Mus.
Nat. Hist., 9:50, December.
1977. S[orex]. m[onticolus]. neomexicanus Hennings and Hoffmann,
Occas. Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.

Holotype.--USNM 100440, original number 7383; adult male
collected 29 May 1900 from "Cloudcroft, New Mexico . . . in the
Sacramento Mountains" by V. Bailey. Type specimen examined and
measured.

Distribution.--Sorex neomexicanus occurs in southcentral New Mexico in the Capitan and Sacramento mountains. The distribution of this shrew might extend east at least as far as the Pecos River (Fig. 9).

Diagnosis.--Sorex neomexicanus can be distinguished from congeners in New Mexico by the following characters: from S. nanus, S. cinereus, and S. merriami by 3rd upper unicuspid smaller than 4th; from S. palustris by upper unicuspid wider than long, and possessing a brownish pelage instead of black and white; and from S. monticolus obscurus by a combination of the length of unicuspid tooththrow and breadth at U1-U1.

Description.--Sorex neomexicanus is larger overall than any other Sorex in New Mexico except S. palustris. S. neomexicanus is considerably larger (Table 1) and slightly darker than S. m. obscurus; the dorsum is "dull sepia brown" with less reddish coloration than S. m. obscurus, and the venter has a "brownish suffusion" instead of the gray venter of S. m. obscurus (Bailey, 1913:134). Mean measurements for skull and mandible characteristics ($\bar{n}$ = 46) are (in mm): maxillary breadth = 5.24, least interorbital breadth = 3.80, cranial breadth = 8.81, breadth at zygomatic plate = 2.15, breadth at M2-M2 = 4.89, breadth at U1-U1 = 2.15, breadth at U4-U4 = 2.60, width across I1-I1 = 1.74, length of P4-M3 = 4.30, length of unicuspid tooththrow = 2.85, cranial depth = 5.12, greatest length of skull = 17.97, length of mandible = 8.52, length of




Fig. 9.--Distribution of localities for *S. neomexicanus* specimens included in this analysis.

mandibular toothrow = 5.46, height of coronoid process = 4.47, greatest condylar depth = 2.21, width of lower condylar facet = 1.32, coronoid process-condylar length = 3.10. Mean measurements for skin characteristics are (in mm): total length = 111.58 ($\underline{n}$ = 24), tail length = 43.76 ($\underline{n}$ = 25), and hind foot length = 13.32 ($\underline{n}$ = 25; Fig. 10).

Comparisons and remarks.--Sorex m. obscurus is the only subspecies of S. monticolus in New Mexico, therefore the only one with which S. neomexicanus might be confused. In a discriminant analysis between these two taxa ($\underline{F}$ -ratio = 80.901, Lambda = 0.5392039; $\underline{d.f.}$ = 18, 1704), 99.88% of the individuals were classified into their a priori groups. Length of unicuspid toothrow, breadth at U1-U1, length of mandibular toothrow, height of coronoid process, and breadth at U4-U4 accounted for the greatest degree of segregation. It is possible, however, to classify 97.1% of individuals as either S. neomexicanus or S. m. obscurus based solely on a combination of length of unicuspid toothrow and breadth at U1-U1 ($\underline{F}$ -ratio = 491.236, Lambda = 0.6364543; $\underline{d.f.}$ = 2, 1720):

length of unicuspid toothrow (5.27408) + breadth at

U1-U1 (5.99613) - 23.3964 = discriminant score

where, if score < -3.1 = S. neomexicanus

if score > -2.1 = S. m. obscurus.

If the score for an individual is between these two values, it is not possible to assign it to either taxon.




Fig. 10.--Camera lucida tracing of skull and mandible of S. neomexicanus (MSB 41081).

When S. bairdi ($n = 110$), S. monticolus ($n = 3454$), and S. neomexicanus ($n = 46$) are considered jointly, 99% of individuals were classified into their correct a priori group (Fig. 11). Height of coronoid process, breadth at U4-U4, breadth at U1-U1, and length of mandible accounted for the greatest degree of segregation. A cluster analysis based on the generalized-distance matrix derived from the multigroup discriminant analysis resulted in the joining of S. bairdi and S. neomexicanus at Mahalanobis' distance 4.587 and the subsequent joining of S. monticolus at Mahalanobis' distance 5.163 (Fig. 12).

Specimens examined.--NEW MEXICO (46).--Lincoln Co.: Capitan Mts., 3 (MSB 4755--4757); southwest slope Capitan Mts., 1 (USNM 128188); 5 mi W Alto, 3 (MSB 16561, 16562, 18970); Otero Co.: Cloudcroft, 3 (USNM 100440--100442); 1 mi S Cloudcroft, 1 (FHSU 5435); Deer Head Campground, 1 mi S Cloudcroft, 1 (RHMC 7492); 2 mi N Cloudcroft, 1 (MSB 18043); 3.2 mi E (by road) Cloudcroft, 6 (MSB 37327--37330, 37340, 37341); 5.5 mi NE (by road) Cloudcroft, 22 (MSB 37343, 37345, 37347--37357, 37359, 37360, 41069, 41071, 41078, 41081, 41084, 41086, 41087); Silversprings Creek, 5.5 mi NE (by road) Cloudcroft, 2 (MSB 41056, 41057); 10 mi NE Cloudcroft, 3 (USNM 118789--118791).




Fig. 11.--Confidence ellipses (95%) for *S. bairdi*, *S. neomexicanus*, and the remaining *S. monticolus*.




Fig. 12.--A dendrogram generated by a cluster analysis based on the generalized-distance matrix derived from the multigroup discriminant analysis for S. bairdi, S. neomexicanus, and the remaining S. monticolus.

Sorex bairdi Merriam

Baird's shrew

1895. Sorex bairdi Merriam, N. Amer. Fauna, 10:77, December.
1918. Sorex obscurus bairdi Jackson, Proc. Biol. Soc. Washington, 31:127, November.
1955. Sorex vagrans bairdi Findley, Univ. Kansas Publ., Mus. Nat. Hist., 9:35, December.
1977. S[orex]. m[onticolus]. bairdii Hennings and Hoffmann, Occas. Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.
Incorrect use of terminal "i" on bairdi.
1990. Sorex bairdii bairdii Carraway, Spec. Publ., The Mus., Texas Tech. Univ., 32:39.
Incorrect use of terminal "i" on bairdi.

Holotype.--USNM 17414/24318, original number 270; adult female collected on 2 August 1889 from "Astoria, Oregon" by T. S. Palmer.
Type specimen not examined.

1918. Sorex obscurus permiliensis Jackson, Proc. Biol. Soc. Washington, 31:128, November.
1955. Sorex vagrans permiliensis Findley, Univ. Kansas Publ., Mus. Nat. Hist., 9:36, December.
1977. S[orex]. m[onticolus]. permiliensis Hennings and Hoffmann, Occas. Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.
1977. S[orex]. m[onticolus]. permilliensis Hennings and Hoffmann,

Occas. Papers Mus. Nat. Hist., Univ. Kansas, 68:14, July.

Incorrect spelling.

1990. Sorex bairdii permiliensis Carraway, Spec. Publ., The Mus., Texas Tech. Univ., 32:40.

Incorrect use of terminal "i" on bairdi.

Holotype.--USNM 91048, original number 4756; adult male collected 2 October 1897 from "Permilia [=Pamelia] Lake, west base of Mount Jefferson, Cascade Range, Oregon" by J. A. Loring. Type specimen not examined.

Distribution.--Sorex b. bairdi occurs along the Pacific Coast in northwestern Oregon, east into the Coast Range and along the Columbia River into Multnomah and Clackamas counties, and south as far as Benton County. Sorex b. permiliensis occurs in the Cascade Range of Oregon from the Columbia River south to southern Lane County.

Comparisons and remarks.--Sorex monticolus setosus is the nearest subspecies of S. monticolus geographically with which S. bairdi could be confused. S. m. setosus occurs primarily in western Washington and British Columbia but includes a narrow zone in northwestern Oregon where it presumably crossed the Columbia River during a landslide about 1700 years ago (Carraway, 1990). In this region of Oregon, S. m. setosus and S. bairdi are sympatric.

Identification of specimens to either species requires use of a combination of characters:

1. breadth at U4-U4 (1.78388) + greatest length of skull (0.70523) + height of coronoid process (2.55808) - 26.9517 = discriminant score

where, if score < 0.51 = S. monticolus setosus

if score > 1.15 = S. bairdi

2. maxillary breadth (0.60984) + cranial breadth (-1.60895) + breadth at M2-M2 (-0.32228) + breadth at U1-U1 (-1.76036) + breadth at U4-U4 (3.07972) + width across I1-I1 (1.10145) + length of unicuspid toothrow (-0.30093) + greatest length of skull (0.70490) + length of mandible (0.70595) + length of mandibular toothrow (-0.06265) + height of coronoid process (3.31541) + coronoid process-condylar length (-0.54096) - 22.4232 = discriminant score

where, if score < 0.66 = S. monticolus setosus

if score > 1.23 = S. bairdi

By use of the first formula, with three characters, 90.6% of specimens can be classified to species (Fig. 4c) and by use of the second formula, with 12 characters, 94.3% of specimens can be classified to species. In either instance, if the discriminant score of an individual is between the two values presented, it is not possible to assign it to either taxon.

In the original description of S. bairdi (Merriam, 1895), only one terminal "i" was used. This remained until 1977 when it was spelled with both one terminal "i" (Hennings and Hoffmann, 1977:4,6,14,18,26) and two terminal "i's" (Hennings and Hoffmann, 1977:10,15,19). Carraway (1990) employed two "i's" when she resurrected the species S. bairdii. Because only one terminal "i" was used in the original description, I am returning the name to its original and correct spelling. In an attempt to avoid perpetuating the spelling error, I have used S. bairdi throughout the text except when referring to a specific taxonomic treatment in which two "i's" were used.

Specimens examined.--OREGON (110).--Clackamas Co.: 2 mi N, 4 mi E Sandy, 1 (CRCM 89-1508); 3.5 mi N, 8 mi E Sandy, 1 (CRCM 89-1456); 5 mi S, 6 mi E Sandy, 2 (CRCM 89-1583, 89-3016); 6 mi S, 7 mi E Sandy, 6 (CRCM 89-1532, 89-1533, 89-1535, 89-1541, 89-1546, 89-1547); 5 mi S, 6 mi E Zigzag, 3 (CRCM 89-1597, 89-1609, 89-1610); 9 mi N, 4 mi W Zigzag, 4 (CRCM 89-1678, 89-1684, 89-1685, 89-1691); Lane Co.: 1.5 mi N, 2 mi E Blue River, 1 (CRCM 89-2668); 2.5 mi S, 7 mi E Blue River, 1 (CRCM 89-2858); McKenzie Bridge, 1 (PSM 8115); 4 mi N Nimrod, 2 (CRCM 89-2356, 89-2357); T15S, R5E, NW1/4 Sec. 24, 7 (PSM 19870, 19872, 19873, 19875, 19877--19879); T15S, R5E, NW1/4 Sec. 28, 1 (PSM 19701); T15S, R5E, NE1/4 Sec. 32, 14 (PSM 19681, 19684--19687, 19689, 19840, 19848, 19849, 19851, 19852, 19854, 19856, 19858); T15S, R5E, SW1/4 Sec. 33, 3 (PSM 19767, 19768, 19774); T16S, R4E, NE1/4 Sec. 24, 3 (PSM 16729, 19738, 19759); T16S,

R5E, SW1/4 Sec. 6, 5 (PSM 19690, 19691, 19693, 19695, 19881);
Lincoln Co.: Cascade Head Experimental Forest, 15 (PSM 4419, 13587, 13588, 13590, 13591, 13594, 13595, 13597, 13726, 13741, 14451--14454, 14458); Linn Co.: 7 mi N, 8 mi E Blue River, 1 (CRCM 89-1948); 8 mi N, 8 mi E Blue River, 2 (CRCM 89-2184, 89-2185); 9 mi N, 8 mi E Blue River, 3 (CRCM 89-2596--89-2598); T14S, R6E, NE1/4 Sec. 20, 3 (PSM 19816, 19818, 19820); T14S, R6E, NE1/4 Sec. 28, 1 (PSM 19860); T15S, R5E, SE1/4 Sec. 11, 4 (PSM 19835--19838); T15S, R5E, SW1/4 Sec. 13, 5 (PSM 19795, 19800--19802, 19807); T15S, R6E, NE1/4 Sec. 7, 8 (PSM 19706--19708, 19711, 19824, 19829, 19830, 19831); Trout Creek Forest Camp, T13S, R3E, NW1/4 Sec. 32, 2 (PSM 11974, 11975); Multnomah Co.: 10 mi N, 10 mi E Sandy, 3 (CRCM 89-1280, 89-1283, 89-1284); 10 mi N, 11 mi E Sandy, 1 (CRCM 89-1324); 10 mi N, 1.5 mi W Zigzag, 1 (CRCM 89-1146); Tillamook Co.: 2 mi upstream Miami River, 1 (PSM 7003); Netarts, 1 (PSM 3339); Netarts Summit, 1 (PSM 6308); 1 mi E Netarts, 2 (PSM 3341, 3342); Tillamook, 1 (PSM 8116).

Sorex monticolus Merriam

Dusky Shrew

1890. Sorex monticolus Merriam, N. Amer. Fauna, 3:43, September.

1895. Sorex vagrans monticola Merriam, N. Amer. Fauna,
 10:69, December.

1895. Sorex obscurus Merriam, N. Amer. Fauna, 10:72, December.

1977. Sorex monticolus Hennings and Hoffmann, Occas.

Papers Mus. Nat. Hist., Univ. Kansas, 68:2, July.

Sorex monticolus alascensis Merriam

1895. Sorex obscurus alascensis Merriam, N. Amer. Fauna, 10:76.

1900. Sorex glacialis Merriam, Proc. Washington Acad. Sci., 2:16,
March.

1900. S[orex]. alascensis Merriam, Proc. Washington Acad. Sci.,
2:18, March.

1901. Sorex glacialis alascensis Elliot, Field Columb. Mus., Publ.
45, Zool. Ser., 2:372.

1902. Sorex obscurus alascensis Allen, Bull. Amer. Mus. Nat. Hist.,
16:229.

1955. Sorex vagrans alascensis Findley, Univ. Kansas Publ., Mus.
Nat. Hist., 9:41, December.

1977. S[orex]. m[onticolus]. alascensis Hennings and Hoffmann,
Occas. Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.

Holotype.--USNM 73539; young adult male collected 10 July 1895
from "Yakutat Bay, Alaska" by C. P. Streater. Type specimen
examined and measured.

Distribution.--Sorex m. alascensis occurs in a narrow band in
southeastern Alaska from western Prince William Sound, southeast

along the Pacific Coast through Glacier Bay to approximately Juneau and Admiralty Island. The southwestern edge of the northwestern triangle of British Columbia is included in the range of S. m. alascensis (Fig. 13).

Subspecific comparisons and remarks.--Sorex m. alascensis and S. m. elassodon are adjacent taxa south of Admiralty Island, along Frederick Sound, and probably south of Glacier Bay along Cross Sound, Icy Strait, and Chatham Strait. Hall (1981) did not include Chichagof or Kruzof islands in the distributions of these taxa. No shrews from either of these islands were examined in this study so I am unable to assess their subspecific status. S. m. alascensis averages significantly larger than S. m. elassodon in all characters included in this analysis except greatest length of skull; there was no significant difference between the means of this character (Table 2).

Sorex m. alascensis and S. m. longicaudus are adjacent taxa in a narrow band across southeast Alaska in the vicinity of Taku Inlet and Stephens Passage. The only apparent zone of intergradation between these two subspecies includes the region near Juneau and Holkham Bay. Stephens Passage separates these taxa along most of their adjacent distributions. S. m. alascensis averages smaller than S. m. longicaudus in some characters and larger in others (Table 2).

Sorex m. alascensis and S. m. obscurus are adjacent taxa in Alaska from just north of Prince William Sound southeast to




Fig. 13.--Distribution of localities for *S. m. alascensis* (squares) and *S. m. shumaginensis* (circles) specimens included in this analysis representing 63 and 52 unique localities, respectively.

northwestern British Columbia. The zone of intergradation between these two subspecies includes Valdez Narrows, Alaska, a region south of Tats Lake, British Columbia, and the area near Haines, British Columbia. Shrews from Tats Lake had strong affinities with S. m. obscurus and did not seem to exhibit significant intergradation. This results in a relatively narrow band of S. m. alascensis and probably some intergradation along the coastline near Yakutat Bay. S. m. obscurus averages significantly smaller than S. m. alascensis in all characters (Table 2).

Sorex m. alascensis and S. m. shumaginensis are adjacent taxa north and west of Prince William Sound, Alaska. The zone of intergradation between these two subspecies includes Seward, Soldotna, Hope, Anchorage, Coghill Point, and Wasilla, Alaska. S. m. shumaginensis tends to be smaller than S. m. alascensis (Table 2).

Specimens examined.--ALASKA (252).--head of Seymour Canal, Admiralty Island, 1 (UAM 14449); Alaganik, Copper River Delta, Chugach National Forest, 1 (OSUFW X1736); Alaganik, Pete Dahl Cutoff Cabin, Copper River Delta, Chugach National Forest, 4 (OSUFW X1711, X1713, X1715, X1719); Alaganik Cabin, Copper River Delta, Chugach National Forest, 5 (OSUFW X1722, X1723, X1725--X1727); Pete Dahl Cutoff Cabin, Copper River Delta, Chugach National Forest, 8 (OSUFW X1729--X1732, OSUFW 7521--7524); Million Dollar Bridge, Copper River, Chugach Mts., 1 (OSUFW X1710); unnamed creek near terminus Sheridan Glacier, 22 km E Cordova, 4 (UAM 14342, 14344--14346);

Martin River Slough, 55 mi S Cordova, 2 (CRCM 53-316, 55-317); Coghill Point, Port Wells, 85 mi NW Cordova, 2 (CRCM 53-320, USNM 298425); Eagle Creek, Juneau, Douglas Island, 3 (USNM 290264, 290265, 290267); Nederstadt-Lindstrom's site, Dry Bay, 4 (CRCM 85-169, 85-341--85-343); Sandspit 1, Dry Bay, 1 (CRCM 85-344); Sandspit 2, Dry Bay, 18 (CRCM 85-346--85-351, 85-353, 85-355, 85-356, 85-358--85-361, 85-363--85-366, 85-368); Astrolabe Bay, Glacier Bay National Monument, 4 (KU 136812, 136814, 136830, 136843); Bartlett Cove, Glacier Bay, 13 (MVZ 339, 342, 345, 350, 352, 353, 357, 358, 360, 397, 405, 406, 435); Bartlett Cove, Glacier Bay National Monument, 1 (USNM 506565); Boussole Lake, Glacier Bay National Monument, 3 (KU 136818, 136820, 136824); Boussole Muskeg, Glacier Bay National Monument, 1 (KU 136819); Doame River Forest, Glacier Bay National Park & Preserve, 3 (CRCM 85-403--85-405); Dundas Bay, Glacier Bay National Monument, 1 (KU 137693); East River Cabin, Glacier Bay National Park & Preserve, 1 (CRCM 85-398); Gustavus, Glacier Bay National Monument, 1 (KU 137692); Point Gustavus, Glacier Bay, 1 (USNM 97709); Inner Point, Glacier Bay National Monument*, 7 (KU 136804--136807, 136809--136811); Lester Island, Glacier Bay National Monument, 1 (KU 136836); North of Lituya Bay, Glacier Bay National Monument, 3 (KU 136851, 136858, 137685); South of Lituya Bay, Glacier Bay National Monument, 3 (KU 136854, 136871, 137686); Ranney's Property, Glacier Bay National Park & Preserve*, 3 (CRCM 85-399, 85-400, 85-402); Torch Bay, Glacier Bay National Monument, 12 (KU 136828, 136829, 136832, 136837, 136841, 136842, 136861--136864, 136873, 137694); Young Island, Glacier Bay National

Monument, 1 (KU 136835); 1 mi S Haines, 10 (KU 28499, 28500, 28502--28504, 28506, 28509, 28513, 28516, 28517); 4 mi N, 9 mi W Haines, east side Chilkat River, 8 (KU 28476--28478, 28480, 28481, 28484, 28495, 28497); 7 mi SSE Haines, Chilkat Peninsula, 2 (KU 42599, 42600); 2.0 mi from Bar light, Holkham Bay, 1 (MVZ 112104); Juneau, 24 (USNM 74386, 74388, 74389, 74391, 74393, 74395--74397, 74399, 74690, 97551, 203067, 235774, 235775, 235780--235782, 235784, 235785, 235790, 235791, 235794, 235797, 235798); Gold Creek, Juneau, 5 (UIMNH 149--152, USNM 290269); Mendenhall River, 1 (USNM 235253); Camp No. 2, Muir Inlet, 2 (USNM 506970, 506971); Nunatak Camp, Muir Inlet, 1 (USNM 506972); Orca, 1 (USNM 97991); Head of Cordova Inlet, Prince William Sound, 1 (MVZ 527); Eleanor Island, Prince William Sound, 2 (MVZ 859, 860); North end, Elrington Island, Prince William Sound, 1 (MVZ 710); East end Green Island, Prince William Sound, 1 (MVZ 663); East side of Canoe Passage, Hawkins Island, Prince William Sound, 3 (MVZ 538, 539, 543); NE Bay, Hinchinbrook Island, Prince William Sound, 6 (MVZ 571, 573, 596, 604, 927, 940); Galena Bay, Prince William Sound, 1 (CRCM 56-366); Drier Bay, Knight Island, Prince William Sound, 7 (MVZ 781, 790--793, 799, 804); Herring Bay, Knight Island, Prince William Sound, 1 (MVZ 839); La Touche P.O., La Touche Island [=Latouche], Prince William Sound, 1 (MVZ 948); Hanning Bay, Montague Island, Prince William Sound, 8 (MVZ 635, 642, 644, 648, 649, 651--653); Patton Bay, Montague Island, Prince William Sound, 1 (OSUFW X2121); San Juan Bay, Montague Island, Prince William Sound, 9 (OSUFW X2124, X2125, X2127--X2133); Zaikof Bay, Montague Island, Prince William Sound, 3

(MVZ 613, 615, USNM 137329); Valdez Narrows, Prince William Sound, 5 (MVZ 884, 895, 901, 902, 909); NE end Sullivan Island, 7 (KU 42604--42610); Sawmill Bay, Valdez, 3 (CRCM 53-324, USNM 298426, 298427); Yakutat, 6 (USNM 73532, 73536, 73538, 73541, 73543, 97989); Antler Creek, Yakutat, 1 (CRCM 85-418); Cannon Beach, Yakutat, 6 (CRCM 85-420--85-425); Yakutat Airport, 3 (CRCM 85-406, 85-407, 85-409); Yakutat Bay, 1 (USNM 73539); Yakutat Dump Forest, 7 (CRCM 85-410--85-413, 85-417, 85-426, 85-427).

Sorex monticolus calvertensis Cowan

1941. Sorex obscurus calvertensis Cowan, Proc. Biol. Soc. Wash., 54:103, July.
1955. Sorex vagrans calvertensis Findley, Univ. Kansas Publ., Mus. Nat. Hist., 9:39, December.
1977. S[orex]. m[onticolus]. calvertensis Hennings and Hoffmann, Occas. Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.

Holotype.--RBCM 1947; adult male collected 14 July 1937 from "Safety Cove, Calvert Island, British Columbia" by T. T. McCabe and E. B. McCabe. Type specimen examined and measured.

Distribution.--Sorex m. calvertensis occurs on two widely separated islands, Banks Island and Calvert Island, off the coast of British Columbia (Fig. 14).




Fig. 14.--Distribution of localities for *S. m. calvertensis* (triangles and solid arrows), *S. m. insularis* (squares and open arrow), *S. m. elassodon* (circles), and *S. m. prevostensis* (asterisks and hatched arrow) specimens included in this analysis representing 5, 3, 55, and 3 unique localities, respectively.

Subspecific comparisons and remarks.--This taxon is nearly identical morphometrically with S. m. elassodon. The large distance separating the two islands known to be occupied by S. m. calvertensis contributes to the inclination to question the validity of this taxon. Both the summer and winter pelages of these shrews, however, are much paler than that of S. m. elassodon or S. m. longicaudus. Because of the small sample size of S. m. calvertensis included in this analysis, a conservative interpretation was employed, and this taxon is recognized as a distinct subspecies although further research may indicate otherwise.

Ranges of S. m. calvertensis and S. m. longicaudus are adjacent along Principe Channel east of Banks Island, and along Fitzhugh Sound east of Calvert Island. The shrews from Banks and Calvert islands demonstrated some weak affinities with S. m. longicaudus. Morphometrically, S. m. calvertensis is more similar to S. m. elassodon than to S. m. longicaudus. S. m. calvertensis averages smaller than S. m. longicaudus in all characters (Table 2).

Specimens examined.--British Columbia (26).--Banks Island, 2 (RBCM 4479, 4480); Larsen Harbor, Banks Island, 9 (RBCM 4469--4477); Safety Cove, Calvert Island, 6 (CNM 15130, 15136, RBCM 1947, UBC 833, 834, USNM 287817); 1.5 mi W Safety Cove, Calvert Island, 2 (RBCM 1948, 1959); 5 mi W Safety Cove, Calvert Island, 7 (RBCM 1950, 1951, 1954, 1956, 1961, 1963, 1965).

Sorex monticolus elassodon Osgood

1901. Sorex longicauda elassodon Osgood, N. Amer. Fauna, 21:35, September.
1905. Sorex obscurus elassodon Elliot, Field Columb. Mus., Publ. 105, Zool. Ser., 6:450.
1955. Sorex vagrans elassodon Findley, Univ. Kansas Publ., Mus. Nat. Hist., 9:40, December.
1977. S[orex]. m[onticolus]. elassodon Hennings and Hoffmann, Occas. Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.

Holotype.--USNM 100597, original number 1030; young adult male collected 13 June 1900 from "Cumshewa Inlet, Moresby Island, Queen Charlotte Islands, British Columbia" by W. H. Osgood and E. Heller. Type specimen examined and measured:

Distribution.--Sorex m. elassodon occurs from Baranof, Kupreanof, and Mitkof islands, Alaska, southward through the Pacific coastal islands of Alaska primarily west of Stikine and Clarence straits (including Prince of Wales, Forrester, and San Fernando islands). The distribution extends south into British Columbia to include two isolated island groups, Porcher and Dewdney islands (from the Estevan group), as well as the Queen Charlotte Islands (excluding Kunghit Island) of British Columbia (Fig. 14).

Subspecific comparisons and remarks.---Sorex m. elassodon and S. m. longicaudus are adjacent taxa along the eastern reach of Frederick Sound, Stikine Strait, and along Clarence Strait, Alaska. Shrews from Dewdney and Porcher islands are referable to S. m. elassodon. These islands are adjacent to islands inhabited by S. m. longicaudus along the Pacific Coast of British Columbia. Campania Island is directly east of the Estevan group of islands (including Dewdney Island) and shrews there exhibited some affinities with both S. m. longicaudus and S. m. elassodon. Shrews on Prince of Wales Island, Alaska, also exhibited some affinities with S. m. longicaudus but they remain referable to S. m. elassodon. S. m. elassodon averages significantly smaller than S. m. longicaudus in all characters (Table 2).

The ranges of S. m. elassodon and S. m. malitiosus are separated by Summer Strait, Sea Otter Sound, and Iphigenia Bay, Alaska. Coronation Island is relatively isolated from all other islands and shrews from there were classified strongly as S. m. malitiosus. Warren Island is much closer to the islands inhabited by S. m. elassodon and shrews from there exhibited some affinities with S. m. elassodon but remain referable to S. m. malitiosus. S. m. elassodon averages significantly smaller than S. m. malitiosus in most characters (Table 2).

The ranges of S. m. elassodon and S. m. prevostensis are separated by the Houston Stewart Channel of the Queen Charlotte Islands, British Columbia. No intergradation between these two subspecies was indicated by the classifications of these shrews.

Shrews in the S. m. prevostensis a priori group misclassified in the 12-group discriminant analysis were more similar to S. m. longicaudus than to S. m. elassodon. S. m. prevostensis tends to be larger than S. m. elassodon in all characters (Table 2).

Specimens examined.--ALASKA (36).--Forrester Island, 1 (USNM 110398); Kupreanof Island, 14 (USNM 126852--126854, 127075, 127078--127083, 127085--130837, 130839); Scow Bay, Kupreanof Island, 1 (USNM 203068); Mitkof Island, 1 (USNM 130834); Petersburg, Mitkof Island, 6 (USNM 126842, 126844, 126845, 126847, 126848, 126855); Kasaan Bay, Prince of Wales Island, 8 (USNM 126951, 126953, 126957, 126960, 126962, 127547--127549); Klawak Lake, Prince of Wales Island, 1 (USNM 203070); Thorne Harbor, Prince of Wales Island, 2 (UMMZ 106575, 106576); Point Amargura, San Fernando Island, Alexander Archipelago, 2 (UAM 15388, 15389).

BRITISH COLUMBIA (142).--Estevan Group, Dewdney Island, 6 (RBCM 3135--3138, 3140, 3141); Freeman Pass, Porcher Island, 4 (RBCM 4481, 4482, 4484, 4485); Bischoff Island, Queen Charlotte Islands, 3 (UBC 8364, 8413, 8414); Frederick Island, Queen Charlotte Islands, 2 (UBC 2069, 2229); Tasu Inlet, Gowing Island, Queen Charlotte Islands, 1 (UBC 8411); Graham Island, Queen Charlotte Islands, 1 (RBCM 9811); Bradley's Ranch, Graham Island, Queen Charlotte Islands*, 2 (UBC 2101, 2102); Juskatla, [Graham Island], Queen Charlotte Islands, 1 (UBC 7834); Masset, [Graham Island], Queen Charlotte Islands, 4 (CNM 3682, UBC 2098, 7799, 8426); Massett [Masset, Graham Island], Queen Charlotte Islands, 1 (USNM 35597); South of Masset, [Graham Island],

Queen Charlotte Islands, 10 (RBCM 15236, 15238, 15249, 15263, 15268, 15269, 15271--15273, 15279); Waton Creek, 9 mi S Masset, [Graham Island], Queen Charlotte Islands, 1 (UBC 7781); McClinton County, Graham Island, Queen Charlotte Islands, 4 (USNM 290731--290734); Meares Point, Graham Island, Queen Charlotte Islands, 3 (UBC 2091--2093); 11 mi N Port Clements, [Graham Island], Queen Charlotte Islands, 2 (UBC 7726, 7727); Queen Charlotte, [Graham Island], Queen Charlotte Islands, 1 (UBC 2105); Skidgate, Graham Island, Queen Charlotte Islands, 1 (USNM 100614); Gold Creek, Tlell-Port Clements, [Graham Island], Queen Charlotte Islands, 1 (UBC 7869); White Creek, Graham Island, Queen Charlotte Islands*, 2 (UBC 2094, 2096); Yakoun River, Graham Island, Queen Charlotte Islands, 1 (RBCM 9362); Helgesen Island, Queen Charlotte Islands, 5 (RBCM 6527--6531); Hibben Island, Queen Charlotte Islands, 2 (UBC 8229, 8440); Hidden Island, Queen Charlotte Islands, 1 (UBC 7930); Hippa Island, Queen Charlotte Islands, 6 (UBC 8175, 8177, 8263, 8266, 8268, 8271); Kunga Island, Queen Charlotte Islands, 2 (UBC 8243, 8262); Langara Island, Queen Charlotte Islands, 2 (UBC 436, 2088); Andrews Point, Langara Island, Queen Charlotte Islands, 1 (UBC 2089); Henslun [=Henslung, Langara Island], Queen Charlotte Islands, 1 (UBC 8438); Henslung Bay, Langara Island, Queen Charlotte Islands, 20 (UBC 2071, 2073, 2075, 2076, 2080--2085, 2125, 7791--7793, 7795, 8203--8206, 8439); Skidgate Inlet, Legace Island, Queen Charlotte Islands, 1 (UBC 8410); Lina Island, Queen Charlotte Islands, 1 (UBC 7895); Lyell Island, Queen Charlotte Islands, 1 (UBC 8412); Mackenzie Island, Queen Charlotte Islands, 1 (UBC 2225); Main Bolkus Island, Queen

Charlotte Islands, 2 (UBC 7987, 7988); Maude Island, Queen Charlotte Islands, 1 (UBC 7847); Moresby Island, Queen Charlotte Islands, 2 (UBC 2219, 2221); Cumshewa Inlet, Moresby Island, Queen Charlotte Islands, 20 (FMNH 24279, USNM 100586, 100588--100590, 100593, 100595--100597, 100599--100608, 107251); Harriet Harbour, [Moresby Island], Queen Charlotte Islands, 1 (UBC 7871); Sandspit, Moresby Island, Queen Charlotte Islands, 1 (UBC 2106); Tasu, [Moresby Island], Queen Charlotte Islands, 4 (UBC 8230, 8264, 8267, 8270); Ogilvie Island, Queen Charlotte Islands, 2 (UBC 2222, 2223); Queen Island, Queen Charlotte Islands, 1 (UBC 2228); Ramsay Island, Queen Charlotte Islands, 8 (UBC 8172, 8173, 8201, 8202, 8211--8214); Saunders Island, Queen Charlotte Islands, 2 (RBCM 6534, 6535); Thurston Harbour, Talunkwan Island, Queen Charlotte Islands, 1 (UBC 7948); Thurston Harbour, [Talunkwan Island], Queen Charlotte Islands, 2 (UBC 8208, 8209).

Sorex monticolus insularis Cowan

1941. Sorex obscurus insularis Cowan, Proc. Biol. Soc. Wash., 54:103, July.
1955. Sorex vagrans insularis Findley, Univ. Kansas Publ., Mus. Nat. Hist., 9:39, December.
1977. S[orex]. m[onticolus]. insularis Hennings and Hoffmann, Occas. Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.

Holotype.--RBCM 3110; adult female collected 24 August 1938 from "Smythe Island, Bardswell group, British Columbia" by T. T. McCabe and E. B. McCabe. Type specimen examined and measured.

Distribution.--Sorex m. insularis is restricted to the western portion of the Bardswell group of islands off the Pacific Coast of British Columbia including Reginald, Smythe [=Athlone], and Townsend Islands (Fig. 14).

Subspecific comparisons and remarks.--Ranges of S. m. insularis and S. m. longicaudus are separated by the Seaforth Channel to the north, Gale Passage to the east, and Waskesiu Passage to the south. Dufferin and Horsfall islands are part of the Bardswell group but are separated from islands inhabited by S. m. insularis by Gale Passage; shrews from Dufferin and Horsfall islands have been considered S. m. longicaudus. In this analysis, 72.7% ($n = 11$) of the shrews from these islands had classification probabilities of $>77\%$ as S. m. longicaudus; however, three individuals were classified as S. m. insularis. Gale Passage separates Smythe Island, the eastern-most island occupied by S. m. insularis, and Dufferin Island, but is relatively narrow in some places and probably does not eliminate all gene flow. Some of the 19 shrews from Smythe Island included in this analysis were classified as S. m. longicaudus, some as S. m. insularis, and several had intermediate classifications. The latter individuals might be intermediates between S. m. insularis and S. m. longicaudus. S. m.

insularis averages smaller than S. m. longicaudus in some characters and larger in others (Table 2).

Specimens examined.--BRITISH COLUMBIA (54).--Milbanke Sound, Reginald Island, Bardswell Group, 8 (RBCM 3126--3129, 4488--4491); Milbanke Sound, Smythe [=Athlone] Island, Bardswell Group, 19 (RBCM 3103, 3104, 3107, 3109--3113, 3116--3118, 3120--3122, 3125, 4496, 4497, 4498, USNM 287819); Milbanke Sound, Townsend Island, Bardswell Group, 27 (RBCM 3082--3085, 3087--3091, 3093, 3094, 3096, 3098, 3099, 4501--4513).

Sorex monticolus isolatus Jackson

1922. Sorex obscurus isolatus Jackson, J. Washington Acad. Sci., 12:263, June.
1955. Sorex vagrans isolatus Findley, Univ. Kansas Publ., Mus. Nat. Hist., 9:38, December.
1977. S[orex]. m[onticolus]. isolatus Hennings and Hoffmann, Occas. Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.

Holotype.--USNM 177719, original number 517; adult male collected 21 May 1911 from the "Mouth of Millstone Creek, Nanaimo, Vancouver Island, British Columbia" by F. A. Wetmore. Type specimen examined and measured.

Distribution.--S. m. isolatus occurs on Vancouver Island, British Columbia, Denman Island, on the eastern coast of Vancouver Island, and on many small islands along the western coast of Vancouver Island. Whidbey Island also may be inhabited by S. m. isolatus (Fig. 15).

Subspecific comparisons and remarks.--In an analysis of S. monticolus and S. vagrans, George and Smith (1991:87) indicated that the shrews from the San Juan Islands "approach S. monticolus in cranial size" in a univariate analysis and that "most of the Gulf and San Juan Island populations segregate from other S. vagrans centroids and ordinate toward the S. monticolus centroids" in a multivariate approach. Even though they were larger in some of their characters than either Vancouver Island or mainland samples of S. vagrans, George and Smith (1991) found that shrews from San Juan, Lopez, Pender, Saturna, and Samuel islands were more similar to S. vagrans than to S. monticolus and suggested that S. monticolus does not occur on these islands. The same authors also examined 10 shrews from Whidbey Island, Washington; they classified nine of these shrews as S. vagrans and one as S. monticolus. They concluded that more shrews from Whidbey Island needed to be examined before drawing conclusions regarding the presence of S. monticolus on Whidbey Island.

In this study, only one shrew from Whidbey Island was examined and it appeared to be S. monticolus. In the discriminant analysis between S. m. isolatus and S. m. setosus this individual had a




Fig. 15.--Distribution of localities for *S. m. isolatus* (squares) and *S. m. setosus* (circles) specimens included in this analysis representing 59 and 150 unique localities, respectively.

strong classification probability (89%) of being S. m. isolatus. I have referred this individual to S. m. isolatus; however, I agree with George and Smith (1991) that the presence of S. monticolus on Whidbey Island may be open to question.

Ranges of S. m. isolatus and S. m. setosus are adjacent east and south of Vancouver Island, British Columbia. They are separated by the Queen Charlotte Strait, Johnstone Strait, Strait of Georgia, and the Strait of Juan de Fuca. Shrews from Sonora, Maurelle, Stuart, Cortes, Bowen, and Marina islands were classified strongly as S. m. setosus in this analysis. S. m. isolatus averages significantly smaller than S. m. setosus in all characters (Table 2).

Specimens examined.--BRITISH COLUMBIA (151).--East Bunsby, Bunsby Island, 6 (RBCM 5857--5860, 5862, 6245); West Bunsby, Bunsby Island, 3 (RBCM 5863, 5864, 5867); Little Bunsby Island, 1 (RBCM 5865); Cape Scott, Cox Island, 1 (RBCM 5643); Madigan Farm, Denman Island, 2 (RBCM 13702, 13703); 3 mi S, 2.5 mi E Post Office, Denman Island, 5 (TCWC 45664, 45667, 45670, 45671, 45673); Dodd Island, 2 (RBCM 7319, 7320); South Gibraltar Island, 1 (RBCM 8353); Haines Island, 1 (RBCM 7980); Kyuquot Channel, Hohoae Island, 2 (RBCM 6382, 6383); Jaques Island, 1 (RBCM 8338); Nantes Island, 2 (RBCM 8178, 8401); Nettle Island, 1 (RBCM 8322); Onion Island, 2 (RBCM 8358, 8366); Prideaux Island, 1 (RBCM 8328); Reeks Island, 1 (RBCM 8319); Spring Island, 2 (RBCM 6218, 6220); Turtle Island Meadow, 6 (RBCM 8214, 8307, 8308, 8416, 8417, 8420); Tzartus Island, 2 (RBCM 7203, 7207); Holford Bay, Tzartus Island, 2 (RBCM 7209, 7213); Holford

Creek, Tzartus Island, 1 (RBCM 7212); Beaver Creek, Alberni, Vancouver Island, 1 (UBC 6349); Alberni Valley, Vancouver Island, 1 (MVZ 12578); Brooks Peninsula, Vancouver Island, 4 (RBCM 10836, 10838, 10847, 10851); Orchard Point Beach, Brooks Peninsula, Vancouver Island, 2 (RBCM 10844, 10845); Phillips Creek, Buttle Lake, Vancouver Island, 1 (UBC 9284); 3.3 mi NW Cowichan Lake [Vancouver Island], 1 (KU 4279); Comox, Vancouver Island, 4 (CNM 13366, 13367, 13373, USNM 73705); Dudley Marsh, Vancouver Island, 1 (RBCM 12430); Coombs, Dudley Marsh, Vancouver Island, 2 (RBCM 12777, 12778); Qualicum Beach, Dudley Marsh, Vancouver Island, 1 (RBCM 12458); Errington, Vancouver Island, 1 (MVZ 12496); Golden Eagle Mine, Vancouver Island, 2 (MVZ 12582, 12585); Goldstream, Vancouver Island, 5 (MVZ 81083, USNM 71911, 71912, 71914, 71916); SE shore Gooseneck Lake, Vancouver Island, 2 (TCWC 45683, 45684); 9 mi W Holberg, San Josef Campsite, Vancouver Island, 11 (TCWC 45804--45813, 45815); Keogh River, Vancouver Island, 1 (UBC 6348); Long Beach, Vancouver Island, 1 (UBC 6346); Long Beach, 6.5 mi S, 8 mi E Tofino, Vancouver Island, 2 (BYU 9700, 9701); Marble River Recreation Area, Vancouver Island, 14 (TCWC 45364--45367, 45816--45822, 45880, 45883, 45887); mouth Millstone Creek, Nanaimo, Vancouver Island, 2 (USNM 177720, 177721); Nanaimo, Vancouver Island, 1 (USNM 177719); Nanoose Estuary, Vancouver Island, 6 (RBCM 12731, 12747, 12751, 12753, 12765, 12775); Paradise Mine, Vancouver Island, 2 (RBCM 5045, 5047); Parksville, Vancouver Island, 2 (MVZ 12485, 12551); Quatsino, Vancouver Island, 1 (MVZ 77514); Shushartie, Vancouver Island, 1 (CNM 13457); Somenos Lake, Vancouver

Island, 1 (RBCM 12523); Sooke, Vancouver Island, 1 (RBCM 11398); 1 mi SE Thomas Point, Vancouver Island, 1 (UBC 6342); 1 mi E Tofino (near Heidlebrand Ranch), Vancouver Island, 1 (UBC 6345); 1 mi SE Tofino, Vancouver Island, 2 (UBC 6343, 6344); Burnside Road, Victoria, Vancouver Island, 3 (RBCM 11514, 11516, 15826); Mt. Washington, Vancouver Island, 1 (KU 10143); 0.7 mi N Winter Harbour, Vancouver Island, 9 (TCWC 45827--45829, 45831--45835, 45894); 2.9 mi S Zeballos, Vancouver Island, 3 (TCWC 45846, 45847, 45902); 4.4 mi N Zeballos, Vancouver Island, 11 (TCWC 45369, 45371, 45836, 45838, 45839, 45842--45844, 45898--45900); Wickaninnish Island, 1 (RBCM 6658).

WASHINGTON (1).--Island Co.: Deception Pass State Park, Whidbey Island, 1 (TCWC 45849).

Sorex monticolus longicaudus Merriam

1895. Sorex obscurus longicauda Merriam, N. Amer. Fauna, 10:74, December.
1900. S[orex]. longicauda Merriam, Proc. Washington Acad. Sci., 2:16, March.
1901. Sorex obscurus longicaudus Elliot, Field Columb. Mus., Publ. 45, Zool. Ser., 2:372.
1955. Sorex vagrans longicauda Findley, Univ. Kansas Publ., Mus. Nat. Hist., 9:37, December.

1977. S[orex]. m[onticolus]. longicauda Hennings and Hoffmann,
Occas. Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.
First use of name combination, incorrect gender.

Holotype.--USNM 74711, original number 4891; adult male
collected 9 September 1895 from "Wrangel, southeast Alaska" by C. P.
Streator. Type specimen examined and measured.

Distribution.--S. m. longicaudus occurs along the Pacific Coast
of southeast Alaska from Taku Inlet (south of Juneau) south along
the coastline east of Stephens Passage, Frederick Sound, and
Clarence Strait. In British Columbia, the distribution of S. m.
longicaudus extends south along the coastline east of Chatham Sound,
Principe Channel, and Estevan Sound, including Pitt, Campania,
Princess Royal, Swindle, Dufferin, Horsfall, Campbell, and Spider
islands, and the Hunter group of islands, but excludes Porcher
Island, the Estevan group, the western portion of the Bardswell
group, Goose Island, and Hecate Island. The distribution of S. m.
longicaudus continues south to approximately Rivers Inlet. S. m.
longicaudus extends east from Taku Inlet, Alaska, to approximately
the Coast Mountains, east of the Alaska-British Columbia border; it
continues south roughly following the border, west of Meziadin Lake,
Smithers, and Eutsuk Lake to approximately Stuie, Hagensborg, and
Bella Coola (Fig. 16).




Fig. 16.--Distribution of localities for *S. m. longicaudus* (circles) and *S. m. malitiosus* (squares and solid arrow) specimens included in this analysis representing 57 and 2 unique localities, respectively.

Subspecific comparisons and remarks.--In Merriam's (1895) original description of the currently recognized subspecies S. m. longicaudus (Merriam, 1895) the name was spelled "longicauda." When Elliot (1901) returned it to subspecific status he corrected the spelling to "longicaudus" to agree in gender with obscurus. The spelling returned to "longicauda" (appropriately) when Findley (1955) subsumed obscurus into Sorex vagrans. Hennings and Hoffmann (1977) separated S. monticolus and Sorex vagrans but retained the spelling longicauda. The spelling of this name should be "longicaudus" to agree in gender with the masculine Sorex and monticolus. In an attempt to avoid perpetuating the spelling error, I have used longicaudus throughout the text except when referring to a specific taxonomic treatment in which the feminine form was used.

The ranges of S. m. longicaudus and S. m. obscurus are parapatric in southeastern Alaska and western British Columbia from about Taku Inlet, Alaska, southward along the Alaska-British Columbia border, just west of the Coast Mountains, and southward along the Hazelton Mountains of British Columbia. A zone of intergradation between these two subspecies includes the Stikine and Klappan River valleys, Meziadin, Lakelse, and Eutsuk lakes, and Kimsquit, British Columbia. S. m. longicaudus is larger than S. m. obscurus in most characters (Table 2).

Ranges of S. m. longicaudus and S. m. setosus are parapatric in southwestern British Columbia at Rivers Inlet and eastward to approximately Hagensborg and Stuie. All shrews from Rivers Inlet were classified into the taxon with which they were most similar by

use of a diagnosis file generated in the discriminant analysis between S. m. longicaudus and S. m. setosus. Some shrews from Rivers Inlet were classified strongly, and some weakly, as one or the other of the two subspecies. This indicates that this region is part of a zone of intergradation between these two subspecies. Also included in this zone of intergradation are Bella Coola, Hagensborg, and Stuie. S. m. longicaudus averages larger than S. m. setosus in all characters (Table 2).

Specimens examined.--ALASKA (62).--Anan Creek, 1 (USNM 217422); Boca de Quadra, 1 (MVZ 8409); Bradfield Canal, 1 (MVZ 8433); Burroughs Bay, 2 (FMNH 21340, 21342); Helm Bay, 4 (MVZ 451, 452, 458, 471); 2.2 mi from Bar light, Holkham Bay, 1 (MVZ 112102); Game Management Unit 1, Ketchikan, 1 (FMNH 21341); Loring, 7 (USNM 74906, 74914, 74915, 74917, 74918, 74920, 74924); Port Snettisham, 1 (MVZ 8464); Quadra Cannery, 2 (USNM 217416, 526793); Portage Cove, Revillagigedo Island, 2 (MVZ 8419, 8420); Mouth of Stikine River, Sergief Island, 2 (MVZ 30552, 30553); Thomas Bay, 5 (MVZ 444, 8452, 8455, 8458, 8459); Wrangell, 23 (USNM 21749, 74691, 74693--74695, 74697--74699, 74701--74703, 74705--74709, 74711--74713, 74718, 74908, 74910, UW 2059); Game Management Unit 3, Wrangell, 3 (FMNH 21338, 21339, 24277); Wrangell Island, 1 (MVZ 8426); Fools Inlet, Wrangell Island, 2 (MVZ 8428, 8429); Wrangell, Wrangell Island, 3 (MVZ 24562, 24564, 24572).

BRITISH COLUMBIA (191).--Otter Channel, Campania Island, 2 (RBCM 3142, 4514); Ormidale Harbour, Campbell Island, 8 (RBCM 4221--4223,

4225--4228, 4234); Murder Cove, Chatfield Island, 2 (RBCM 4467, 4468); Raven Cove, Chatfield Island, 2 (RBCM 4483, 4487); Don Peninsula, Neekis River, 11 (RBCM 4195, 4196, 4200, 4201, 4203, 4205--4210); Dufferin Island, 3 (RBCM 4235, 4236, 5425); Joassa Channel, Dufferin Island, 1 (UBC 2768); Hot Springs, Eucott Bay, Dean Channel, 1 (CNM 16592); Fitzhugh Sound, Koeve River, 9 (RBCM 3985, 3988, 3989, 3991, 3993--3997); Fitzhugh Sound, Schooner Pass, 2 (RBCM 1945, 1946); Fort [=Port] Simpson, 13 (USNM 90200, 90202, 90208, 90210, 90211, 90213, 90221, 90222, 90225, 90226, 90229, 90231, 90232); Grenville Channel, Lowe Inlet, 9 (RBCM 3151, 3154--3156, 3159, 3160, 3165, 3166, 3168); Hagensborg, Bella Coola River, 6 (CNM 15533, 15534, 15539, 15580, 15803, 15805); Dundivan Inlet, Horsfall Island, 7 (RBCM 4211--4215, 4217, 4218); Hunter Group, B Island, 7 (RBCM 3967--3969, 3971, 3972, 3974, UBC 836); Hunter Group, C Island [=Ruth Island], 2 (RBCM 3970, UBC 835); Hunter Island, 2 (RBCM 4008, 4011); Inverness, mouth Skeena River, 7 (USNM 90215, 90216, 90218, 90235, 90236, 90240, 90242); 1.5 mi S Prince Rupert, Kaien Island, 2 (BYU 9698, 9699); Khutze Inlet, 2 (UMMZ 106581, 106582); Kimsquit, mouth of Dean River, 9 (CNM 16524, 16533, 16547, 16548, 16733, 16756, 16759, 16766, 16775); west Kinahan Island, 1 (UBC 8850); near Prince Rupert, west Kinahan Island, 1 (UBC 8848); Kynoch Inlet, 5 (RBCM 5426--5428, UBC 2766, 2767); Lakelse Lake, 2 (USNM 290724, 290726); Captain Cove, Pitt Island, 1 (RBCM 4486); Union Pass, Pitt Island, 1 (RBCM 3146); 10 mi E Prince Rupert, 2 (UBC 8845, 8849); 12 mi E Prince Rupert, 2 (UBC 8846, 8847); Haque Point, Princess Royal Island, 1 (RBCM 3133); Surf

Inlet, Princess Royal Island, 1 (UBC 6365); Port Belmont, Surf
 Inlet, Princess Royal Island, 2 (UBC 6366, 6367); head of Rivers
 Inlet, 25 (CNM 15038, 15039, 15051, 15069, 15072, 15074--15076,
 15095, 15096, 15202, USNM 90173, 90175, 90180, 90182, 90183, 90186,
 90187, 90192, 90195--90199, 92805); Spider Island, 24 (RBCM 3978--
 3980, 3983, 3998, 4001, 4003, 4006, 4007, 4009, 4010, 4012--4018,
 4020--4022, 4024, 4025, 4027); Stikine River at Great Glacier, 1
 (MVZ 30542); Caribou Mt., near Stuie, 1 (CNM 15674); Junction of
 Atnarko and Whitewater Rivers, Stuie, 3 (CNM 15644, 15651, 15652);
 Meyers Pass, Swindle Island, 1 (RBCM 3131); Wigham Cove, Yeo Island,
 10 (RBCM 4175, 4177, 4186, 4188--4191, 4193, 4492, 4493).

Sorex monticolus malitiosus Jackson

1919. Sorex obscurus malitiosus Jackson, Proc. Biol. Soc.
 Washington, 32:23, April.
1955. Sorex vagrans malitiosus Findley, Univ. Kansas Publ., Mus.
 Nat. Hist., 9:40, December.
1977. S[orex]. m[onticolus]. malitiosus Hennings and Hoffmann,
 Occas. Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.

Holotype.--MVZ 8401, original number 7532; adult female
 collected 21 May 1909 from "east side of Warren Island, Alaska" by
 H. S. Swarth. Type specimen not examined.

Distribution.--S. m. malitiosus is restricted to Coronation and Warren islands in southeast Alaska (Fig. 16).

Subspecific comparisons and remarks.--The shrews from Coronation Island grouped strongly with S. m. malitiosus in the longicaudus-elassodon-malitiosus discriminant analysis employed in this study. The shrews from Warren Island had slightly weaker classifications and tended toward S. m. longicaudus (instead of S. m. elassodon as might be expected on the basis of its proximity) but are still referable to S. m. malitiosus.

Specimens examined.--ALASKA (17).--Egg Harbor, Coronation Island, 11 (MVZ 8388, 8389, 8391--8394, 8396, 8397, UAM 14455--14457); East side Warren Island, 6 (MVZ 8398, 8399, 8400, 8402, 8405, 8406).

Sorex monticolus monticolus Merriam

1890. Sorex monticolus Merriam, N. Amer. Fauna, 3:43, September.

1895. Sorex vagrans monticola Merriam, N. Amer. Fauna, 10:69, December.

1977. S[orex]. m[onticolus]. monticolus Hennings and Hoffmann, Occas. Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.

Holotype.--USNM 17599/24535, original number 406; adult male collected 28 August 1889 from "San Francisco Mountain, Arizona" by C. H. Merriam and V. Bailey. Type specimen examined and measured.

Distribution.--S. m. monticolus is restricted to southeastern Arizona and an isolated population in the San Francisco Mountains (Fig. 17).

Subspecific comparisons and remarks.--Sorex m. monticolus and S. m. obscurus are separated in east central Arizona by the Mogollon Plateau. Their adjacent ranges extend southward approximately along the Arizona-New Mexico border. These two subspecies do not seem to intergrade along the Mogollon Plateau but along the southeastern border of Arizona and New Mexico a zone of intergradation includes the region around Mogollon, New Mexico and the Chiricahua Mountains in Cochise County, Arizona. S. m. monticolus averages smaller than S. m. obscurus in all characters (Table 2).

Specimens examined.--ARIZONA (120).--Apache Co.: Alpine Job Corp.*, 1 (UIMNH 47103); 4 mi S, 16 mi W Alpine, 6 (MSB 40941, 40944, 40946, 40948, 40992, 40993); Apache National Forest, 1 (BSFC 2952); West Fork Black River, 1 (MVZ 60377); near head Burro Creek, White Mts., 1 (USNM 209331); Greer, 9 (UIMNH 54033--54039, 54041, 54042); 1 mi S Greer, 1 (UIMNH 54043); Hall Creek, 3 mi N Greer, 1 (UIMNH 50049); Hall Creek, 3.5 mi N Greer, 2 (UIMNH 54031, 54032); Sheep Crossing, 4.5 mi SW Greer, 6 (UIMNH 54044--54049); Phelps




Fig. 17.--Distribution of localities for *S. m. monticolus* (squares and solid arrows), *S. m. obscurus* (circles), and *S. m. parvidens* (triangles and open arrow) specimens included in this analysis representing 39, 5, and 794 unique localities, respectively.

Botanical Area, 6.5 mi SW Greer, 1 (UIMNH 54050); Sunrise Lake, 7 mi E Greer, 1 (UIMNH 50050); Phelps Botanical Area, 8 mi SW Greer, 1 (UIMNH 50051); Little Colorado River, White Mts., 3 (USNM 158587, 158589, 158590); Sheep Crossing on Little Colorado River, 9 (UIMNH 29058, 29060, 29062--29065, 29067, 29069, 29070); Mount Thomas, White Mts., 1 (USNM 208664); east slope Mount Thomas, White Mts., 2 (USNM 209336, 209337); Springerville, 1 (USNM 24829); Horseshoe Cienega, White River, White Mts., 1 (USNM 209326); Cochise Co.: Fly Park, Chiracahua Mts., 1 (USNM 66090); Long Park, Chiracahua Mts., 1 (UMMZ 77375); Rustler's Park, Chiracahua Mts., Coronado National Forest, 12 (MSB 40091--40093, 40892, 40893, 40895, 40897, 40899, 47301--47304); Coconino Co.: San Francisco Mt., 1 (USNM 17599); Graham Co.: near head Ash Creek, Graham Mts., 1 (USNM 204189); Hospital Flat, Graham Mts., 1 (UIMNH 4774); Marijilda Canyon, Graham Mts., 1 (MVZ 50249); Snow Flat, Graham Mts., 2 (UIMNH 4775, 4776); Greenlee Co.: Hannagan Creek, 8 (MVZ 55193--55196, 55198, 55199, 60378, 60379); Hannagan Meadow, 10 (MVZ 55201, 55203, UIMNH 6037--6044); 0.75 mi NE Hannagan Meadow, 2 (UIMNH 6035, 6036); Horse Cienega, 1 mi NNE Hannagan Meadow, 6 (UIMNH 6028--6032, 6034); Lost Lake Road, 1.5 mi N Hannagan Meadow, 2 (UIMNH 6021, 6023); Lost Lake Road, 2 mi N Hannagan Meadow, 2 (UIMNH 6024, 6025); Cache Cienega, 3 mi SSW Hannagan Meadow, 3 (UIMNH 6045, 6047, 6049); 3 mi S, 2 mi W Hannagan Meadow, 2 (UIMNH 6056, 6057); 4 mi NE Hannagan Meadow, 3 (UIMNH 6016, 6019, 6020); Prieto Plateau, South end Blue Range, 1 (USNM 205838); Rose Peak, 1 (MVZ 55189); Navajo Co.: North Fork White River, White Mts., 10 (SDNHM 10624, 10658, 10678, 10718,

10725, 10732, 21564--21567); Pima Co.: Summerhaven, Santa Catalina Mts., 1 (USNM 244070).

Sorex monticolus obscurus Merriam

1891. Sorex vagrans similis Merriam, N. Amer. Fauna, 5:34, July.

Name preoccupied by Sorex similis Hensel [Neomys similis].

1895. Sorex obscurus Merriam, N. Amer. Fauna, 10:72, December.

1955. Sorex vagrans obscurus Findley, Univ. Kansas Publ., Mus.

Nat. Hist., 9:43, December.

1977. S[orex]. m[onticolus]. obscurus Hennings and Hoffmann, Occas.

Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.

Holotype.--USNM 23525/30943, original number 1670; adult female, collected 26 August 1890 from "Timber Creek, Salmon River Mountains, Idaho" by B. H. Dutcher. Type specimen examined and measured but not included in analysis because of broken skull.

Distribution.--S. m. obscurus occurs from eastern Alaska, south through western Canada including the Yukon Territory, Northwest Territories, British Columbia, Alberta, and southwestern Saskatchewan, but excluding the western and southwestern coastline of British Columbia. The distribution of S. m. obscurus continues southward into eastern Washington and through the Rocky Mountains of the United States, including Idaho, Montana, Wyoming, Utah,

Colorado, Arizona, and New Mexico, and south into Mexico. Several isolated groups of S. m. obscurus occur in mountainous offshoots of the Rockies including the Wallowa, Blue, and Steens mountains of Oregon, the Ruby, Santa Rosa, and Carson mountains of Nevada, and the Sierra Nevada mountains of California (Fig. 17).

Subspecific comparisons and remarks.--Ranges of S. m. obscurus and S. m. setosus are adjacent in southwestern British Columbia and central Washington. A zone of intergradation between these two subspecies in British Columbia includes Anahim Lake, Itcha Mountain, and Chezacut. The Fraser Plateau and the Cascade Range of British Columbia and Washington separates S. m. setosus and S. m. obscurus southward. A zone of intergradation in this region includes Hope and Manning Provincial Park in British Columbia, and Barron, Slate Peak, Stehekin, head of Lake Chelan, Cloudy Pass, Lyman Lake, Wenatchee Lake, Keechelus Lake, Easton, Lester, Corral Pass, Owyhigh Lake, Crystal Mountain, Council Pass, Signal Peak, and southward to approximately the White Salmon River in southern Washington. S. m. obscurus averages smaller than S. m. setosus in some characters and larger in others (Table 2).

Ranges of Sorex m. obscurus and S. m. shumaginensis are adjacent in central Alaska. The zone of intergradation includes Fishhook, head of Toklat River, Maclaren, Glenn Highway at Eureka, Denali National Park, Healy, Kakagrak, and the Noatak Valley in northwestern Alaska. Populations from these localities contained some shrews classified as S. m. shumaginensis and some as S. m.

obscurus. In most cases, these classifications were weak; this suggests that these shrews are different only at the subspecific level and are not sympatric species. The univariate Student-Newman-Keuls multiple-range test indicated that S. m. shumaginensis is significantly smaller than S. m. obscurus in some characters but larger in others (Table 2).

Specimens examined.--ALASKA (57).--Alatna, 1 (USNM 245526); Anaktuvuk Pass, 4 (USNM 290246, 290250, 290257, 290258); Bettles, 5 (KU 43198, USNM 245527--245529, 245533); Chandler Lake, 1 (KU 43197); Chilkoot Lake, 13.5 km NW Haines, 1 (CMNH 70694); Denali Hwy, east of Maclaren, 63°10'N, 146°30'W, 3 (CNM 40276--40278); Denali Hwy, east of Maclaren, 63°08'N, 146°15'W, 3 (CNM 40279--40281); Mts. near Eagle, 7 (USNM 130999, 131014, 131029, 131048, 131069, 131070, 131082); Glenn Hwy, near Eureka Lodge (61°49'N, 147°20'W), 1 (CNM 40282); 1 mi S, 1 mi W Healy, 3 (CMNH 76006--76008); Hogatza, 70 mile (Hughes Quad), 1 (UAM 15318); Kilikmak Valley, large lake ca. 5 mi NE Kakagrak (headwaters Kilikmak Cr.), 1 (USNM 505019); Mt. McKinley Park, 1 (UBC 3573); Richardson, Tanana River, 6 (USNM 234687, 234688, 234693, 234695, 234698, 241790); Savage River, 4 (USNM 244042--244044, 244049); Sawtooth Mts., 1 (USNM 245479); Skagway River Valley, 4 km NE Skagway, 2 (CMNH 70701, 70703); Tanana River, 1 (USNM 38754); Head of Toklat River, 8 (USNM 157191--157193, 157196--157198, 157269, 157271); Wahoo Lake, Brooks Range, 1 (KU 50409); Wells, Chilkat Valley, 1 (USNM 217412); Yukon River, opposite mouth of Charlie Creek, 1 (USNM 293178).

ALBERTA (71).--Akamina Creek, Waterton Lake National Park, 1 (UBC 1681); Beaverlodge, 55°13'N, 119°26'W, 1 (CNM 51474); Belly River, Waterton Lake Park, 1 (CNM 5131); Belly River, Waterton Lake National Park, 3 (UAMZ 2830, 3509, 3521); Blakiston Br., Waterton Lake National Park, 1 (UAMZ 3517); Bryant Creek, Banff National Park, 50°52'N, 115°27'W, 2 (CNM 16174, 16179); Cameron Lake W.P. [=Waterton Lake Park], 1 (UAMZ 1388); Cardston, 2 (UAMZ 1368, 1369); Coleman, Vicary West Ridge, 1 (UBC 8808); 16 mi N Colman [=Coleman], 1 (UAMZ 539); Cottonwood Creek, Jasper National Park, 1 (UAMZ 2832); Crowsnest Pass, 49.4°N, 114.5°W, 1 (UAMZ 1361); Elbow River and Prairie Creek fork, 1 (CNM 25887); Fiddle Creek, (Ath. [=Athabasca] R.) Jasper National Park, 1 (UAMZ 4007); 7 km S, 5 km W Grimshaw, 56°08'N, 117°41'W, 1 (CNM 51473); Henry House, 2 (USNM 75317, 75318); 15 mi S Henry House, Rocky Mts., 4 (USNM 81511, 81512, 92118, 92119); 20 mi N Hotchkiss, 2 (CMNH 75970, 75971); Jasper, 1 (CNM 10786); 1 mi N, 10 mi E Kinuso, Assinean River, 2 (KU 21079, 21080); Kvass Summit, 70 mi NW Jasper (by air), 2 (CMNH 22822, 22825); Lost L Creek x Bauerman Br. Waterton Lakes National Park, 2 (UAMZ 2835, 2842); Maligne Lake, Jasper National Park, 52.4°N, 117.3°W, 4 (UAMZ 4220--4223); Medicine Lake, 14 mi S Buck Lake, 52.4°N, 117.1°W, 1 (UAMZ 4707); Muskeg Creek, 15 mi from mouth, 1 (USNM 81353); Muskeg Creek, 20 mi from mouth, 3 (USNM 81359, 81364, 81369); Muskeg Creek, Fishing Lake; 90 mi N Jasper, 1 (USNM 81349); Rock Lake, 6 mi N of NE boundary Jasper Park, 1 (CMNH 22820); Rodent Valley, 25 mi W Henry House, 1 (USNM 81373); Slave Lake, 3 (UAMZ 941, 942, 1390); Snaring, Jasper National Park, 1 (UAMZ 2837); Spray

Lake, 1 (UAMZ 1366); Sulpher Prairie, 1 (USNM 81401); Sulpher Prairie, Grand Cache River, 2 (USNM 81347, 81348); Sunwapta Pass, Banff National Park, 1 (ROM 22868); N end Tyrell Lake, 1 (UAMZ 5493); Waterton, 4 (UAMZ 557, 1380, 1383, 1385); Waterton Lakes, 1 (ROM 22867); Waterton Lakes Park, 4 (CNM 4654, 4675, 4725, 4734); Waterton Lakes Park, Bertha Creek, 2 (CNM 5087, 5094); Headwaters of West Castle River, 2 (UAMZ 6904, 6905); Whirlpool River, Jasper National Park, 3 (UAMZ 2829, 2833, 2834); Whiskey Gap, 1 (UAMZ 1373).

ARIZONA (12).--Apache Co.: Spruce Creek, Tunitcha Mts., 3 (USNM 227460, 227461, 227466); Tsaile Creek, 9.5 mi SE Lukachukai, 1 (UIMNH 28431); Wheatfield Creek, west slope Tunitcha Mts., 1 (USNM 247800); Coconino Co.: Baker's Butte, 2 (MVZ 55184, 55187); Double Cabin, 6 mi NW Promontory Lookout, 2 (UIMNH 26319, 26320); Sawmill Springs, 8 mi SE Mormon Lake, 1 (MVZ 55183); Vincent Ranch, 39 mi S, 16 mi W Winslow, 1 (UIMNH 33552); 28 mi S, 24 mi W Winslow, 1 (UIMNH 33550).

BRITISH COLUMBIA (151).--Akamina Pass, Alberta border, 1 (UBC 3589); Alpha Lake, 1 (UBC 37); Anahim Lake, 1 (UBC 2630); Apex Mountain, 1 (RBCM 7555); Ashcroft, Hat Creek Valley, 1 (RBCM 9587); Atlin Lake, 1 (RBCM 9895); 6 mi N Babine Trail, Babine Mts., 1 (USNM 202800); Barkerville, 3 (USNM 106220, 106222, 106223); Bear Lake, site of Fort Connolly, 2 (USNM 202802, 202811); Bennett, 4 (USNM 127087, 128208, 128584, 130027); Bennett City, 2 (USNM 99310, 99311); Buffalo Lake, 1 (RBCM 7556); Burnell Lake, 1 (RBCM 15906); Caraboo Lake, 1 (USNM 67432); Mts. near head Chapa-atan River; near

head branch of Stikine River, 1 (USNM 170720); Charlie Lake, 2 (RBCM 2589, 2590); Chezacut Lake, Chilcot, 7 (UBC 6353--6356, 6358--6360); Chilcotin, Itcha Mtn., 1 (UBC 6351); Dease Lake, 58°27'N, 130°02'W, 1 (CNM 35932); Eutsuk Lake, 1 (RBCM 1715); Fort Grahame, 2 (USNM 170722, 170725); NE Fort Nelson, 1 (RBCM 11324); NE of Fort Nelson, 59°25'N, 120°47'W, 3 (RBCM 11319, 11320, 11325); E of Fort Nelson, 58°49'N, 121°22'W, 1 (RBCM 11138); Glacier, 4 (USNM 69134, 69135, 69218, 290716); Golden, 1 (USNM 69141); Haines Road, mile 61, 1 (RBCM 6141); Hazelton, 1 (USNM 202799); Hedley, Stirling Creek, 2 (CNM 8864, 8868); Hope-Princeton Highway, 1 (RBCM 5504); mile 15, Hope-Princeton Highway, 1 (RBCM 5506); mile 20, Hope-Princeton Highway, 1 (RBCM 5505); Hudson's Hope, 1 (USNM 170744); Indian Point Lake, 2 (RBCM 1221, 1224); John Sandy Creek, Pemberton Meadows, 1 (UBC 13308); July Pass, N. Fork Morkill River, 40 mi above mouth, near Snowshoe, 3 (CMNH 19628--19630); Lac du Bois, Kamloops, 1 (UBC 2673); Kaskonook, Kootenay Lake, 1 (RBCM 5097); 6 mi W Kelly Lake, 1 (UBC 13414); North end Kerry Lake, E bank Crooked River, 3 (KU 63902--63904); Klappan River Valley, Tset-ee-yeh [=Tsetia] River, (branch of Klappan River), 1 (USNM 170717); Kootenay National Park, 1 (RBCM 14748); Lac La Hache, 1 (UBC 2633); Laurier Pass, 2 (USNM 256623, 256624); Link Creek, 1 mi S, 5 mi W LeMoray, 1 (CMNH 75974); Manning Park, Allison Pass, 1 (RBCM 5144); Manning Park, Mountain Beaver Valley, 1 (RBCM 5145); Manning Park, Ranger Station, 5 (UBC 3387, 3394--3397); 0.5 mi W Manning Park Lodge, 2 (UBC 9548, 9550); McDame Creek, Dease River; Quartz Creek, 1 (USNM 206099); McDame Post, Dease River, 2 (USNM 206103, 206104); Meziadin Lake, 1 (RBCM

9928); Meziadin Lake, east side near campground, Nass Basin, 1 (PSM 20929); Monashee Pass, 2 (RBCM 2332, UBC 3741); Moose Lake, 2 (USNM 174412, 174414); Mount Gordon, 3 (RBCM 5509--5511); Mount Revelstoke, 1 (USNM 290719); Mount Revelstoke National Park, 1 (USNM 290721); Mount Wardle, Kootenay National Park, 1 (RBCM 14917); North Fork Moose River, 1 (USNM 174402); 9 mi S, 44 mi W Muskwa, 2 (KU 63900, 63901); Nelson, 1 (USNM 69131); 6 mi S Nelson, Silver King Mine, 3 (USNM 66662, 69215, 69216); 165 mi N of Nelson [=Fort Nelson], Alaska Hwy, 58°51'N, 125°44'W, 1 (CNM 17399); Okanagan, 1 (RBCM 955); Okanagan Landing, 1 (RBCM 921); Omineca Mts., 1 (RBCM 4833); Ootsa Lake, 2 (RBCM 1717, 1718); Petitot, 59°44'N, 121°51'W, 1 (RBCM 11134); Purden Lake, 35 mi E Prince George, 2 (CMNH 75987, 75988); Quick, 20 mi E Smithers, 1 (RBCM 9942); Sicamous, 1 (USNM 69319); Smithers, 1600 ft., 1 (TCWC 26980); 10 mi W Smithers, 1 (SUVN 1415); Smoky River, 7.5 mi above mouth, near Snowshoe, 1 (CMNH 19626); Spatsizi Plateau, 1 (RBCM 6548); Spout Lake, 1 (UBC 2999); Stikine River at Great Glacier, 1 (MVZ 30546); Cassiar Dist., 5.5 mi W of Jct. Stonehouse Cr. and Kelsall River, 4 (KU 28534, 28535, 28537, 28545); Summit Lake, mi 392 Alaska Hwy, 1 (CNM 34554); West end Summit Lake, Alaska Hwy mi 393, 1 (KU 63905); Takla Lake, Omineca Mts., 1 (RBCM 4837); Tats Lake, 11 (RBCM 11668, 11679, 11688, 11704, 11719, 11721, 11728, 11740, 11745, 11746, 11752); Tetana Lake, Driftwood River, 1 (RBCM 4648); 3 mi WNW Jct. Trout and Liard Rivers, Hot Springs, 1 (KU 28551); 15 mi N, 5 mi E Trutsh, 1 (CMNH 75991); Tutshi Lake, 1 (RBCM 9937); Vermilion Crossing, Kootenay National Park, 2 (USNM 290727, 290728); Vermilion River, 30

mi N Radium Hot Springs, 1 (USNM 319523); Whiskers Point, 9 mi S, 6 mi E McLeod Lake, 9 (CMNH 75976--75979, 75982--75986); Yellowhead Lake, 1 (USNM 174419).

CALIFORNIA (116).--Amador Co.: Silver Lake, 1 (UMNH 13635); Fresno Co.: Bullfrog Lake, 1 (MVZ 24797); Cascade Valley, 1 (UIMNH 28283); Charlotte Creek, at Bubb's Creek, 1 (UMMZ 106602); Charlotte Dome, Jesse's Woods, 1 (UMMZ 106594); Cottonwood Basin, 2 (UMMZ 106603, 106605); Grouse Creek, T8S, R25E, Sec. 23, 2 (CMNH 84038, 84039); Horse Corral Meadows, 2 (USNM 42195, 42196); Kearsarge Lake, 1 (UMMZ 106591); Kearsarge Basin, Kearsarge Lake, 1 (UMMZ 106587); Kearsarge Basin, Kearsarge Ledge Camp, 1 (UIMNH 23550); Kearsarge Basin Ledges, 1 (UMMZ 106601); Russian Tank Camp, 1 (UMMZ 106596); Sierra Nevada Mts., San Joaquin River, 2 (USNM 42075, 42078); Inyo Co.: 3 mi S, 8 mi W Big Pine, 1 (MVZ 98945); 7 mi W Big Pine, 2 (RHMC 8166, 8917); 10 mi W Big Pine, 4 (KU 1373, 1374, 1376, 1377); 11 mi SW Big Pine, Big Pine Creek Campground, 1 (RHMC 6345); Gilbert Lake, 1 (UMMZ 106586); 1.25 mi S, 5 mi W Independence, 1 (MVZ 98946); near Independence, Little Onion Valley, Sierra Nevada, 1 (MVZ 17774); near Independence, Onion Valley, Sierra Nevada, 1 (MVZ 17778); Independence Creek, Gilbert Lake, 3 (UMMZ 106585, 106599, 106600); Kearsarge Pass, near Independence, Onion Valley, Sierra Nevada, 1 (MVZ 17779); Lone Pine Creek, 1 (MVZ 17783); Lone Pine Creek, 1.25 mi S, 9.5 mi W Lone Pine, 2 (MVZ 98947, 98948); Onion Valley, 4 (UMMZ 106583, 106584, 106597, 106598); Round Valley, 1 (USNM 42411); Sageflat Campground, E side Sierra Nevada Mts., Inyo National Forest, 1 (RHMC 7494); Sierra Nevada Mts., Bishop Creek, 2

(USNM 42070, 42073); Kern Co.: Kern Lakes [=Tulare Co.], 1 (USNM 42410); Onyx, South Fork Kern River, 1 (USNM 108815); Madera Co.: Mount Lyell, 6 (USNM 109529, 109533, 110290--110292, 116022); Mariposa Co.: East Fork Indian Canyon, 3 (MVZ 22026, 22027, 22029); Lake Tenaya, 3 (USNM 108936, 110295, 116023); 1 mi E Merced Lake, Yosemite Park, 2 (MVZ 22990, 22999); near Porcupine Flat, Yosemite Park, 3 (MVZ 22032, 22041, 22042); Mono Co.: Pine City, near Mammoth, 1 (MVZ 32902); Warren Fork of Seevining [=Lee Vining] Creek, 1 (MVZ 23011); Williams Butte, 1 (MVZ 23007); Placer Co.: Robinson Flat Campground, T15N, R13E, SE1/4 Sec. 10, 1 (SDNHM 22519); Tulare Co.: Aster Lake, Sequoia National Park, 1 (MVZ 108976); Big Meadows, Sequoia National Forest, 7 (SDNHM 12007, 12008, 13573, 13575, 13576, 13578, 13581); Cahoon Meadow, Sequoia National Park, 3 (MVZ 108977, 108978, USNM 274873); Halstead Meadows, Sequoia National Park, 1 (USNM 42205); Kaweah River, East Fork, 5 (USNM 42303, 42305--42307, 42309); Kern River, South Fork, 3 (USNM 41630--41632); Little Brush Meadow, Olancho Peak, Sierra Nevada Mts., 1 (MVZ 16278); Moltke [=Mulkey] Meadows, Sierra Nevada Mts., 1 (USNM 42739); Mount Whitney, 3 (USNM 42369, 42550, 42551); Mount Whitney, Whitney Cr., 4 (FMNH 13356--13359); Round Meadow, Sequoia National Park, 2 (MVZ 108980, USNM 274874); Sequoia National Park, 1 (USNM 42202); 3 mi SE Three Rivers, 1 (RHMC 7357); Upper Junston Meadow, Kern River, 2 (MVZ 108982, 108983); Tuolumne Co.: Fletcher [=Fletcher] Creek, near Vojeberry Lake, Yosemite Park, 1 (MVZ 22993); Glen Aulin, Tuolumne River, Yosemite National Park, 1 (MVZ 23015); Near Lyell and Dana Forks Tuolumne River, 2 (CMNH 7185,

7207); Near Mono Meadow, Yosemite Park, 3 (MVZ 22020, 22024, 22025); Mount Dana, 1 (USNM 109268); Tuolumne Meadows, N Base Mount Lyell, 1 (USNM 110289); Tuolumne Meadows, Soda Springs, 1 (USNM 108937); Tuolumne Meadows, Yosemite National Park, 3 (MVZ 22046, 22049, 22054).

COLORADO (280).--Archuleta Co.: 2 mi E, 0.5 mi N Chimney Rock, 1 (MSB 6758); Boulder Co.: Allens Park, 3 (UIMNH 56226--56228); 0.75 mi N, 2 mi W Allens Park, 5 (KU 50312--50316); Boulder, 2 (FMNH 11672, USNM 112096); 5 mi W Boulder, 1 (USNM 137335); Eldora, 1 (USNM 142519); Longs Peak, at timberline, 1 (USNM 73774); Meadow Mtn., 0.5 mi N, 0.5 mi W Allens Park, 2 (UIMNH 51042, 51044); Nederland, 6 (USNM 137656, 137657, 142518, FMNH 11665, 11682, 11683); Ward, 1 (USNM 53941); 3 mi S Ward, 1 (KU 19939); Chaffee Co.: Poncha Creek, 10 mi SW Salida, 5 (KU 113610, 113611, 113613--113615); Saint Elmo, 2 (USNM 150753, 150760); 17 mi W Salida, E side Monarch Pass, 2 (CMNH 15536, 15537); Clear Creek Co.: 1 mi SW Berthoud Pass, 1 (BYU 9695); 1013 Griffith Street, Georgetown, 1 (BSFC 10942); Conejos Co.: 5 mi S, 24 mi W Antonito, 1 (KU 41567); Platoro, 6 (FHSU 6819, 6820, 7164, 7166--7168); 3-5 mi SW Platoro, 13 (FHSU 6815--6818, 7150--7152, 7154--7159); Costilla Co.: Fort Garland, 1 (USNM 47269); Delta Co.: Collbran [=Mesa Co.], 12 mi S [=N], 5.5 mi E Skyway, Grand Mesa, 1 (KU 59667); 12 mi S, 5.5 mi E Collbran, Grand Mesa, 2 (KU 59646, 59647); 0.5 mi S, 8 mi E Skyway, Grand Mesa, 2 (KU 59648, 59650); 1.5 mi S, 8 mi E Skyway, Grand Mesa, 2 (KU 59655, 59665); 2 mi S, 8 mi E Skyway, Grand Mesa, 1 (KU 59653); Dolores Co.: Stoner Mesa, 1 (BSFC 4237); T40N, R13W, Sec.

13, 1 (KU 120928); Fremont Co.: 3.5 mi S Coaldale, 1 (TCWC 4217);
Garfield Co.: Baxter Pass, Book Plateau, 1 (USNM 148157); Henderson
 Ridge, 1 (BSFC 10610); S slope Trapper Creek, 8.5 mi N, 7 mi W
 Rifle, 2 (BSFC 2174, 2175); T3S, R93W, Sec. 11, 2 (BSFC 6675, 6676);
Grand Co.: Fraser Experimental Forest, 3 (BSFC 2915, 2916, 2917);
 Spruce Creek, Fraser Experimental Forest, 1 (BSFC 2919); 1 mi N
 Radium, along Blacktail Creek, 2 (BSFC 6069, 6070); Gunnison Co.:
 Beaver Creek, 7.5 mi W Gunnison, 1 (BSFC 24); Calhoun Line*, 3 (USNM
 485366, 485367, 485370); Dry Gulch and Gunnison River, 2 (UMNH
 17979, 17980); Gothic, 7 (MSB 9068, USNM 303466--303468, 303470--
 303472); 0.25 mi N Gothic, 2 (MVZ 125637, 125638); 0.3 mi NNW
 Gothic, 2 (USNM 485303, 485306); 2.7 mi NW Gothic, 2 (USNM 485258,
 485310); Gunnison, 7.5 mi W Beaver Creek, 1 (BSFC 1335); 5.8 mi NW
 Gunnison; Maggie Gulch, 1 (USNM 485288); 1.5 mi N Rocky Mt.
 Biological Lab., near Gothic, 7 (KU 116710--116716); 8 mi NW
 Sapinero; Black Mesa Experimental Range, 1 (USNM 287590); Willow
 Creek, 0.5 mi above jct. Gunnison River, 1 (UMNH 17981); Huerfano
Co.: 4 mi S Cuchara Camp, 2 (KU 68438, 68441); 5 mi S, 1 mi W
 Cuchara Camp, 8 (KU 59656, 59659--59662, 59664, 59669, 59670);
Jackson Co.: 2 mi N, 2 mi E Gould, 1 (KU 116709); La Plata Co.:
 Columbine Ranger Station, Cascade Creek, 4 (MVZ 60389, 60390, 60392,
 60393); Lake Co.: Halfmoon Creek, 8 mi SW Leadville, 3 (KU 113605,
 113606, 113607); 12 mi S, 1 mi W Leadville, 1 (KU 57884); 3 mi W
 Twin Lakes, 2 (KU 57885, 57886); Larimer Co.: Estes Park, Aspenglen
 Campground, 1 (CMNH 74012); 3 mi S, 3 mi W Estes Park, YMCA Camp, 2
 (UIMNH 7558, 7559); 3.5 mi S, 4 mi W Estes Park Village, 2 (KU

50306, 50309); North Fork Poudre River Campgrd, Roosevelt National Forest, 1 (BSFC 10783); Sheep Creek, Arapaho-Roosevelt National Forest, 2 (BSFC 14461, 14817); Sheep Creek, ca. 8 mi N Red Feather Lakes, Arapaho-Roosevelt National Forest, 10 (BSFC 12625, 12627, 12629--12634, 12636, 13495); Willow Park, Rocky Mt. National Park, 7 (UMMZ 57035, 57036, 57038, 57039, 57042, 57043, USNM 293215); Mesa Co.: T5S, R100W, Sec. 35 [=Garfield Co.], 1 (BSFC 10611); Uncompaghre Plateau, Uncompaghre Butte, 1 (USNM 149974); Mineral Co.: Borns Lake, 3 (MVZ 60383--60385); Mount McLellan [=McClellan], 2 (USNM 137434, 137435); 10.3 mi SW South Fork, 1 (KU 4397); 4 mi S, 6 mi E Wagon Wheel Gap, 1 (KU 120929); Moffat Co.: Beaver Creek, 25 mi N, 49.5 mi W Maybell, 1 (FHSU 25125); 1 mi S Craig, 2 (BSFC 4607, 4608); Maddox Place, Douglas Mt., 3 (BSFC 13069, 13071, 13072); 22.5 mi N, 49.5 mi W Maybell, 1 (FHSU 25767); 24.5 mi N, 49 mi W Maybell, 6 (FHSU 25763--25766, 25770, 25771); Pool Creek Ranch, Dinosaur National Monument, 2 (BSFC 13108, 13109); Streatel Canyon, 1 (BSFC 3578); Tanks Peak, Dinosaur National Monument, 1 (BSFC 12803); Montezuma Co.: 1 mi W Mancos, 1 (KU 75971); Morfield Canyon, Mesa Verde National Park, 1 (KU 75973); Morefield [=Morfield] Canyon, ca. 3 mi S Morefield [=Morfield] Village, Mesa Verde National Park, 8 (BSFC 14216--14218, 14221, 14222, 14229, 14230, 14232); Prater Canyon, 0.25 mi N Middle Well, Mesa Verde National Park, 1 (KU 69239); Prater Canyon, Upper Well, Mesa Verde National Park, 1 (KU 69238); Montrose Co.: 28 mi SW Delta, head Monitor Creek, 2 (USNM 533010, 533011); 29 mi SW Delta, Smokehouse Campground, 10 (USNM 498400, 498401, 498403, 498404, 498406--498409, 533007, 533009); 13

mi N, 7 mi E Norwood, 1 (KU 120918); T48N, R14W, SW1/4 Sec. 11, 1 (KU 120921); Ouray Co.: Billy Creek Wildlife Area, 3 (BSFC 5876, 5877, 5880); Park Co.: Duck Creek, 8 mi NNW Grant, 2 (BSFC 33, 34); Wilkerson Pass, 9 mi NW Lake George, 1 (TCWC 3917); Rio Blanco Co.: no specific locality*, 1 (BSFC 3011); Avery Lake Meadow, 8 (BSFC 3012, 3017--3023); Cb Oil Shale Lease Tr*, 1 (BSFC 2207); Little Hill Res. Station, 2 mi S, 15 mi W Meeker, 1 (BSFC 2208); 1 mi S, 2 mi W Meeker, 1 (BSFC 2179); Oldland Ranch Spring, 7 mi N, 14 mi W Rio Blanco, 2 (BSFC 5353, 5360); 9.5 mi SW Pagoda Peak, 2 (KU 19933, 19934); PL Ranch, Willow Creek Spring, 7 mi N, 16 mi W Rio Blanco, 3 (BSFC 5359, 5361, 5362); 7 mi N Rio Blanco, 1 (BSFC 3456); 7 mi N, 14 mi W Rio Blanco, 4 (BSFC 5354--5356, 5358); Thornburgh Site, 1 (BSFC 3167); Rio Grande Co.: 8 mi S Monte Vista, 2 (UIMNH 33181, 33182); 8 mi S, 11 mi W Monte Vista, Comstack Campground, San Juan Mts., 1 (UIMNH 33180); Rio Grande Wildlife Area, 2 (BSFC 5878, 5879); Windy Mtn., San Juan Mts., 1 (UIMNH 33187); Routt Co.: no specific locality*, 2 (PSM 12027, 12029); 1 mi N Oak Creek (town), 1 (BSFC 5413); 1 mi S Slater [=Moffat Co.], 6 (BSFC 4609--4614); 3 mi N Steamboat Lake, 2 (BSFC 5415, 5416); 3 mi N Steamboat Lake, Deep Creek, 1 (BSFC 5417); 3 mi S Steamboat Springs, 3 (BSFC 5419, 5422, 5423); 3 mi SSW Steamboat Springs, T6N, R84W, Sec. 28, 1 (PSM 12028); 7.9 mi SE Steamboat Springs, 1 (FHSU 17330); Saguache Co.: 5 mi NE Cochetopa Dome, 1 (UMNH 24265); Cochetopa Pass, 2.5 mi E Summit, 2 (MVZ 60381, 60382); Cochetopa Pass, 33 mi W Saguache, 2 (KU 18206, 18207); Funk's Meadow, Cochetopa Park and W. Pass Creek, 1 (UMNH 23928); Gold Basin Creek, 10 mi SSE Gunnison, 1 (BSFC 25);

Monshower Meadows, 27 mi NW Saguache, 3 mi E Cochetopa Pass, 1 (USNM 48182); 3 mi N, 16 mi W Saguache, 1 (KU 41565); Samora Creek, 0.5 mi from Pass Creek, 2 (UMNH 23930, 23931); 0.25 mi from Summit, Lujan Creek, 1 (UMNH 23929); San Juan Co.: Silverton, 3 (USNM 56818--56820); Summit Co.: 0.5 mi S, 0.2 mi W Loveland Pass, 1 (BYU 9706); 0.75 mi S, 0.75 mi E Loveland Pass, 2 (BYU 9709, 9712); Teller Co.: Florissant, 13 mi N Trail Creek, 1 (BSFC 1336); Glen Cove, Pikes Peak, 1 (UMMZ 56308); Trail Creek, 10 mi N Florissant, 2 (BSFC 29, 31).

IDAHO (39).--Adams Co.: 1 mi N Bear Creek Ranger Station, SW slope Smith Mt., 1 (KU 45421); Summit of Smith Mt., 1 (KU 45423); 0.5 mi E Black Lake, 1 (KU 45427); Blaine Co.: Alturas Lake, 1 (MVZ 72109); Alturas Lake, 7000 ft., 1 (TCWC 24607); Perkins Lake, Sawtooth National Forest, 1 (KU 27348); Sawtooth City, 2 (USNM 74999, 75005); Boise Co.: Boise National Forest, Bald Mt. River, 10 mi S Idaho City, 1 (USNM 241793); Bonner Co.: Gold Peak Road, 1 (PSM 3816); Bonneville Co.: 10 mi SE Irwin, 1 (USNM 177269); 8 mi SE Palisades, 1 (TCWC 24604); Caribou Co.: 4.25 mi N, 1.5 mi W Hooper Springs, 1 (FHSU 21948); Custer Co.: Pahsimeroi Mts., 1 (USNM 31942); Fremont Co.: 7 mi W West Yellowstone, 3 (KU 33753, 33754, 33756); 4 mi N, 17 mi E Ashton, 16 (MVZ 88911, 88913, 88917, 88918, 88920, 88921, 88924--88929, 88931--88933, 88935); Idaho Co.: Devils Mts., 1 (USNM 74632); Teton Co.: 3 mi SW Victor, 1 (MVZ 72104); Valley Co.: Salmon River Mt., 1 (USNM 23524); Landmark R.S., Paytett National Forest, 10 mi E Warm Lake, 1 (USNM 265158); Washington Co.: 1 mi NE Heath, SW slope Cuddy Mt., 3 (KU 45429, 45430, 48121).

MEXICO (6).--Chihuahua, Sierra Madre, near Guadalupe y Calvo, 4 (USNM 95322--95324, 95326); Durango, El Salto, 2 (USNM 94539, 94540).

MONTANA (82).--Beaverhead Co.: Birch Creek, 18 mi NW Dillon, 11 (MVZ 106882--106892); Wise River, 2 (UMNH 26566, 26567); Carbon Co.: Beartooth Mountains, 1 (USNM 66710); 8 mi S, 22 mi E Bridger, 1 (KU 136308); Pryor Mountains, 4 (KU 136309, USNM 66489, 66490, 66501); 2 mi E Shriver, 1 (MVZ 106896); Cascade Co.: Neihart, Little Belt Mts., 1 (USNM 233464); Chouteau Co.: 0.5 mi N Arrow Creek Divide, Highwood Mts., 1 (KU 83728); Eagle Creek, 25 mi ESE Big Sandy, 2 (UMMZ 87334, 87335); Highwood Mountains, 6 (USNM 170012, 170013, 170015, 170021, 170023, 170026); Fergus Co.: Big Snowy Mountains, 2 (USNM 67563, 67564); Crystal Lake, Big Snowy Mtn., 1 (UMMZ 87324); 5 mi NW Hilger, Moccasin Mts., 1 (USNM 233460); Kendall, North Moccasin Mts., 1 (KU 86103); 7 mi N, 9 mi E Lewistown, 2 (KU 86104, 86105); Rock Creek, Big Snowy Mtn., 1 (UMMZ 87330); Rocky Creek, Big Snowy Mtn., 1 (UMMZ 87328); Flathead Co.: Gunsight Lake, Glacier National Park, 1 (USNM 244548); Nyack, 2 (UMMZ 75282, 75283); 2 mi S, 1 mi W Summit, 1 (KU 33758); Upper Stillwater Lake, 1 (USNM 72803); Gallatin Co.: Baker Hole, near West Yellowstone, 2 (CMNH 44786, 44787); Baker Hole, on Madison River, near West Yellowstone, 2 (CMNH 44794, 44801); Beaver Creek Camp, Madison River, 1 (BYU 9713); Gallatin National Forest, near West Yellowstone, 1 (CMNH 44789); West Fork, West Gallatin River, 3 (USNM 226675, 226677, 226679); Glacier Co.: 1.5 mi S, 2.5 mi W Babb, 1 (KU 33757); Crossley [=Cosley] Lake, Glacier National Park, 1 (USNM 246773);

Many Glacier, Glacier National Park, 2 (MVZ 88580, 88581); St. Mary's, Glacier Park, 1 (UMMZ 57957); St. Mary Lake Campground, Glacier National Park, 1 (MVZ 126573); Sherburne Lake, Glacier Park, 3 (UMMZ 57960, 57962, 57963); Hill Co.: head Eagle Creek, Bearpaw Mts., 2 (UMMZ 87337, 87339); Judith Basin Co.: 3 mi W Geyser, 1 (KU 42616); 20 mi SW Stanford, Dry Wolf Creek, Little Belt Mts., 1 (USNM 233463); Lake Co.: Saint Mary's Lake, 4 (USNM 72238, 72239, 72246, 72489); Madison Co.: 12 mi SW Alder Hinch Creek, Ruby Mts., 3 (USNM 226673, 226678, 226680); Ward Peak, Madison National Forest, Washington Creek, 1 (USNM 226674); Meagher Co.: 4 mi S Fort Logan, Camas Creek, Big Belt Mts., 3 (USNM 232921, 232924, 232925); Park Co.: 2 mi NE Cooke, 2 (MVZ 106899, 106898); Phillips Co.: Zortman, 1 (USNM 169621); Ravalli Co.: 3 mi E Hamilton, 1 (MVZ 99975); 8 mi NE Stevensville, 1 (USNM 168054).

NEVADA (11).--Elko Co.: 22 mi N Deeth, Mary's River, 1 (MVZ 67764); 1.25 mi S, 5 mi W Haystack Ranch, 1 (FHSU 20618); South Fork Humboldt River, T31N, R36E, Sec. 13, 1 (LACM 74491); South Fork Long Creek, Ruby Mts., 2 (KU 45407, 45408); 6 mi S, 7.5 mi W North Fork, Independence Mts., 1 (KU 139103); Three Lakes, Ruby Mts., 1 (KU 45411); Eureka Co.: Evans, 1 (MVZ 70521); Humboldt Co.: Summer Camp, Mahogany Creek, 1 (KU 133039); Ormsby Co.: Marlette Lake, 1 (MVZ 67039); 0.5 mi S Marlette Lake, 1 (MVZ 67041).

NEW MEXICO (145).--Bernalillo Co.: 1.2 mi E, 0.2 mi S Sandia Crest Benchmark, Sandia Mts., 1 (MSB 21635); Catron Co.: 19 mi ENE Glenwood, 1 (UIMNH 55932); 10 mi E Mogollon, 2 (FHSU 2925, 8113); 12 mi E Mogollon, 4 (MSB 41359, 41360, 41367, 41369); West Fork Gila

River, Mogollon Mts., 1 (FMNH 48101); Willow Creek, 1 (KU 1061);
 Willow Creek, Mogollon Mts., 1 (USNM 148327); near head Willow
 Creek, Mogollon Mts., 1 (USNM 158314); between Willow Creek &
 Mogollon, Mogollon Mts., 2 (MSB 6958, 6960); Cibola Co.: 4 mi N,
 17.5 mi E Grants, T12N, R7W, NW1/4 Sec. 5, 3 (MSB 54673--54675); 6
 mi N, 14 mi E Grants, T12N, R7W, 4 (MSB 49714--49717); Colfax Co.: 1
 mi S, 2 mi E Eagle Nest, 2 (KU 41568, 41569); 1 mi S, 3 mi E Eagle
 Nest, 4 (FHSU 166, 6635--6637); Philmont Scout Ranch, 17 mi NW
 Cimarron, 1 (USNM 554244); Rio Arriba Co.: Arroyo Yeso, N Ghost
 Ranch Headquarters, Carson National Forest, 1 (MSB 20567); 9 mi E, 4
 mi N Canjilou, 2 (MSB 6807, 6808); 6 mi E, 1 mi N Cuba, San Gregorio
 Lake, Jemez Mts., T21N, R1E, 24 (MSB 15683--15686, 15689, 15849,
 15850, 15912, 15915, 15999, 16000, 16629--16631, 16648, 16649,
 16768--16772, 16785--16787); 6 mi E Truchas, 2 (MVZ 116469, 116470);
 north slope Truchas Peak, 1 (MVZ 116472); Upper Canjilon Lakes, 1
 (MSB 32606); San Juan Co.: North end Chuska Mts., 1 (USNM 158740);
 Washington Pass, Chuska Mts., 1 (MSB 6896); San Miguel Co.: Gallinas
 Creek, near Ranger Station, 1 (MSB 483); Harvey's Ranch, 20 mi NW
 Las Vegas, Las Vegas Mts., 1 (KU 1754); Sandoval Co.: Fenton Lake,
 10 mi N Jemez Springs, 14 (MSB 1941, 1942, 1944, 1945, 1948--1950,
 1952, 1955, 2359, 2360, 2403, 2404, 2407); Jemez Creek, 6 mi NW
 Bland, 5 (MVZ 116333--116337); 3 mi N, 8.5 mi E Jemez Springs, Los
 Conchas, 1 (MSB 37449); 3 mi N, 8.5 mi E Jemez Springs, Los Conchas
 Campground, 2 (MSB 37452, 37457); 3 mi N, 9.5 mi E Jemez Springs, 1
 (MSB 43681); 3 mi N, 10.5 mi E Jemez Springs, 4 (MSB 37421, 43703--
 43705); 6 mi N Jemez Springs, 3 (MSB 42328--42330); 7.5 mi N, 6 mi E

Jemez Springs, 2 (MSB 53599, 53601); 12 mi N, 1 mi E Jemez Springs, 1 (MSB 42312); 12.5 mi N Jemez Springs, Fenton Lake, 1 (MSB 41053); 15 mi N, 2 mi E Jemez Springs, Cibola Creek, 1 (MSB 53715); 16 mi N Jemez Springs, 10 (MSB 41539, 41541--41543, 41568, 41574, 41576, 41629, 41630, 41697); 15 mi S, 5 mi W Los Alamos, Jemez Mts., 1 (MSB 22780); Seven Springs Fish Hatchery, 0.75 mi N, 1.5 mi E Jemez, 1 (USNM 399960); Santa Fe Co.: Hyde Park, 5 mi NE Santa Fe, 1 (UIMNH 7217); 16 mi NE Pojoaque, 2 (MVZ 116473, 116474); Rito Pacheco, Pacheco Canyon, 13 mi NE Santa Fe, 1 (UIMNH 7215); Socorro Co.: Copper Canyon, Magdalena Mts., 1 (USNM 160735); Mill Canyon, Magdalena Mts., 2 (MSB 9678, 9679); Taos Co.: 3.5 mi N, 3 mi E Arroyo Hondo, D.H. Lawrence Ranch, 2 (MSB 22336, 22368); 4 mi N, 11 mi E Arroyo Hondo, 4 (MSB 41250, 41323, 41324, 41326); Taos, 1 (USNM 133407); 1 mi N, 2 mi E Tres Ritos, 1 (FHSU 115); 1 mi N, 3 mi E Tres Ritos, 1 (FHSU 8119); 2 mi NE Tres Ritos, Rio La Junta, 2 (MSB 4966, 4969); 4 mi N Tres Ritos, 3 (MSB 966--968); T23N, R3E [=Rio Arriba Co.], 3 (MSB 970--972); Upper LaJunta Campground, near Tres Ritos, 1 (FHSU 14891); Torrance Co.: Fourth of July Campground, Manzano Mts., 1 (MSB 35351); Red Canyon Camp, 5 mi W Manzano, Manzano Mts., 2 (MSB 16541, 16542); Red Canyon, 0.5 mi S, 5 mi W Manzano, Manzano Mts., 2 (MSB 7898, 7899); 6 mi W Tajique on Forest Rd 55, Fourth of July Campground, 1 (MSB 35359); Valencia Co.: Upper Coalmine Canyon, Mt. Taylor [=Cibola Co.], 1 (MSB 10402); 2 mi NE La Mosca Peak [=Torrance Co.], 3 (MSB 10749, 10761, 10773).

NORTHWEST TERRITORIES (6).--Aklavik, 68°12'N, 135°W, 1 (CNM 24362); Mackenzie, Fort Simpson, 3 (USNM 133746, 133752, 140193);

Mackenzie, Mackenzie River; Nahanni River Mts., 1 (USNM 129739); Macmillan Pass, Canol Road, mile 286, 63°20'N, 129°40'W, 1 (CNM 18079).

OREGON (44).--Baker Co.: Anthony, 3 (MVZ 3700, 3705, USNM 154239); Bourne, 1 (USNM 208283); 0.3 mi SE Rock Creek Butte, Elkhorn Mts., 1 (CRCM 82-16); Grant Co.: 8 mi S, 5 mi W Long Creek, T11S, R31E, Sec. 19, 5 (OSUFW 7335, 7341, 7342, 7345, 7346); Malheur River, N.F.K., 1 (MVZ 83547); Strawberry Butte, 1 (USNM 79384); Harney Co.: Fish Lake Region, Steens Mt., 9 (OSUFW 7308--7310, 7312--7316, 7318); 5 mi S, 11 mi E Frenchglen, T32S, R32 3/4E, Sec. 34, 1 (KU 145770); Little Blitzen Gorge, Steens Mt., T33S, R33E, Sec. 10, 2 (OSUFW 4746, 4812); 18 mi N, 4 mi W Riley, T20S, R26E, SE1/4 Sec. 26, 3 (KU 145761, 145762, OSUFW X2512); Steens Mt., 1 (OSUFW 7570); Steens Mt., T32S, R33E, Sec. 32, 7 (OSUFW 4747, 4749, 4762, 4810, 4815, 4817, 4818); Steens Mt., T33S, R33E, Sec. 2, 1 (MVZ 119708); Wallowa Co.: 15 mi S, 2 mi E Lostine, T3S, R43E, Sec. 25, 1 (OSUFW 4798); 19 mi S, 4 mi E Lostine, T4S, R44E, Sec. 17, 1 (OSUFW 4799); Wallowa Lake, 3 (MVZ 81067, 81069, USNM 90724); South of Wallowa Lake, Wallowa Mts., 1 (USNM 90756); Wallowa Mts., 1 (USNM 96003); Wheeler Co.: 7 mi S, 11 mi W Mitchell, 1 (MVZ 83542).

SASKATCHEWAN (4).--Cypress Hills, 4 (CNM 19463, 19468, 19473, 19474).

UTAH (261).--Beaver Co.: Meadow, west and north Big Flat Guard Station, Tushar Mts., 5 (UMNH 16455, 16456, 16458--16460); Delano Ranger Station, Tushar Mts., 4 (UMNH 16463--16466); East shore Kents Lake, Tushar Mts., 1 (UMNH 16470); Merchant Creek at confluence of

Crazy Creek, Tushar Mts., 1 (UMNH 16461); Puffer Lake, Beaver Mts.,
 1 (USNM 158514); Stream side, South of Puffer Lake, Tushar Mts., 2
 (UMNH 16467, 16473); Cache Co.: Cache National Forest, 1 (SUVM
 3452); Daggett Co.: Birch Creek, 4 mi S Utah-Wyoming line, 2 (UMNH
 20364, 20365); Junction Deep Creek and Carter Creek, 1 (UMNH 6071);
 15.8 km S Manila, 1 (SUVM 5442); Sheep Creek, Palisade Campground, 2
 (UMNH 18809, 18810); East side Spirit Lake, 1 (UMNH 18812); 7.3 km
 NE Spirit Lake, 2 (SUVM 7130, 7134); Duchesne Co.: Anthro Mt., 1
 (BYU 10551); Butterfly Lake, 2 (BYU 9696, 9697); Highline Trail,
 Scudder Lake, 1 (BYU DS4-2#5); Horse Ridge, 1 (BYU 10588); Uinta
 Park, 1 (BYU 3339); Emery Co.: Huntingdon Canyon at Corral Creek, 1
 (UMNH 13206); Lake Creek, 11 mi E Mt. Pleasant, 2 (CMNH 14206,
 14211); Olsen Ranch, Lower Joe's Valley, 3 (UMNH 13223--13225); near
 Seely [=Seeley] Mt., 21 mi NE Ephram [=Ephraim, San Pete Co.], 1
 (CMNH 14202); Upper Joe's Ranger Station, 3 (UMNH 13214--13216);
Garfield Co.: E. Fork Boulder Creek, 10 mi N Boulder, 1 (UMNH 9726);
 Daves Hollow, NW visitors center, Bryce Canyon National Park, 2
 (BSFC 13828, 13829); East Creek Wells, Bryce Canyon National Park, 4
 (BSFC 13291, 13744--13746); Griffin Spring, 8 mi NE Widsoe [=
 Widtsoe], 1 (UMNH 9728); King's Pasture, 10 mi N Boulder, 1 (UMNH
 9729); Pine Lake, 6 mi SE Widsoe [=Widtsoe], 1 (UMNH 9109); Round
 Willow Bottom Res., 10 mi N, 13 mi W Escalante, 1 (UMNH 9054); Steep
 Creek, 15 mi N Boulder, 1 (UMNH 9091); Upper Valley Ranger Station,
 15 mi SW Escalante, 1 (UMNH 9049); Grand Co.: Beaver Basin, NW Mans
 Peak, La Sal Mts., 4 (UMNH 12474, 12475, 12477, 12480); Beaver
 Creek, 0.5 mi E Beaver Basin, La Sal Mts., 1 (UMNH 12452); Beaver

Creek, 1.5 mi E La Sal Peak, La Sal Mts., 10 (UMNH 12448, 12450, 12456, 12457, 12459, 12460, 18949--18952); Beaver Creek, 2 mi NE Mt. Waas, La Sal Mts., 4 (UMNH 18932--18935); LaSal Mts., 1 (USNM 149971); 1.5 mi E La Sal Peak, La Sal Mts., 1 (UMNH 14289); 2.5 mi NE La Sal Peak, La Sal Mts., 1 (UMNH 14290); Mill Creek, La Sal Mts., 1 (BYU 11590); Pioche Spring, W. Fork Willow Creek, 17 mi N Thompsons, 1 (CMNH 13289); Warner Ranger Station, La Sal Mts., 3 (BYU 257, UMNH 6680, 6681); Iron Co.: Brian Head, 1 (BYU 3334); Parowan Mts., Brian Head, 2 (USNM 157954, 157955); Kane Co.: Yovimpa Pass Meadow, Bryce Canyon National Park, 4 (BSFC 13248--13251); Millard Co.: Robin's Valley, Pavant Range, 5 (UMNH 16448--16452); Morgan Co.: Salt Lake, near summit of Wasatch Mts., 1 (USNM 186694); Piute Co.: head of Two Mile Canyon, Tushar Mts., 1 (UMNH 16453); Salt Lake Co.: Brigham Fork, Emmigration Canyon, 10 mi E Salt Lake City, 1 (UMNH 24020); Brighton, 3 (UMNH 5139--5141); 1 mi W Brighton, 1 (UMNH 13603); 1 mi E mouth Little Cottonwood Canyon, 1 (UMNH 9432); 1 mi W Boy Scout Camp, Millcreek Canyon, 1 (UMNH 21707); Mill Creek Guard Station, Mill Creek Canyon, 1 (UMNH 18362); 2 mi E Mill Creek Guard Station, Mill Creek Canyon, 1 (UMNH 14168); Spruces, Camp 67, Big Cottonwood Canyon, 1 (UMNH 10699); Wilson Fork, Mill Creek Canyon, 1 (UMNH 9377); San Juan Co.: Cooley Pass, 8 mi W Monticello, 3 (CMNH 15529, 15531, 15533); Horse Creek, La Sal Mts., 2 (BYU 11556, 11557); Kigalia Ranger Station, Elk Ridge, 1 (BYU 3378); La Sal Mts., 4 (BYU 11686--11689); North Creek, 6 mi W Monticello, 2 (UMNH 12466, 12471); 1 mi S Twin Peaks, Abajo Mts., 1 (UMNH 12472); Sanpete Co.: Ephraim Co., 9 mi E Ephraim, 1 (UMNH

9330); Ferron Reservoir, 28 mi W Ferron, 3 (UMNH 13296, 13298, 13299); Gooseberry Creek, 1 mi S Lower Gooseberry Reservoir, 5 (UMNH 13198--13202); Indian Spring Creek, above Ferron Reservoir, 2 (UMNH 13210, 13211); 0.5 mi E Mammoth Ranger Station, 5 (UMNH 13234--13238); Manti, 2 (USNM 186697, 186698); Mt. Baldy Ranger Station, 1 (UMNH 9342); Seeley Creek Ranger Station, 1 (UMNH 13232); 0.25 mi SW Seeley Creek Ranger Station, 1 (UMNH 13204); Straight Fork Pleasant Creek, 9 (UMNH 13239, 13241, 13242, 13244--13247, 13249, 13251); Tom's Hole, 0.25 mi E Seeley Creek, 1 (UMNH 13227); Sevier Co.: Convulsion Canyon, 41 mi E Salina, 1 (BYU 6629); Fish Lake Plateau, 2 (USNM 157956, 157957); Hunt's Lake, 2 mi W Monroe Peak, 1 (UMNH 16468); 1 mi NW Mt. Marvine, 1 (UMNH 9433); Rock Spring Creek, Seven Mile Canyon, 1.5 mi N Johnson's Reservoir, 1 (UMNH 9898); 7 [=Seven] mile Creek, 20 mi SE Salina, 4 (CMNH 15513--15515, 15517); Seven Mile Creek Trib., 1.5 mi N Johnson's Reservoir, 3 (UMNH 9379, 9383, 9384); Summit Co.: Bald Mt., 2 (UMNH 21976, 21977); Jct. Bear River and East Fork, 2 (CMNH 16796, 16797); 2 mi S, Jct. Bear River and Hayden Fork, 1 (CMNH 16801); 4.8 km ENE Bridger Lake, 2 (SUVN 7131, 7132); Burnt Fork, Round Lake, Uintah Mts., 2 (BYU 11788, 11789); China Meadows Campground, 2 (SUVN 5438, 5439); 0.5 mi S Dahlgreen Camp, 2 (UMNH 21978, 21979); 8.9 km ESE Kamas, 2 (SUVN 7137, 7138); Little East Fork, Blacks Fork, 1 (BYU 11825); Little East Fork, East Fork, Blacks Fork, Uintah Mts., 2 (BYU 11781, 11783); 2.25 km S, 2.1 km W Mirror Lake, 1 (SUVN 6501); 13.2 km N, 3.2 km E Mirror Lake, 5 (SUVN 6503--6505, 6508, 6509); 6.9 km N, 13.9 km E Oakley, 3 (SUVN 6499, 6510, 6596); East Fork, South Fork Uintah Mts., 1 (BYU 11780);

Tooele Co.: Head of Mack Canyon, Stansbury Mts., 3 (UMNH 26368, 26538, 26539); Right Fork, Middle Canyon, Oquirrh Mts., 1 (UMNH 26545); Upper Fork Ophir Creek, Oquirrh Mts., 1 (UMNH 26540); South Willow Creek, Stansbury Mts., 1 (UMNH 26577); North slope South Willow Creek, 1 (UMNH 26544); South Willow Creek Canyon, Stansbury Mts., 4 (UMNH 26370, 26530, 26536, 26537); 7.5 mi S, 3 mi E Vernon, 1 (BSFC 2735); 10 mi SE Vernon, 1 (BSFC 2733); Right Fork, Vernon Creek Canyon, Sheep Rock Mts., 1 (UMNH 26542); Uintah Co.: N Fork Ashley Creek, Uintah Mts., 2 (UMNH 26558, 26559); N Fork Ashley Creek, 1.2 mi E Hacking Lake, 2 (UMNH 26551, 26552); Blue Mt., Point of Pines Area, 1 (BYU 10822); Head Colton Draw, Colton Ranger Station, 2 (UMNH 26548, 26549); Hatch Cabin, 3 mi NE PR Springs, 1 (UMNH 15336); Kaler Camp, 20 mi NW Vernal, 1 (UMNH 26560); East base Leidy Peak, Uintah Mts., 1 (UMNH 26553); Paradise Park, 15 mi N, 21 mi W Vernal, 3 (KU 38030--38032); Paradise Park, Uinta Mts., 3 (CMNH 13284, UMNH 6069, 6070); 17.1 km S Robertson, 4 (SUVN 6631, 6632, 6634, 6636); Trout Creek, Trout Creek Ranger Station, 2 (UMNH 26556, 26557); Head Trout Creek, 250 yards W Trout Creek Ranger Station, 1 (UMNH 26546); Trout Creek Park, 200 feet E Trout Creek Ranger Station, 1 (UMNH 26547); 28.3 km NNE Vernal, 2 (SUVN 5436, 5437); Utah Co.: 1 mi E Payson Lake, Nebo Mts., 1 (UMNH 7203); Summit Campground, 13 (BYU 10164, 10166--10169, 10175, 10176, 10179, 10180, 10993, 10994, 10996, 10997); Wasatch Co.: Bald Mt. [=Summit Co.], 25 mi NE Kamas, 1 (CMNH 13287); Washington Co.: 1.5 mi E Pine Valley, 8 (UMNH 21111, 21114--21119, USNM 375531); Inlet, Pine Valley Lake, 1 mi E Pine Valley, 1 (USNM 375528); Pine Valley Mts., 4 mi E Pine

Valley, 2 (USNM 166720, 166721); Pine Valley Mts., 5 mi E Pine Valley, 2 (USNM 166716, 166718); Wayne Co.: Aquarius Guard Station, 10 mi S Bicknell, 1 (UMNH 9724); 1 mi NE Thousand Lake Mts., 1 (UMNH 13733).

WASHINGTON (49).--Asotin Co.: 1 mi S, 11 mi W Anatone, 1 (FHSU 3840); Chelan Co.: head Lake Chelan, 3 (USNM 42244, 42603, 42604); Cloudy Pass, 1 (CRCM 20); Wenatchee, 1 (USNM 91044); Garfield Co.: Spruce Spring, 1 (PSM 5798); King Co.: Lester, 1 (PSM 10634); Little Eagle Lake, 1 (PSM 10646); Kittitas Co.: Easton, 8 (USNM 41619, 41621--41625, 41628, 41629); 4.5 mi N, 6.5 mi W Easton, 1 (KU 57220); Swauk Cr., Dunning Ranch Trail, 1 (PSM 2094); Lincoln Co.: 12 mi N Odessa, 1 (PSM 4068); Okanogan Co.: Bauerman Ridge, west end at Tungsten Mine, 1 (USNM 235205); E. Cirque Slate Peak, 1 (CRCM 82-19); Pend Oreille Co.: Gypsy Meadow, 1 (CRCM 47-143); Pierce Co.: Chrystal [=Crystal] Mt., 1 (PSM 11934); Corral Pass, T18N, R11E, Sec. 30, 1 (PSM 26388); Stevens Co.: Loon Lake, 3 (PSM 2543, 2547, 2849); Spokane Indian Reservation, 2 (CRCM 86-119, 86-120); Spokane Indian Reservation, T27N, R37E, Sec. 13, 1 (CRCM 85-791); Spokane Indian Reservation, T27N, R39E, Sec. 5, 2 (CRCM 85-773, 85-775); Spokane Indian Reservation, T28N, R38E, Sec. 2, 1 (CRCM 86-60); Spokane Indian Reservation, T28N, R38E, Sec. 12, 2 (CRCM 86-178, 86-206); Spokane Indian Reservation, T29N, R39E, Sec. 26, 1 (CRCM 86-26); Spokane Indian Reservation, T29N, R39E, Sec. 35, 2 (CRCM 86-139, 86-150); Spokane Indian Reservation, T29N, R40E, Sec. 20, 2 (CRCM 86-77, 86-78); 4 mi N Wellpinit, 1 (CRCM 86-448); 5 mi NE Wellpinit, 3 (CRCM 86-458, 86-609, 86-693); 6 mi NNE Wellpinit, 4

(CRCM 86-351, 86-479, 86-585, 86-602); Moses Rd., 7.5 mi NE Wellpinit, 3 (CRCM 86-403, 86-466, 86-477); 8.5 mi NE Wellpinit, 1 (CRCM 86-354); Cottonwood Rd., 8.5 mi NE Wellpinit, 3 (CRCM 86-561, 86-675, 86-691); S. Cottonwood, 8.5 mi NE Wellpinit, 1 (CRCM 86-419); 9 mi NE Wellpinit, 1 (CRCM 86-676); Yakima Co.: Yakima Indian Reservation, Signal Peak, 1 (USNM 226855).

WYOMING (269).--Albany Co.: Bear Creek, 3 mi SW Eagle Peak, 2 (USNM 160235, 160236); Blair Campground, 1 (KU 91020); 2.25 mi ESE Browns Peak, 1 (KU 16735); 3 mi ESE Browns Peak, 3 (KU 16747, 16756, 16759); NE Burnett Homestead*, 2 (BYU 9719, 9720); Foxpark, 1 (UW 2262); 2 mi E Happy Jack Ski Area, 1 (UW 2245); 1 mi E Hidden Valley Picnic Ground, 1 (UW 2265); 4 mi SW Laramie, 1 (KU 91014); 6.1 mi SW Laramie, 1 (KU 91017); 6.5 mi S, 8.75 mi E Laramie, 3 (KU 27688, 27689, 27691); 9 mi S Laramie, 1 (UW 2428); 11.8 mi S Laramie on 287, Red Buttes Biological Research Facility, 1 (UW 2426); 13.5 km S Laramie on 287, Flag Ranch, 1 (UW 2427); 20 mi SW Laramie, 1 (UW 2247); 26 mi N, 4.5 mi E Laramie, 1 (KU 27686); 26.75 mi N, 4.5 mi E Laramie, 1 (KU 27685); 26.75 mi N, 6.5 mi E Laramie, 1 (KU 27684); 27 mi N, 5 mi E Laramie, 2 (KU 27682, 27683); 29 mi N, 8.75 mi E Laramie, 2 (KU 27679, 27680); 30 mi N, 10 mi E Laramie, 3 (KU 27427--27429); Lazenby Lake, 10 mi S Laramie, 1 (UW 2248); 0.2 mi N Libby Flats, 2 (BYU 9686, 9687); 700 ft east Libby Lake Outlet, 2 (BYU 9724, 9725); 0.1 mi S, 0.4 mi E Outlet Libby Lake, 2 (BYU 9689, 9691); Nelson Park, Medicine Bow Mts., 1 (KU 91018); 1 mi ESE Pole Mt., 3 (KU 16763, 16764, 16766); 2 mi SW Pole Mt., 2 (KU 16723, 16725); 3 mi S Pole Mt., 1 (KU 16732); 1.3 mi SE Pole Mt. Ranger

Station on Hidden Valley Road, 1 (UW 2249); Red Buttes Biological
 Research Station, 1 (UW 2430); S.H. Knight Science Camp, 1 (UW
 2235); East slope Sugar Loaf Peak, 2 (BYU 9692, 9693); University of
 Wyoming Science Camp, 1 (UW 2252); Vedauwoo picnic ground, 1 (KU
 91022); 0.2 mi E Vedouwod [=Vedauwoo] Camp, 2 (BYU 9721, 9722); near
 Wallis Campground, Pole Mtn. Div. N.F., 1 (KU 91021); Woods Post
 Office, 1 (USNM 186693); Big Horn Co.: Big Horn Mts., 1 (USNM
 56147); Medicine Wheel Ranch, 28 mi E Lovell, 8 (KU 32252, 32253,
 32258, 32260--32262, 32264, 32268); 1 mi N, 12 mi E Shell, 1 (KU
 20906); 4.5 mi S, 17.5 mi E Shell, 1 (KU 19915); head of Trapper's
 Creek, Big Horn Mts., 1 (USNM 168798); Carbon Co.: Bridger Pass, 18
 mi SW Rawlins, 2 (KU 19937, 19938); 8 mi N, 14 mi E Encampment, 1
 (KU 25219); 8 mi N, 14.5 mi E Encampment, 1 (KU 25231); 8 mi N, 16
 mi E Encampment, 3 (KU 25234--25236); 8 mi N, 21.5 mi E Encampment,
 2 (KU 25252, 25256); 9 mi N, 3 mi E Encampment, 1 (KU 25211); 10 mi
 N, 12 mi E Encampment, 1 (KU 25196); 10 mi N, 14 mi E Encampment, 3
 (KU 25202, 25203, 25207); Ferris Mts., 4 (USNM 160573, 160581,
 160583, 160585); 6 mi S, 14 mi E Saratoga, 1 (KU 25194); Loco Creek,
 20 mi W Saratoga, 1 (BSFC 1800); 5 mi N, 10.5 mi E Savery, 1 (KU
 25269); 6.5 mi N, 16 mi E Savery, 1 (KU 25265); 7 mi N, 17 mi E
 Savery, 1 (KU 25263); 8 mi N, 19.5 mi E Savery, 3 (KU 25253, 25254,
 25260); Shirley Mts., 5 (USNM 160241, 160242, 160244, 160246,
 160247); Sierra Madre Mts., S base Bridger Peak, 2 (USNM 176506,
 176508); 1 mi NW Silver Lake, 1 (KU 26687); Trowbridge Ranch, 5 mi
 SW Encampment, 3 (BSFC 14483, 14485, 14488); Converse Co.: 21 mi S,
 24.5 mi W Douglas, 1 (KU 32280); 21.5 mi S, 24.5 mi W Douglas, 4 (KU

32278, 32282, 32286, 32287); 22 mi S, 24.5 mi W Douglas, 2 (KU 32275, 32276); 22.5 mi S, 24.5 mi W Douglas, 1 (KU 32281); Northslope Laramie Peak, 4 (USNM 160226, 160229, 160230, 160232); Springhill, 12 mi N Laramie Peak, 6 (USNM 160215, 160220--160224); Fremont Co.: Bronx, 1 (UMMZ 106644); 17 mi S, 6.5 mi W Lander, 2 (KU 37296, 37297); Mocassin Lake, 4 mi N, 19 mi W Lander, 1 (KU 32272); Mosquito Park Ranger Station, 2.5 mi N, 17.5 mi W Lander, 1 (KU 32273); Rattlesnake Mts., 10 (USNM 160255, 160257--160259, 160261, 160263, 160266, 160267, 160269, 160270); 8 mi E Rougis, Green Mts., 1 (USNM 166869); Sweetwater, near Hwy 187 Fremont Co. border, 1 (UW 1699); Laramie Co.: 4.5 mi E Cheyenne on Herford Road, 2 (UW 2231, 2232); 1 mi N, 5 mi W Horse Creek Post Office, 1 (KU 14779); Lincoln Co.: 10 mi SE Afton, Salt River Mts., 3 (USNM 176970, 176971, 177272); Natrona Co.: 7 mi S Casper, Casper Mts., 2 (USNM 160249, 160250); 7 mi S, 1 mi W Casper, 1 (KU 27426); Park Co.: Beartooth Lake, 3 (USNM 66713--66715); Beartooth Lake, above timberline, 1 (USNM 66718); Beartooth Lake, Shoshone National Forest, 2 (TCWC 11390, 11391); Blacktail Deer Creek, Yellowstone National Park, 1 (MVZ 135367); Ishawooa Creek, 19 mi S, 19 mi W Cody, 2 (KU 39402, 39403); Mammoth Hot Springs, Snow Pass; Yellowstone National Park, 1 (USNM 120590); Mammoth Hot Springs, Yellowstone National Park, 6 (USNM 120591, 120594--120598); Needle Mt., 1 (USNM 169316); Needle Mt., tributary of Boulder Creek, 1 (USNM 169507); Pahaska, Grinnell Creek, 3 (USNM 169880, 170102, 170108); Pahaska, mouth of Grinnell Creek, 7 (USNM 169838, 169845, 169846, 169864, 169865, 169870, 169881); Pahaska Tepee, mouth Grinnell Creek, 4 (USNM 169849,

169853, 169854, 170101); Slough Creek, Yellowstone National Park, 2 (MVZ 135368, 135369); Valley, 7000 ft, 1 (USNM 169278); Valley, Shoshone Mts., 2 (USNM 169280, 169282); SW slope Whirlwind Peak, 1 (KU 21673); Yellowstone National Park, 2 (USNM 67383, UW 1698); Yellowstone Park, Willow Park, 1 (USNM 66723); Platte Co.: NE base Black Mt., Pat O'hare, 3 (USNM 169124, 169127, 169137); Sublette Co.: no specific locality*, 1 (KU 4065); North side Half Moon Lake, 1 (KU 14776); 37 mi SE Jackson, 1 (KU 1545); 2.25 mi NE Pinedale, 2 (KU 14777, 14778); 12 mi NE Pinedale, Surveyor Park, 1 (USNM 176725); 31 mi N Pinedale, 1 (KU 41557); Sweetwater Co.: Black Rock Creek, 2 mi W Pass, 1 (USNM 170554); Teton Co.: Beaver Dick Lake, Grand Teton National Park, 1 (UMMZ 68085); South Shore Emma Matilda Lake, Grand Teton Mts., 1 (UIMNH 7603); Jackson, Williams Slough, 1 (UMMZ 67553); Jackson Hole Wildlife Park, 11 (USNM 303665, 303667, 303669, 303672, 303673, 303675, 303677, 303681--303684); Jackson Hole Wildlife Park, headquarters, 4 (USNM 303655--303657, 303663); Jackson Hole Wildlife Park, 0.5 mi E Moran, 1 (UW 34); Jackson Hole Wildlife Park, 1 mi E Moran, 2 (KU 91011, 91012); 2.5 mi NE Moose Bar BC Ranch, 1 (KU 32940); 4 mi N Moose Timbered Island, 2 (KU 32938, 32939); Moran, 5 (KU 16697, USNM 170268, 170269, 303658, UW 6257); NW Moran, Willows, 1 (UIMNH 7604); 0.25 mi N, 2.5 mi E Moran, 3 (KU 16695, 16696, 16749); 1 mi N Moran, 1 (USNM 170221); 1 mi S, 3.75 mi E Moran, 6 (KU 16703, 16705--16707, 16711, 16712); 3 mi E Moran, 2 (KU 16751, UIMNH 56453); 3.75 mi E Moran, 1 (KU 16750); Old Faithful, Yellowstone Park, 3 (USNM 246757, 246758, 246806); Teton Mts., Moose Creek, 2 (USNM 170249, 170252); Teton Mts., south of

Moose Creek, 3 (USNM 170320--170322); Teton Pass, above Fish Creek, 5 (USNM 170330, 170332, 170335, 170346, 170347); Whetstone Creek, 3 (UMMZ 62025, 62026, USNM 249209); Yellowstone National Park, 1 (UMMZ 83653); Uinta Co.: Fort Bridger, 2 (KU 16716, 16717); 1 mi N Fort Bridger, 1 (KU 16714); 9 mi S, 2.5 mi E Robertson, 1 (KU 25186); 10 mi S, 1 mi W Robertson, 1 (KU 25188); 10.5 mi S, 2 mi E Robertson, 1 (KU 25190); Washakie Co.: 4 mi N, 9 mi E Ten Sleep, 1 (KU 19927); 5 mi N, 9 mi E Ten Sleep, 1 (KU 19925).

YUKON TERRITORY (54).--no locality given, 1 (CNM 29396); Bonnet Plume Lake, 64°20' N, 132° W, 4 (CNM 35252, 35254, 35257, 35259); 5 mi N, 1 mi W Carcross, 1 (CMNH 75972); 20 mi S Chapman Lake, 1 (CNM 33696); 20 mi S Chapman Lake, 64°34' N, 138°15' W, 1 (CNM 29397); 1.5 mi S, 3 mi E Dalton Post, 1 (KU 28524); Dempster Hwy, mile 51, 64°33' N, 138°15' W, 1 (CNM 44986); Dezadeash Lake, 2 (CNM 18184, ROM 16250); Dezadeash Lake, Kluane National Park, 3 (UBC 16531--16533); Haeckel Hill, 8 mi NW Whitehorse, 1 (CNM 31153); Haeckel Hill, 60°46' N, 135°17' W, 1 (CNM 37067); Klondike Keno, Keno Hill, 63°55' N, 135°18' W, 1 (CNM 35272); Kluane, 2 (UBC 16130, 16131); Kluane Range, 25 mi SE Destruction Bay, 1 (CNM 29402); Kluane Range, 25 mi SSE Destruction Bay, 1 (CNM 29403); Little Hyland River, 128 mi N Watson Lake, 2 (CNM 31154, 31156); Macmillan Pass, Canol Road, mile 282, 63°12' N, 130°05' W, 1 (CNM 18093); South Fork, Macmillan River, Canol Road, mile 249, 62°55' N, 130°30' W, 2 (CNM 18049, 18061); McIntyre Creek, 3 mi NW Whitehorse, 1 (KU 21074); Nisutlin River, Canol Road, mile 40, 4 (CNM 17904, 17906, 17924, 17940); North Toobally Lake, 60°20' N, 126°15' W, 1 (CNM 29401); Old Crow, 1

(CNM 29858); Old Crow Mtn. Range, 1 (ROM 74823); Rose [=Ross] River, Canol Road, mile 95, 7 (CNM 17797, 17826, 17833, 17846, 17849--17851); Rose [=Ross] River, Canol Road, mile 95, 61°10' N, 132°59' W, 1 (CNM 17845); Sheldon Lake, Canol Road, mile 222, 2 (CNM 18023, 18024); Slims River, Kluane National Park, 1 (UBC 16534); South Moose Creek, 25 mi N, 32 mi E Tuchitua jct. (Cantung), 61°10'30"N, 128°17'0" W, 8 (CMNH 75993--75997, 75999--76001).

Sorex monticolus parvidens Jackson

1921. Sorex obscurus parvidens Jackson, J. Mammal., 2:161.
 1955. Sorex vagrans parvidens Findley, Univ. Kansas Publ., Mus. Nat. Hist., 9:58, December.
 1977. S[orex]. m[onticolus]. parvidens Hennings and Hoffmann, Occas. Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.

Holotype.--USNM 56561, original number 242; adult male collected 3 October 1893 from "Thurman's Camp, Bluff Lake, . . . western side of San Bernardino Peak, San Bernardino Mountains, California" by J. E. McLellan. Type specimen examined and measured.

Distribution.--S. m. parvidens is restricted to the San Bernardino Mountains and San Gabriel Mountains in southern California (Fig. 17).

Subspecific comparisons and remarks.--Because only six shrews referable to S. m. parvidens were examined, this taxon was not included in the multivariate analysis. This taxon is included in the discussion of geographic variation within the subspecies of S. monticolus; comparisons were made among the cranial and mandibular characters by examination of the summary statistics for univariate characters.

Specimens examined.--CALIFORNIA (6).--Los Angeles Co.: 0.4 mi W Wrightwood, San Gabriel Mts., 1 (RHMC 6423); San Bernardino Co.: Fisherman's Camp on Deep Creek near Lake Arrowhead, 1 (RHMC 4766); San Bernardino Peak, 1 (USNM 56561); San Bernardino Peak, Bluff Lake, 2 (USNM 56558, 56559); Summit, 1 (USNM 55550).

Sorex monticolus prevostensis Osgood

1901. Sorex longicauda prevostensis Osgood, N. Amer. Fauna, 21:35, September.
1905. Sorex obscurus prevostensis Elliot, Field Columb. Mus., Publ. 105, Zool. Ser., 6:450.
1955. Sorex vagrans prevostensis Findley, Univ. Kansas Publ., Mus. Nat. Hist., 9:41, December.
1977. S[orex]. m[onticolus]. prevostensis Hennings and Hoffmann, Occas. Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.

Holotype.--USNM 100618, original number 1089; adult male collected 3 July 1900 from "Prevost Island, Queen Charlotte Islands, British Columbia" by W. H. Osgood and E. Heller. Type specimen examined and measured.

Distribution.--S. m. prevostensis is restricted to Kunghit Island at the southern tip of the Queen Charlotte Islands, British Columbia (Fig. 14).

Subspecific comparisons and remarks.--S. m. prevostensis seems to be allied more closely with S. m. longicaudus, S. m. insularis, and S. m. malitiosus than with S. m. elassodon, the subspecies that occurs on the remainder of the Queen Charlotte Islands.

Specimens examined.--BRITISH COLUMBIA (25).--Prevost Island [=Kunghit Island], Queen Charlotte Islands, 8 (USNM 100611, 100612, 100615, 100616, 100618, 100619, 107253, 107254); Kunghit, Queen Charlotte Islands, 1 (CNM 27253); Kunghit Island, Queen Charlotte Islands, 12 (UBC 2109, 2110, 2112--2114, 2117, 2118, 2120--2122, 2130, 2133); Rose Harbour, Kunghit Island, Queen Charlotte Islands, 4 (CNM 30802, 30803, UBC 7931, 7932).

Sorex monticolus setosus Elliot

1899. Sorex setosus Elliot, Field Columb. Mus., Zool. Ser., 1:274,
May.
1918. Sorex obscurus setosus Jackson, Proc. Biol. Soc. Washington,
31:127, November.
1938. Sorex obscurus mixtus Hall, American Nat., 72:462. (Type: MVZ
70376).
1955. Sorex vagrans setosus Findley, Univ. Kansas Publ., Mus. Nat.
Hist., 9:36, December.
1955. Sorex vagrans mixtus Findley, Univ. Kansas Publ., Mus. Nat.
Hist., 9:38, December.
1977. S[orex]. m[onticolus]. setosus Hennings and Hoffmann, Occas.
Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.
1977. S[orex]. m[onticolus]. mixtus Hennings and Hoffmann, Occas.
Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.

Holotype.--FMNH 6213/238; adult male collected 18 August 1898
from "Happy Lake, Olympic Mountains, Clallam County, Washington" by
D. G. Elliot. Type specimen not examined.

Distribution.--S. m. setosus occurs from approximately Rivers
Inlet, British Columbia, east to the Fraser River and southward to
Washington where it occurs primarily on the west side of the Cascade
Range. S. m. setosus also occurs in a narrow band near the Columbia
River in northwestern Oregon (Fig. 15).

Specimens examined.--BRITISH COLUMBIA (227).--Alpha Lake, Mons, 1 (UBC 6407); Alta Lake, 19 (RBCM 1537, 1539, 1540, 1543, 1549, 2403, UBC 42, 493, 498, 649, 1150, 1845, 1846, 1849, 1852, 1853, 6454, 7434, 7435); Alta Lake, Mons, 27 (UBC 6368, 6369, 6371, 6373--6379, 6381, 6383--6386, 6388--6399); Sprout Mountain, Alta Lake, 1 (UBC 6406); Bowen Island, 2 (UBC 50, 64); Cortes Island, 2 (RBCM 9621, 12949); Cultus Lake, 1 (UBC 540); 1.5 km upstream from mouth Foley Creek, 1 (RBCM 11022); Black Tusk Meadow, Garibaldi Provincial Park, 1 (UBC 3603); Parnassis Cr., Black Tusk Meadow, Garibaldi Provincial Park, 1 (RBCM 1148); Outlet Green Lake, North Garibaldi Provincial Park, 1 (BYU 9702); Goose Island, 21 (RBCM 3962--3965, 5413--5418, 5421--5423, UBC 2705--2711, 2714); UBC Research Forest, Haney, 6 (UBC 10103, 10107, 10108, 10135, 10143, 10765); Hecate Island, 1 (RBCM 4237); Horseshoe Lake, 3 (CNM 14167, 14183, 14190); 8 mi N Stillwater, Horseshoe Lake, 7 (CNM 14367, 14380, 14516--14518, 14529, 14546); Black Mountain, Howe Sound, 1 (UBC 52); Gibson's Landing, Howe Sound, 9 (USNM 90250, 90251, 90253, 90255--90257, 90261, 92806, 92808); Huntingdon, 4 (UBC 56, 416, 417, 767); Lesser Garibaldi Lake, 1 (RBCM 6404); Lund, 1 (USNM 92812); Lund, Malaspina Inlet, 5 (USNM 89475, 89479, 89482, 90243, 90249); 3 mi E Lund, Okeover Arm Park, 5 (TCWC 45651, 45654, 45658, 45660, 45661); Marina Island, 1 (RBCM 12956); Maurelle Island, 1 (RBCM 12962); Mons, 4 (UBC 6400, 6402--6404); Nita Lake, 1 (UBC 6409); North Vancouver, 13 (UBC 3713, 3715, 3716, 3718, 6414, 10088, 10090, 10091, 10093--10097); Makay Creek, North Vancouver, 1 (UBC 48); Port

Moody, 7 (USNM 67048, 88838--88841, 88853, 88855); Port Neville, 13 (RBCM 9543--9545, 9594, 9596--9599, 9601, 9605, 9608, 9611, 9841); Powell River, 1 (UBC 49); Quadra Island, 2 (RBCM 9151, 9158); Head Rivers Inlet, 17 (CNM 15042, 15050, 15054, 15065, 15070, 15081, 15092, 15189, 15193, 15210, USNM 90174, 90176--90178, 90189, 90193, 90428); Thurston Harbor, Sonora Island, 1 (RBCM 12953); Stuart Island, 1 (PSM 15507); Sumas, 4 (USNM 88710, 102676, 102677, 102680); Gillies Bay, Texada Island, 1 (BYU 9703); Texada Island, 2 mi SSE Gillies, Harwood Pt., 3 (TCWC 45679--45681); Vananda, Texada Island, 2 (MVZ 70373, 70374); Vancouver, 3 (CNM 27254, RBCM 1636, UBC 6473); Grantham's Lodge, Vancouver, 8 (UMMZ 106703--106710); Point Grey, Vancouver, 10 (UBC 38, 174, 6415--6417, 6472, 6475--6477, 7436); Stanley Park, Vancouver, 1 (RBCM 5354); UBC Campus, Vancouver, 1 (UBC 8335); UBC Endowment Lands, Vancouver, 6 (UBC 10101, 10117, 10118, 10125, 10126, 10129); 23 km E Chilliwack Lake Rd., Vedder Crossing, 2 (RBCM 10879, 11010); Vedder Road, 1 (UBC 8910); Mile 2, Vedder Road, 1 (UBC 8909).

OREGON (125).--Clackamas Co.: 8 mi SE Molalla, 1 (PSM 6244); Still Creek Forest Camp, Mt. Hood National Forest, 1 (PSM 8632); 2 mi N, 4 mi E Sandy, 5 (CRCM 89-1502, 89-1504, 89-1507, 89-1511, 89-1514); 3.5 mi N, 8 mi E Sandy, 1 (CRCM 89-1457); 3.8 mi N, 8 mi E Sandy, 4 (CRCM 89-1655, 89-1657--89-1659); 4.2 mi N, 8 mi E Sandy, 1 (CRCM 89-1477); 6 mi S, 7 mi E Sandy, 10 (CRCM 89-1534, 89-1536--89-1540, 89-1543, 89-1544, 89-1548, 89-1549); 5 mi S, 6 mi E Zigzag, 24 (CRCM 89-1575, 89-1576, 89-1580, 89-1582, 89-1584--89-1589, 89-1592--89-1596, 89-1598--89-1601, 89-1604--89-1608); 9 mi N, 4 mi W

Zigzag, 12 (CRCM 89-1675, 89-1676, 89-1680--89-1682, 89-1686--89-1688, 89-1692, 89-1695--89-1697); Clatsop Co.: Seaside, 1 (PSM 8631); T8N, R7W, NE1/4 Sec. 18, 1 (PSM 26586); T8N, R10W, W1/2 Sec 6, 3 (PSM 26580, 26582, 26583); T9N, R6W, N1/2 Sec. 19, 1 (PSM 26585); Columbia Co.: 1 mi W Rainier, 1 (PSM 11931); T6N, R1W, SW1/4 Sec. 7, 2 (PSM 26575, 26578); 7 mi NE Vernonia, 1 (PSM 15502); Hood River Co.: 1 mi E Cascade Locks, Oxbow Fish Hatchery, 1 (PSM 6893); Multnomah Co.: 6 mi N, 11 mi E Sandy, 4 (CRCM 89-1435, 89-1436, 89-1438, 89-1439); 8 mi N, 11 mi E Sandy, 2 (CRCM 89-1168, 89-1170); 9 mi N, 10 mi E Sandy, 4 (CRCM 89-1744, 89-1745, 89-1748, 89-1750); 10 mi N, 10 mi E Sandy, 11 (CRCM 89-1263, 89-1267--89-1271, 89-1273, 89-1277, 89-1278, 89-1282, 89-1286); 10 mi N, 11 mi E Sandy, 7 (CRCM 89-1330, 89-1331, 89-1333--89-1337); 10 mi N, 11 mi E Sandy (3.2 mi SSE Ainsworth Park), 4 (CRCM 89-1322, 89-1325, 89-1326, 89-1328); 11 mi N, 10 mi E Sandy, 5 (CRCM 89-1216--89-1220); 11 mi N, 10 mi E Sandy (2.8 mi S Ainsworth Park), 1 (CRCM 89-1215); 8 mi N, 3 mi E Zigzag, 10 (CRCM 89-1388--89-1397); 10 mi N, 1.5 mi W Zigzag, 4 (CRCM 89-1144, 89-1147--89-1149); 11 mi N, Zigzag, 2 (CRCM 89-1196, 89-1197); Tillamook Co.: 2 mi upstream Miami River, 1 (PSM 7004).

WASHINGTON (149).--Chelan Co.: 22 mi W Lucerne, Lyman Lake, 1 (SDNHM 17023); Wenatchee Lake, 1 (OSUFW 6063); Clallam Co.: Deer Lake, 4 (UMMZ 106622--106624, 106627); Canyon Creek, 1 (CRCM 33B); Johnson's Ranch*, 3 (FMNH 6220, 6222, 6223); La Push, 1 (USNM 89153); Neah Bay, 5 (USNM 88497, 88523, 88527, 88536, 88537); Olympic National Park, 1 (PSM 25775); Deer Lake, Olympic National Park, 2 (PSM 2164, 2166); Heart O' the Hills Campground, Olympic

National Park, 3 (PSM 21049, 21050, 21052); Hurricane Ridge, Olympic
 National Park, 1 (PSM 3155); Moose Lake, Olympic National Park, 2
 (PSM 26396, 28503); Sand Point Trail, Ozette Lake, Olympic
 Peninsula, 4 (PSM 1404, 1412, 1421, 1422); Sol Duc Park, Olympic
 National Park, 1 (PSM 2890); Ozette Lake, Swan Bay, 2 (UMMZ 106639,
 106640); Port Angeles, 2 (CRCM 573, 5657); Sol Duc Hot Springs, 2
 (KU 10718, UMMZ 106636); Cowlitz Co.: Gilbert Lookout, 3 (PSM 11879,
 11882, 11885); Grays Harbor Co.: Aberdeen, 1 (USNM 24326); 3 mi S
 Copalis, Iron Springs, 2 (PSM 1428, 1439); Quiniaiet Lake
 [=Quinault Lake], 4 (USNM 89636, 89639, 89645, 89647); Jefferson
Co.: Blue Glacier, 1 (UMMZ 106621); Jackson Ranger Station, National
 Forest, Hoh River, 1 (UMMZ 106620); Kalalock [=Kalaloch], 3 (PSM
 2071, 2072, 2075); Mt. Kimta [=Kimta Peak], 2 (UMMZ 106628, 106629);
 Marmot Lake, Olympic National Park, 2 (PSM 3850, 3851); Rainbow
 Forest Camp, 1 (PSM 3297); Reflection Lake, 2 (UMMZ 106630, 106631);
King Co.: Baldi, 1 (PSM 10654); Lake Keechelus, 1 (UMMZ 57672);
 Lester, 1 (PSM 10636); Little Eagle Lake, 1 (PSM 13851); Lynn Lake,
 1 (PSM 10639); 8 mi SE North Bend, 3 (KU 4208, 4211, 4213); 1.7 mi E
 Scenic, 2 (KU 4224, 4235); 8.9 mi W Scenic, 1 (KU 4263); Mason Co.:
 Detroit, 3 mi SW Allyn, 1 (PSM 984); Lake Cushman, 2 (UMMZ 52905,
 USNM 66186); Pacific Co.: Fort Canby State Park, 2 (PSM 26573,
 26574); Long Beach, Gile Road, 2 (PSM 1447, 1475); Pierce Co.: Big
 Creek Forest Camp, 4.5 mi NE Ashford, 3 (UMMZ 88564, 88566, 88567);
 Chrystal Mt. [=Crystal], 2 (PSM 11935, 11936); Corral Pass, 1 (PSM
 25777); Corral Pass Campground, 1 (PSM 19266); Sunrise Beach Drive,
 Gig Harbor, 1 (PSM 28509); Longmire, 1 (CRCM 271); 1.5 mi up from

Nisqually gate, T15N, R7E, SE1/4 Sec. 33, 1 (PSM 16380); Paradise Creek, Mount Rainier, 1 (USNM 89577); Bench Lake, Rainier National Park, [Lewis Co.] 1 (PSM 16459); Kautz Creek headwaters, Rainier National Park, 3 (PSM 4279--4281); Owyhigh Lake, near outlet, Rainier National Park, 1 (PSM 5941); Tahoma Creek, Rainier National Park, 1 (PSM 4282); Tacoma, 1 (UMMZ 57663); Tacoma headworks, 1 (PSM 13852); Skagit Co.: 0.7 mi NE Rosario Beach, Fidalgo Island, 2 (CRCM 85-592, 85-599); Hamilton, 1 (USNM 24311); Mount Vernon, 5 (USNM 24309, 76318, 76490, 76491, 88808); Skamania Co.: 5.5 mi N, 6.5 mi W Carson; T4N, R7E, NE1/4 Sec. 33, 5 (KU 145774--145777, 145779); 5.5 mi N, 7.5 mi W Carson; T4N, R7E, E1/2 Sec. 32, 6 (KU 145782--145787); 8 mi N, 8 mi W Carson; T4N, R7E, N1/2 Sec. 20, 5 (KU 145788--145790, 145792, 145794); 10 mi N, 2 mi W Carson; T4N, R8E, SW1/4 Sec. 7, 6 (KU 145797--145802); 11 mi N, 9.5 mi W Carson; T5N, R6E, NW1/4 Sec. 25, 7 (KU 145804--145809, OSUFW X1887); 12 mi N, 9.5 mi W Carson; T5N, R6E, N1/2 Sec. 24, 3 (KU 145815, 145816, 145818); 13 mi N, 8.5 mi W Carson; T5N, R7E, N1/2 Sec. 18, 2 (KU 145819, 145820); 44 mi N, 9.5 mi W Carson; T7N, R6E, SW1/4 Sec. 12, 10 (KU 145821--145830); Council Pass, 1 (PSM 1396); 5 mi N Skamania, 1 (CRCM 82-462); Snohomish Co.: Granville, 1 (USNM 89651); 3 mi up Rapid River from Beckler River, 1 (PSM 14146); Thurston Co.: Tenino, 1 (USNM 89649); Whatcom Co.: Mt. Baker, 1 (CRCM 58); Skyline Ridge, Mt. Baker, 1 (CRCM 51).

Sorex monticolus shumaginensis Merriam

1900. Sorex alascensis shumaginensis Merriam, Proc. Washington Acad. Sci., 2:18, March.
1901. Sorex glacialis shumaginensis Elliot, Field Columb. Mus., Publ. 45, Zool. Ser., 2:372.
1902. Sorex obscurus shumaginensis Allen, Bull. Amer. Mus. Nat. Hist., 16:228.
1955. Sorex vagrans shumaginensis Findley, Univ. Kansas Publ., Mus. Nat. Hist., 9:42, December.
1977. S[orex]. m[onticolus]. shumaginensis Hennings and Hoffmann, Occas. Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.

Holotype.--USNM 97993, original number 2210 (Fisher catalog); adult male collected July 1899 from "Popof Island, Shumagin Islands, Alaska" by D. Saunders. Type specimen examined and measured.

Distribution.--S. m. shumaginensis occurs in western Alaska from western Prince William Sound, west to the Bering Sea (including the Kenai and Alaska peninsulas), and north to the Noatak Valley in the western Brooks Range (Fig. 13).

Subspecific comparisons and remarks.--Some shrews from the Kenai and Alaska Peninsulas had weaker Geisser classifications than those of the mainland populations but remain referable to S. m. shumaginensis. Because peninsular populations are relatively

isolated, it is not surprising that these shrews have diverged slightly from those composing the remainder of the mainland population.

Specimens examined.--ALASKA (194).--Lake Aleknagik, 1 (USNM 224892); 1 mi NE Anchorage, 1 (KU 21070); 20 mi NE Anchorage, 1 (KU 42598); Conner's Lake Bog, Raspberry Road, Anchorage, 1 (MVZ 162064); Aniak, 1 (USNM 180429); Anuk Lake, Noatak Valley, eastern most lake, headwaters Kaluich Creek, 1 (USNM 505018); Becharof Lake, Alaska Peninsula, 7 (USNM 119638, 119641, 119642, 119644--119647); Campbell Klatt Bog, 300 m SE Klatt Rd & Victor Rd jct., Anchorage, 2 (MVZ 162066, 162067); Chignik, 2 (USNM 159542, 179961); Chignik, upper Lake, 1 (USNM 176657); Chignik, Chignik Bay, 1 (USNM 176658); Cold Bay, Alaska Peninsula, 8 (USNM 119656--119658, 119661, 119662, 119664, 119666, 120031); Fishhook, 3 (CNM 40283--40285); East base Frosty Peak, Alaska Peninsula, 9 (USNM 177036--177038, 177041, 177042, 177045--177047, 177219); Halibut Cove, Kachemak Bay, Cook Inlet, 1 (UIMNH 17912); Hazen Point, Izembek Bay, 2 (USNM 246471, 246477); Hope, Cook Inlet, 8 (USNM 107135, 107139, 107141, 107143, 107145, 107146, 107153, 107156); near Hope, 5 (CNM 40286--40290); mountains near Hope, Cook Inlet, 4 (USNM 107105, 107107, 107111, 107155); Iliamna, 1 (CMNH 62274); Kakhtul River, 4 (USNM 119723, 119726, 119727, 119732); Kanatak, Portage Bay, Alaska Peninsula, 2 (USNM 119649, 119652); King Cove, Alaska Peninsula, 11 (USNM 177020--177022, 177025, 177027, 177028, 177213, 177301, 177305--177307); Kokwok, Nushagak River, 2 (USNM 119668, 119670); 80 mi

upstream Kokwok, 1 (USNM 180561); Morzhovoi Bay, Alaska Peninsula, 4 (USNM 177030, 177031, 177034, 177035); Ninilchik, 22 (CMNH 76010--76012, 76015--76017, 76020--76035); Nome, 1 (SDNHM 46); Nushagak, 13 (USNM 119678, 119683, 119685--119687, 119691--119694, 119696--119699); Pedro Bay, 3 (CMNH 62278--62280); Pilot Station, Yukon River, 2 (UIMNH 53554, 53555); Popof Island, 3 (TCWC 20639, 20640, USNM 273099); Popof Island, Shumagin Islands, 1 (USNM 97993); Sanak, 1 (USNM 273096); Sanak Island, 1 (USNM 263300); Sapsuk River, Alaska Peninsula, 3 (UAM 13767, 13773, 13775); Sapsuk River, Port Moller Quad, Alaska Peninsula, 5 (UAM 13760--13762, 13765, 13771); Scammon Bay, 2 (CMNH 70695, 70696); Seward, 7 (CMNH 62275, 62276, USNM 273092--273094, 273100, 273101); 13 km NW Seward, Kenai Peninsula, 1 (CMNH 60950); Mountain Climber Roadhouse, Skwentha River, 1 (USNM 242751); 2 mi W Soldotna, Kenai Peninsula, 2 (CRCM 80-27, 80-753); Upper Susitna River, Talkeetna Mts., 1 (UAM 13717); Upper Susitna River, Devil's Canyon area, Talkeetna Mts., 1 (UAM 14802); Upper Susitna River, Tsusena Creek area, Talkeetna Mts., 3 (UAM 14801, 14803, 14804); 7 mi N, 4 mi W Talkeetna, 10 (CMNH 76036--76039, 76042, 76044--76048); Tyonek, Cook Inlet, 2 (FMNH 24280, 24281); Tyoonok [=Tyonek], Cook Inlet, 21 (USNM 107049--107051, 107053--107056, 107063, 107065--107068, 107114--107117, 107121, 107123, 107124, 107202, 107203); Ugaguk [=Egegik] River, Alaska Peninsula, 1 (USNM 119636); Unga, Unga Island, 1 (USNM 177300); Urilia Bay, Unimak Island, 1 (USNM 246467); Cottonwood Creek, 1 mi SE Wasilla, 1 (UMNH 28197).

Sorex monticolus soperi Anderson and Rand

1945. Sorex obscurus soperi Anderson and Rand, Can. Field-Nat.,
59(2):47, March-April.
1955. Sorex vagrans soperi Findley, Univ. Kansas Publ., Mus. Nat.
Hist., 9:48, December.
1977. S[orex]. m[onticolus]. soperi Hennings and Hoffmann, Occas.
Papers Mus. Nat. Hist., Univ. Kansas, 68:4, July.

Holotype.--CNM 18249, original number 4264; adult male collected
21 September 1940 from "2.5 mi NW Lake Audy, Riding Mountain
National Park, southwestern Manitoba, Canada" by J. D. Soper. Type
specimen not examined.

Distribution.--S. m. soperi occurs in the vicinity of Prince
Albert National Park, Saskatchewan and extends east to include
Riding Mountain National Park, Duck Mountain, Porcupine Mountains,
The Pas, and Thompson, Manitoba (Wrigley et al., 1979).

Subspecific comparisons and remarks.--I was able to locate 18
specimens classified as S. m. soperi; 14 were not received in
requested loan material. Because the loan requests for this study
specified unbroken skulls, it is possible that like many soricid
skulls in collections, all of these specimens have broken crania.
Only one unbroken specimen referable to S. m. soperi was examined in
this study. The distribution presented is based wholly on published

records of specimens (Wrigley et al., 1979). Therefore, the distribution should be viewed as tentative.

Specimens examined.--SASKATCHEWAN (1).--Spruce River, 1 (UAMZ 1363).

DISCUSSION

Sorex neomexicanus was recognized previously as a subspecies of Sorex monticolus (Bailey, 1913; Hennings and Hoffmann, 1977) but is herein recognized as a distinct species. S. neomexicanus occurs in the Sacramento and Capitan mountains, an isolated mountainous region in southcentral New Mexico. This region is approximately 282 km south of the known extent of Pleistocene glaciation in New Mexico except for possible glaciation on the highest peak of the Capitan Mountains, Sierra Blanca (Ellis, 1935). It is possible that during the Pleistocene glaciation, southern New Mexico acted as a boreal-forest refugium for S. monticolus-type shrews. During the warmer interglacial period, after the most recent glaciation, the valleys became too arid for survival and these shrews survived in forested, montane regions of New Mexico. The Capitan and Sacramento mountains are sufficiently isolated from other montane regions in the state to reduce or eliminate gene flow between these populations of shrews.

Sorex monticolus as herein defined exhibits little morphometric variation in comparison with several nearby congeners (S. sonomae, S. bairdi, S. pacificus--Carraway, 1990; S. bendirii--Jackson, 1928). Within nominate races, even those widely distributed geographically (e.g., S. m. obscurus), coefficients of variation for all characters are ≤ 7.88 (Table 1). Even among nominate races, the only obvious morphometric variation is a partial north-south cline in greatest length of skull (Fig. 18). There is a general trend of increasing size from south to north. The southern subspecies




Fig. 18.--Mean and range of greatest length of skull for all subspecies of Sorex monticolus and the approximate distribution of each subspecies demonstrating the trend of increasing skull length from south to north.

restricted to isolated mountains (S. m. monticolus and S. m. parvidens; Fig. 18) have the shortest skulls of all S. monticolus (Table 1). Greatest lengths of skull for these two taxa were significantly shorter than those of all other taxa; they averaged 10% shorter than the taxa with the longest skulls. The subspecies found in the northern coastal and insular areas of southeast Alaska and British Columbia (S. m. longicaudus and S. m. prevostensis) have the longest skulls (Table 1, Fig. 18). Greatest lengths of skull for these two taxa were significantly greater than those of all other taxa. The remaining subspecies, in decreasing skull length are S. m. malitiosus, S. m. insularis, S. m. calvertensis, S. m. alascensis, S. m. elassodon, S. m. setosus, S. m. isolatus, S. m. soperi, S. m. shumaginensis, and S. m. obscurus (Table 1). These taxa do not conform to the north-south reduction of skull length (Fig. 18). Among these taxa, the insular populations tend to have longer skulls (except S. m. elassodon) and the mainland populations tend to have shorter skulls. Races with long skulls that occur on the mainland (S. m. longicaudus and S. m. alascensis) also occur on some islands, and the mainland portion of their distributions are restricted to a narrow band along the coasts of Alaska and British Columbia (Fig. 18).

Bergmann's rule states that in warm-blooded species, races from colder climates (increased latitude or altitude) will be larger than races of the same species from warmer climates (Mayr, 1963; McNab, 1971). In some cases this correlation of body size and latitude is a reflection of the size of available prey and is correlated

negatively as frequently as it is correlated positively (McNab, 1971). In European shrews (Soricinae), Hanski (1985) found that the most northern species are smaller than southern species. Many birds and mammals (especially rodents) also attain a large body size on islands and mountains (Foster, 1964; Grant, 1965; McNab, 1971). These regions typically have a low diversity of species, possibly resulting in less competition for food. The available food might exist in a greater range of food-particle size which would be an advantage to a larger body size (Grant, 1965; Hanski, 1985; McNab, 1971). There is much confusion, however, regarding the size of many mammals on islands (Foster, 1964). In a survey of the literature, Foster (1964) found a tendency toward insular dwarfism among carnivores, artiodactyls, and lagomorphs, a tendency toward gigantism in rodents, and no conclusive evidence for other orders of mammals. Among insectivores, Foster (1964) found four species smaller than mainland counterparts, four the same, and one larger. Insular adaptations (e.g., dwarfism and gigantism) can occur more rapidly than they would on the mainland if the population is relatively small and gene flow is restricted (Foster, 1964).

The pattern of geographic variation observed in the northern half of the range of S. monticolus is largely consistent with the tendency towards insular gigantism. Except for S. m. elassodon, insular taxa are larger than mainland taxa. S. m. elassodon occurs on the Queen Charlotte Islands and the outer islands of the Alexander Archipelago but remains relatively small; this may be a result of having a comparatively large distribution. If the gene

pools for all of the large insular forms are small, it would be possible for a change in body size to occur in a relatively short period of time. Insular gigantism could be explained by having a shortage of food available on islands that would result in an advantage for larger individuals. In contradiction, moderate temperatures and abundant precipitation on the coastal islands should favor populations of invertebrates on which soricines feed. S. m. shumaginensis occurs in the northwestern portion of Alaska; its distribution is relatively large and includes interior mainland as well as coastal and peninsular regions. This taxon does not conform to Bergmann's rule but instead remains consistent with the pattern of insular gigantism and smaller shrews in the mainland populations.

The morphometric variation among nominate races is sufficient to warrant continued separation at the subspecies level of all taxa except S. m. calvertensis and S. m. elassodon. Were it not for differences in pelage color, based on my morphometric analysis S. m. calvertensis and S. m. elassodon should be synonymized.

LITERATURE CITED

- Allen, J. A. 1902. List of mammals collected in Alaska by the Andrew J. Stone expedition of 1901. Bull. Am. Mus. Nat. Hist. 16:215-230.
- Anderson, R. M., and A. L. Rand. 1945. A new form of dusky shrew from the prairie provinces of Canada. Can. Field-Nat. 59:47-48.
- Bailey, V. 1913. Ten new mammals from New Mexico. Proc. Biol. Soc. Washington 26:129-134.
- Carraway, L. N. 1990. A morphologic and morphometric analysis of the "Sorex vagrans species complex" in the Pacific Coast region. Spec. Publ., Mus., Texas Tech Univ. 32:1-76.
- Choate, J. R. 1972. Variation within and among populations of the short-tailed shrew in Connecticut. J. Mamm. 53:116-128.
- Corel Corporation. 1992. Corel Draw. User's manual. Ontario, Canada: Corel Corporation.
- Cowan, I. M. 1941. Insularity in the genus Sorex on the north coast of British Columbia. Proc. Biol. Soc. Washington 54:95-108.
- Diersing, V. E. 1980. Systematics and evolution of the pygmy shrews (subgenus Microsorex) of North America. J. Mamm. 61:76-101.
- Elliot, D. G. 1899. Catalogue of mammals from the Olympic Mountains Washington with descriptions of new species. Field Columbian Mus., Publ. No. 32, Zool. Ser. 1:239-276 + pls. 41-61.
- . 1901. A synopsis of the mammals of North America and the adjacent seas. Field Columbian Mus., Publ. No. 45, Zool. Ser. 2:1-522.
- . 1905. A check list of mammals of the North American continent, the West Indies, and the neighboring seas. Field Columbian Mus., Publ. 105, Zool. Ser. 6:1-761.
- Ellis, R. W. 1935. Glaciation in New Mexico. Univ. New Mexico Bull., No. 276, Geol. Ser. 5(1):1-32.
- Findley, J. S. 1955. Speciation of the wandering shrew. Univ. Kansas Publ., Mus. Nat. Hist. 9:1-68.

- Foster, J. B. 1964. Evolution of mammals on islands. *Nature* 202:234-235.
- George, S. B., and J. D. Smith. 1991. Inter- and intraspecific variation among coastal and island populations of Sorex monticolus and Sorex vagrans in the Pacific Northwest. Pp. 75-91 in *The Biology of the Soricidae* (J.S. Findley and T. L. Yates, eds.). Albuquerque: Univ. of New Mexico Press.
- Grant, P. R. 1965. The adaptive significance of some size trends in island birds. *Evolution* 19:355-367.
- Hall, E. R. 1938. Variation among insular mammals of Georgia Strait, British Columbia. *Am. Nat.* 72:453-463.
- Hall, E. R. 1981. *The mammals of North America*. Second ed. New York: John Wiley & Sons.
- Hanski, I. 1985. What does a shrew do in an energy crisis? Pp. 247-252 in *Behavioural ecology, ecological consequences of adaptive behaviour*. 25th Symp. British Ecol. Soc., R. M. Sibly and R. H. Smith, eds. Oxford: Blackwell Sci. Publ.
- Hennings, D., and R. S. Hoffmann. 1977. A review of the taxonomy of the Sorex vagrans species complex from western North America. *Occas. Papers Mus. Nat. Hist., Univ. Kansas* 68:1-35.
- Jackson, H. H. T. 1918. Two new shrews from Oregon. *Proc. Biol. Soc. Washington* 31:127-130.
- . 1919. An unrecognized shrew from Warren Island, Alaska. *Proc. Biol. Soc. Washington* 32:23-24.
- . 1921. Two unrecognized shrews from California. *J. Mamm.* 2:161-162.
- . 1922. New species and subspecies of Sorex from western America. *J. Washington Acad. Sci.* 12(11):262-264.
- . 1928. A taxonomic review of the American long-tailed shrews. *N. Am. Fauna* 51:1-238.
- Johnson, M. L., and B. T. Ostenson. 1959. Comments on the nomenclature of some mammals of the Pacific Northwest. *J. Mamm.* 40:571-577.
- Mayr, E. 1963. *Animal species and evolution*. Cambridge, Massachusetts: Belknap Press.
- McNab, B. K. 1971. On the ecological significance of Bergmann's rule. *Ecology* 52:845-854.

- Merriam, C. H. 1890. Results of a biological survey of the San Francisco Mountain region and desert of the Little Colorado, Arizona. N. Am. Fauna 3:1-136 pp., 13 pls., 5 maps.
- . 1891. Results of a biological reconnoissance of Idaho, south of latitude 45 and east of the thirty-eight meridian, made during the summer of 1890, with annotated lists of the mammals and birds, and descriptions of new species. N. Am. Fauna 5:1-113.
- . 1895. Synopsis of the American shrews of the genus Sorex. N. Am. Fauna 10:57-98.
- . 1900. Papers from the Harriman Alaska expedition. I. Descriptions of twenty-six new mammals from Alaska and British North America. Proc. Washington Acad. Sci. 2:13-30.
- Microsoft Corporation. 1988. Microsoft QuickBASIC, version 4.5. Redmond, Washington: Microsoft Corp.
- Osgood, W. H. 1901. Mammals of the Queen Charlotte Islands. N. Am. Fauna 21:25-37.
- Owen, J. G., and M. A. Chmielewski. 1985. On canonical variates analysis and the construction of confidence ellipses in systematic studies. Syst. Zool. 34:366-374.
- Pimentel, R. A., and J. D. Smith. 1986. BIOETAT II: a multivariate statistical toolbox. Second ed. Placentia, California: Sigma Soft.
- Rudd, R. L. 1955. Age, sex, and weight comparisons in three species of shrews. J. Mamm. 36:323-339.
- Statistical Graphics Corporation. 1987. Statgraphics: statistical graphics system. User's guide. Rockville, Maryland: Statistical Graphics Corp.
- van Zyll de Jong, C. G. 1980. Systematic relationships of woodland and prairie forms of the common shrew, Sorex cinereus cinereus Kerr and S. c. haydeni Baird, in the Canadian prairie provinces. J. Mamm. 61:66-75.
- Wrigley, R. E., J. E. Dubois, and H. W. R. Copland. 1979. Habitat, abundance, and distribution of six species of shrews in Manitoba. J. Mamm. 60:505-520.