

Walnut

2014 Pest Management Guide for the Willamette Valley

EM 8421 · Revised March 2014

Jay W. Pscheidt, Ed Peachey, and Vaughn Walton

The chemicals, formulations, and rates listed for insect, mite, and disease control are among the best recommendations based on label directions, research, and orchard use experience. Only a thorough knowledge of the orchard, its variety, tree size and density, canopy characteristics, pest complex, and past pest problems will enable you to correctly select chemicals, rates, amount of water used per acre, and method of application for optimum pest control. Occasionally, different formulations of a product or like formulations containing a different amount of active ingredient also are registered and effective for use on the pests listed. These products also may be used; we do not intend to discriminate against them. You may wish to consult their labels and determine whether their use confers advantages over the products listed in this guide.

Always refer to the pesticide label for use instructions. It is the legal document regarding use patterns. Two questions frequently are asked about the chemical control of insects and diseases: “How much chemical do I use per acre?” and “What is the least amount of water I need per acre to apply in my concentrate sprayer?” Notice that the schedule below suggests an amount of formulated product (not active ingredient) to use per acre. This amount is based on a “typical” middle age and density orchard with moderate pest pressure. Common sense indicates that less material may be needed (than that given) for 1- to 4-year-old orchards. Conversely, more chemical (within label limits) may be required for large, mature trees experiencing heavy pest pressure from multiple pests.

Stages

Early Prebloom
Late Prebloom
Postbloom

Not shown

June–July; July–August



Early Prebloom



Late Prebloom



Postbloom

Many insecticide labels today indicate the minimum amount of water needed per acre to apply concentrate sprays of insecticides, as well as how to calculate the amount of chemical needed per acre in a concentrate sprayer. **CHECK LABEL BEFORE SPRAYING!** Some label directions indicate dilute applications only.

Also:

1. Make sure any tank-mixes of pesticides are compatible. For example, the elevated pH of some boron spray solutions weakens many insecticides.
2. Use adjuvants and spreader stickers with caution.

Important information

1. Diazinon is not labeled for use on walnuts. Existing supplies with the older label still may be used according to label instructions.
2. Blackline of walnuts is a serious disease that can't be controlled by use of pesticides. For more information, see EM 8907, *Growing Walnuts in Oregon*. Order from Publication Orders, Extension & Station Communications, Oregon State University, 422 Kerr Administration, Corvallis, OR 97331-2119 (puborders@oregonstate.edu; 800-561-6719 or 541-737-2513).

Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.

Walnut Pest Control Recommendations

Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.

Early Prebloom (late March to early April, when catkins begin to enlarge)

Pest or disease/ Material	Amount of product per acre	Comments/Reentry interval/Preharvest interval (PHI)
Blight and downy leaf spot		
bordeaux 4-2-100	—	The low-lime formula reduces the possibility of bordeaux foliage injury. See footnote 2, page 4.
C-O-C-S WDG	4–7.7 lb	48-hour reentry.
Copper-Count-N	8–12 qt	48-hour reentry.
Cuprofix Ultra 40 Disperss	5–10 lb	48-hour reentry.
Kocide 3000	3.5–7 lb	48-hour reentry.
Nordox 75 WG	5–8 lb	12-hour reentry
Nu-Cop 50DF	4–8 lb	48-hour reentry.
Python 27 AG	30–50 fl oz/ 100 gal water	48-hour reentry.

Late Prebloom (early to mid-May, when shoots begin to expand)

Pest or disease/ Material	Amount of product per acre	Comments/Reentry interval/Preharvest interval (PHI)
Blotch (anthracnose)		
<i>Note:</i> No materials are registered specifically for this disease. The following are suggested, but there is no information about efficacy.		
Adament 50WG	4–8 oz	Group 3 + 11 fungicide. 12-hour reentry. 60-day PHI.
Gem 500SC	1.9–3.8 oz	12-hour reentry.
Luna Experience	8.8–17 fl oz	Group 3 + 7 fungicide. 12-hour reentry. 35-day PHI.
Luna Sensation	5.6–7.6 fl oz	Group 7 + 11 fungicide. 12-hour reentry. 60-day PHI.
Pristine	10.5–14.5 oz	Group 7 + 11 fungicide. 12-hour reentry. 14-day PHI.
Quadris Top	12–14 fl oz	Group 3 + 11 fungicide. 12-hour reentry. 45-day PHI.
Quilt Xcel	14–21 oz	Group 3 + 11 fungicide. 12-hour reentry. 60-day PHI.
Regalia	1–4 qt	Use with a surfactant and on 7-day intervals. 4-hour reentry.
Syllit FL	4.5 pt	48-hour reentry. 7-day PHI.
Tilt	4–8 fl oz	12-hour reentry. 60-day PHI.

Blight
See materials listed for Early Prebloom Stage.

Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.

Postbloom (late May)

Pest or disease/ Material	Amount of product per acre	Comments/Reentry interval/Preharvest interval (PHI)
--------------------------------------	---------------------------------------	--

Blight

See materials listed for Early Prebloom Stage.

June–July

Pest or disease/ Material	Amount of product per acre	Comments/Reentry interval/Preharvest interval (PHI)
--------------------------------------	---------------------------------------	--

Aphids

Asana XL	10–16 oz	Do not use more than 0.2 lb ai/A per season. Do not feed or graze livestock on treated orchard floors. 21-day PHI.
Assail 70WP	1.1–4.1 oz	14-day PHI.
Brigade WSB	0.05–0.2 lb ai/A	7-day PHI.
Cobalt	26–57 oz	Do not make more than 2 applications per season. 14-day PHI.
Esteem 35WP	3.2–4 oz	Do not exceed 2 applications per season. 12-hour reentry. 21-day PHI.
insecticidal soap (M-Pede)	2%	0-day PHI.
Lorsban 75WG	2–2.67 lb	Make no more than 1 dormant stage application and no more than 2 foliar spray applications on walnuts per season. Do not graze orchard floors. 24-hour reentry. 14-day PHI.
malathion 57% EC	See labels.	See footnote 1, page 4. 7-day PHI.
Movento	6–9 oz	7-day PHI.
Provado 1.6	3.5–7 oz	7-day PHI.
Success 2L	4–10 oz	Do not apply fewer than 14 days apart. Do not use more than 29 oz/A per season. 14-day PHI.
Supracide 2E	4–8 pt	Apply as a cover spray when aphids appear. Do not graze orchard floors. 48-hour reentry, depending on rate. 7-day PHI.
Warrior II	1.28–2.56 fl oz	14-day PHI.

Aphids, codling moth

Note: Monitor codling moth with pheromone traps beginning in June. See footnote 3, page 4.

Scales (apply when crawlers appear; chemical control of scales usually is not necessary)

Esteem 35WP	3.2–4 oz	Do not exceed 2 applications per season. 12-hour reentry. 21-day PHI.
Intrepid 2F	8–16 fl oz	4-hour reentry. 14-day PHI.
Movento	6–9 oz	7-day PHI.
Supracide 2E	4–8 pt	Do not graze orchard floors. Do not make more than 2 cover sprays during the growing season. 2- to 14-day reentry, depending on rate. 7-day PHI.

Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.

July–August		
Pest or disease/ Material	Amount of product per acre	Comments/Reentry interval/Preharvest interval (PHI)
Walnut husk fly		
Ambush 25 WG	12.8–25.6 oz	Do not graze orchard floor. 12-hour reentry. 12-day PHI.
Asana XL	9.6–19.2 oz	See restrictions under aphids. 21-day PHI.
Baythroid XL	2.4–2.8 oz	14-day PHI.
Brigade WSB	0.1–0.2 lb ai/A	7-day PHI.
Cobalt	26–57 oz	Do not make more than 2 applications per season. 14-day PHI.
Delegate WG	3–7 oz	No more than 4 applications per season. 14-day PHI.
Imidan 70W	4.33–8.5 lb	Do not apply more than 5 times per season. 28-day PHI.
Lorsban 75WG	2.67 lb	Make no more than 1 dormant stage application and no more than 2 foliar spray applications on walnuts per season. Do not graze orchard floors. 14-day PHI.
Lorsban 4E	4 pt	Make no more than 1 dormant stage application and no more than 2 foliar spray applications on walnuts per season. Do not graze orchard floors. 14-day PHI.
malathion 57% EC	See labels.	See footnote 1, page 4. 7-day PHI.
Pounce 25 WG	12.8–16.0 oz	Do not graze orchard floor. 1-day PHI.
Success Naturalyte Insect Control	4–10 oz	Do not apply treatments fewer than 7 days apart. Do not use more than 29 oz/A per season. Entrust is the organic formulation. GF-120 NF is the spinosad bait formulation. 1-day PHI.
Warrior II	1.28–2.56 fl oz	14-day PHI.
Fall webworm		
<i>Note:</i> Insecticides labeled for walnut husk fly, diazinon, and <i>Bacillus thuringiensis</i> will control this pest. Only spot treatments are necessary. Completely drench the infested branch. Applications made when the larvae are small are most effective.		

Footnotes

1. More than one type of formulation usually is available for most insecticides. Lower rates can be used on smaller trees.
2. Bordeaux 4-2-100 means 4 pounds of copper sulfate plus 2 pounds of hydrated lime in 100 gallons of water. In any bordeaux formula, ingredients always are listed in the same order—copper sulfate, hydrated lime, then gallons of water.
3. Early summer nut drop can be a result of codling moth infestation. Inspect nuts for larvae and monitor moths with pheromone traps. *Walnut Production Manual* (University of California Publication 3373) provides information on codling moths as a pest of walnuts.

Note: Bacteria resistant to copper-based products has occurred in California and easily could occur in Oregon.

Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.

Table 1. Quick Reference Guide for Herbicides Labeled for Use in Fruit and Nut Crops

- Shaded boxes indicate the herbicide is labeled for use in that crop.
- Nonbearing (NB) indicates the herbicide is labeled only for crops that will not be harvested for 1 year. It has a 365-day preharvest interval.
- Herbicides in *bold, italic* type are recommended for new plantings.
- For more complete information, please refer to the *PNW Weed Management Handbook*: <http://pnwhandbooks.org/weed/>.

Ingredient common name (herbicide mode of action)	Product name example	Nuts			Pome fruits		Stone fruits						Rates
		Chestnut	Hazelnut	Walnut	Apple	Pear	Apricot	Cherry	Nectarine	Peach	Plum	Prune	
Applications that persist in soil and are soil active													
diuron (7)	Karmex												See label for crop-specific application rates.
dichlobenil (20)	Casoron												4–6 lb ai/A (100–150 lb/A Casoron). Apply in cold, wet weather.
<i>isoxaben</i> (21)	<i>Trellis, Gallery</i>				NB	NB	NB	NB	NB	NB	NB	NB	0.5–1 lb ai/A (0.66–1.33 lb/A product)
indaziflam (29)	Alion												0.065–0.085 lb ai/A (5–6.5 oz/A product) depending on soil texture
<i>napropamide</i> (3)	<i>Devrinol</i>												5 lb ai/A (50 lb/A Devrinol 10-G)
norflurazon (12)	Solicam												1.97–7.8 lb ai/A (2.5–10 lb/A Solicam)
<i>oryzalin</i> (3)	<i>Surflan</i>												2–6 lb ai/A (2–6 qt/A Surflan)
<i>pendimethalin</i> (3)	<i>Prowl</i>												Prowl H ₂ O: 1.9–6 lb ai/A (2–6.3 qt/A) depending on desired length of weed control and crop
<i>pronamide</i> (3)	<i>Kerb</i>												1–4 lb ai/A (2–8 lb/A). Rate depends on species present and soil texture.
simazine (5)	Princep												See product labels for rates. Princep Caliber 90 is a Special Local Needs label (OR-080038) for sweet cherries only.
terbacil (5)	Sinbar					NB	NB	NB				NB	0.4–0.8 lb ai/A (0.5–1 lb/A), newly established; 2–4 lb/A Sinbar, bearing, depending on soil type
<i>trifluralin</i> (3)	<i>Treflan</i>												0.5–1 lb ai/A (1–2 pt/A Treflan 4EC)
trifluralin (3) + isoxaben (21) + oxyfluorfen (14)	Showcase	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	2.5–5 lb ai/A (100–200 lb/A Showcase)
Applications that persist in soil and have both soil and foliar activity													
<i>flumioxazin</i> (14)	<i>Chateau</i>												0.188–0.38 lb ai/A (6–12 oz/A Chateau WDG). Note differences in rates and uses in SW and WDG labels. Avoid contact with green bark on small trees.
oxyfluorfen (14)	Goal												1.25–2 lb ai/A (5–8 pt/A Goal 2XL)

Table continues on next page

Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.

Ingredient common name (herbicide mode of action)	Product name example	Nuts			Pome fruits		Stone fruits						Rates
		Chestnut	Hazelnut	Walnut	Apple	Pear	Apricot	Cherry	Nectarine	Peach	Plum	Prune	
CONTINUED—Applications that persist in soil and have both soil and foliar activity													
penoxsulam (2)	Pindar GT												(1.5–3.0 pints/A)
rimsulfuron (2)	Matrix												0.063 lb ai/A (4 oz/A Matrix FNV per year)
saflufenacil (14)	Treevix												0.045 lb ai/A (1 oz/A Treevix)
Postemergence contact and translocated herbicides													
acetic acid	WeedPharm												
carfentrazone (14)	Aim												Green sucker control in hazelnuts. 0.031 lb ai/A (2 fl oz/A Aim EC)
clethodim (1)	Select Max		NB	NB	NB	NB	NB	NB		NB		NB	0.068–0.121 lb ai/A (9–16 fl oz/A Select Max)
clopyralid (4)	Stinger												Apples: 0.094–0.25 lb ae/A (0.25–0.66 pints/A Stinger) Others: 0.12–0.25 lbs ae/A (0.33–0.66 pints/A Stinger)
diquat (22)	Reglone		NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	0.375–0.5 lbs ai/A (1.5–2 pints)
fluazifop (1)	Fusilade DX		NB	NB	NB	NB							0.25–0.375 lb ai/A (16–24 oz/A Fusilade DX). Refer to specific grassy weeds listed on label.
glufosinate (10)	Rely 280												Sucker control. 0.88–1.46 lb ai/A (1.5 to 2.5 quarts/A Rely)
glyphosate (9)	Roundup												General weed control and grass suppression in row middles. Read label carefully for crops listed and geographic location.
halosulfuron (2)	Sandea												Apples: 0.035–0.094 lb ai/A (0.75–2 oz/A) Nut crops: 0.031–0.063 lb ai/A (2/3–1 1/2 oz/A)
paraquat (22)	Gramoxone												Green sucker control in hazelnuts. 0.625–1 lb cation/A (2.5–4 pt/A Gramoxone; 1.7–2.7 pt/A Firestorm)
pyraflufen (14)	Venue												0.001–0.005 lb ai/A (0.7–4 fl oz product/A)
sethoxydim (1)	Poast										NB	NB	Grass suppression in row middles. 0.28–0.47 lb ai/A (1.5–2.5 pt/A product)
2,4-D (4)	2,4-D												Green sucker control in hazelnuts. 0.7–0.95 lb ai/A (1.5–2 pt/A Saber)

Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.

OSU Internet Resources for Plant Protection

Information regarding plant protection is available from several sources at OSU. The following listings are excellent examples:

- OSU Integrated Plant Protection Center. Online weather data and degree day information for insect pests and diseases (<http://uspest.org/wea/>)
- Codling moth development information (<http://ippc2.orst.edu/cgi-bin/ddmodel.pl?clm>)
- Pacific Northwest Plant Disease Management Handbook (<http://pnwhandbooks.org/plantdisease>)
- Pacific Northwest Insect Management Handbook (<http://pnwhandbooks.org/insect>)
- Pacific Northwest Weed Management Handbook (<http://pnwhandbooks.org/weed>)

Oregon Poison Center

The Oregon Health & Science University
3181 S.W. Sam Jackson Park Road
Portland, OR 97239
Phone: 1-800-222-1222

If a person has collapsed or is not breathing, dial 911.

Basic Elements of Safe Pesticide Use

- Always read the label with care. This is the first step in selecting the right material for the job. Never rely on your memory. Before opening the container, pay strict attention to warnings and cautions printed on the label.
- Keep all pesticide and spray materials out of the reach of children, pets, and irresponsible persons. Storage outside of the home, away from food and feed, and under lock and key is the safest method.
- Store only in the original container and keep tightly closed.
- NEVER smoke, eat, or drink while applying pesticides.
- Avoid inhalation or direct contact. Always wear protective clothing and safety devices as recommended on the label.
- Avoid spills. If spills occur, take immediate action to remove contaminated clothing and wash thoroughly.
- After each application, bathe and change to clean clothing. Wash clothing after each use. Always use fresh clothing when starting new application.
- Avoid contamination of fish ponds and water supplies. Cover feed and water containers when treating around livestock or pet areas.
- Keep separate equipment for use with hormone-type herbicides to avoid accidental injury to susceptible plants. Also avoid applications under wind conditions that could create drift to nontarget areas.
- Rinse empty containers three times before disposing of them. Add the rinse to the spray tank and dispose of containers according to local regulations to avoid hazard to humans, animals, and the environment.
- Follow label directions for mixing and application to keep residues within the limits prescribed by law.
- Plan ahead. Discuss with your physician the materials you will be using during the season so that he or she can be prepared to provide the appropriate treatment in case of accidental exposure. If symptoms of illness occur, call the physician or get the patient to a hospital immediately. Always provide the medical personnel with as much information as possible.
- Be cautious when you apply pesticides. Know your legal responsibility as a pesticide applicator. You may be liable for injury or damage resulting from pesticide use.

Prepared by Jay W. Pscheidt, Professor and Extension plant pathology specialist, Department of Botany and Plant Pathology; Ed Peachey, assistant professor of weed science, Department of Horticulture; and Vaughn Walton, Associate Professor and Extension Entomologist, Department of Horticulture; all of Oregon State University. The information in this pest management guide is valid for 2014. Trade-name products and services are mentioned as illustrations only. This does not mean that the Oregon State University Extension Service either endorses these products and services or intends to discriminate against products and services not mentioned. Due to constantly changing laws and regulations, the Oregon State University Extension Service can assume no liability for the suggested use of chemicals contained in this guide. Pesticides should be applied according to the label directions on the pesticide container.

Dedicated to the memory of Jeff Olsen, Extension horticulturist, Yamhill County.

© 2014 Oregon State University. This publication was produced and distributed in furtherance of the Acts of Congress of May 8 and June 30, 1914. Extension work is a cooperative program of Oregon State University, the U.S. Department of Agriculture, and Oregon counties. Oregon State University Extension Service offers educational programs, activities, and materials without discrimination based on age, color, disability, gender identity or expression, genetic information, marital status, national origin, race, religion, sex, sexual orientation, or veteran's status. Oregon State University Extension Service is an Equal Opportunity Employer.

Revised March 2014.

Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.