

departments, utility companies, and other agencies that might spray rights-of-way or roadsides know where your vineyard is. If rights-of-way or roadsides run through your property, keep them free of weeds so they are less likely to be sprayed. Encourage your neighbors to use [drift-reduction spray nozzles](#) (nozzles that produce large droplets) and to select herbicides that are less likely to injure grapes.

Minimize drift injury from herbicides used in the vineyard.

Glyphosate is registered for use in nonbearing and bearing grapes as a directed spray. However, if it is not applied properly, severe damage can occur. To avoid injury, grape growers should observe the following guidelines.

- Avoid glyphosate contact with any green parts of the vine or through drift.
- If possible, avoid summer and fall applications (when grapes are most susceptible to injury).
- Avoid glyphosate applications when shoots begin to trail, especially with downward shoot-training systems.
- Use a shield mounted to a wand for a backpack sprayer application or a commercial shielded sprayer such as a dome sprayer.
- Avoid spraying in windy conditions or during temperature inversions.
- Use drift-reduction nozzles (for example, drift-guard or air induction types) that operate at lower pressure (15 to 30 psi) and produce large droplets.
- Use grow tubes to protect the green shoots of first-year vines from herbicide contact.
- If you practice chemical weed control in your vineyard, begin the weed management program with preemergence herbicides (consult the [PNW Weed Management Handbook](#)) and follow up with postemergence herbicides before bud break.
- In midseason, use a contact herbicide (not systemic) to treat weed escapes. Examples are products such as Gramoxone (paraquat) or Rely (glufosinate).
- If you use 2,4-D in your vineyard, apply it before active shoot growth occurs, use low spray pressures, and be extremely careful to avoid treatment when weather conditions favor drift, such as during high temperatures, breezy conditions, and temperature inversions.



Figure 3. Glyphosate injury symptoms: (A) Distorted leaves. (B) Lateral shoot growth with unusual burst of latent buds on nodes, short internodes, and distorted leaves.



Figure 4. ALS inhibitor injury symptoms from sulfonylurea herbicide spray drift: chlorosis of leaf veins and change in leaf appearance from smooth to crinkled.

Other herbicide drift resources

Diagnosing Herbicide Injury on Garden and Landscape Plants. ID-184-W. Purdue Plant & Pest Diagnostic Laboratory. http://www.extension.purdue.edu/extmedia/ID/ID_184_W.pdf

Leaf Index and Severity Rating, Washington State University. <http://feql.wsu.edu/eb/>

Oregon Department of Agriculture (503-986-4653). Direct investigations of suspected Oregon drift incidents here.

Questions and Answers About Vineyard Injury from Herbicide Drift. Kansas State University. <http://www.ksre.ksu.edu/bookstore/pubs/mf2588.pdf>

Washington Pesticide Management Division (Washington Department of Agriculture, 509-225-2647, toll-free 1-877-301-4555). Direct investigations of suspected Washington drift incidents here.

Use pesticides safely!

- Wear protective clothing and safety devices as recommended on the label. Bathe or shower after each use.
- Read the pesticide label—even if you've used the pesticide before. Follow closely the instructions on the label (and any other directions you have).
- Be cautious when you apply pesticides. Know your legal responsibility as a pesticide applicator. You may be liable for injury or damage resulting from pesticide use.

All photos in this publication by Imed Dami, © The Ohio State University

Trade-name products and services are mentioned as illustrations only. This does not mean that the Oregon State University Extension Service either endorses these products and services or intends to discriminate against products and services not mentioned.

© 2014 Oregon State University. This publication was produced and distributed in furtherance of the Acts of Congress of May 8 and June 30, 1914. Extension work is a cooperative program of Oregon State University, the U.S. Department of Agriculture, and Oregon counties. Oregon State University Extension Service offers educational programs, activities, and materials without discrimination based on age, color, disability, gender identity or expression, genetic information, marital status, national origin, race, religion, sex, sexual orientation, or veteran's status. Oregon State University Extension Service is an Equal Opportunity Employer.

Published March 2004. Revised February 2014.